

Chironomids of the Subfamily Diamesinae (Diptera, Chironomidae) from Japan

III. *Boreoheptagyia unica* sp. nov.

By

Eugenyi A. MAKARCHENKO

Institute of Biology and Pedology, Far East Branch of Russian
Academy of Sciences, Vladivostok-22, 690022 Russia

Abstract Male of *Boreoheptagyia unica* sp. nov. is described from the Nikko National Park, Honshu, Japan.

Key words: Diptera; Chironomidae; Diamesinae; *Boreoheptagyia*; taxonomy; morphology; distribution.

Introduction

The genus *Boreoheptagyia* BRUNDIN, 1966, is distributed in the Holarctic and comprises sixteen species (SERRA-TOSIO, 1989). *Boreoheptagyia brevitarsis* (TOKUNAGA), comb. nov., *B. eburnea* (TOKUNAGA), comb. nov., and *B. nipponica* (TOKUNAGA), comb. nov., were described from Japan by TOKUNAGA (1936, 1937, 1939) and were known as species of the *Heptagyia* genus (SASA, 1989). *Boreoheptagyia brevitarsis* is distributed in Japan and in some regions of the Russian Far East (MAKARCHENKO, 1985). For the two other Japanese species, only females are known and they cannot be correctly identified now.

Below I am describing a new species of *Boreoheptagyia*, *B. unica* sp. nov., from the Nikko National Park, Honshu, Japan.

The terminology follows SAETHER (1980). Material was fixed by 70% ethanol.

Boreoheptagyia unica sp. nov.

(Figs. 1–4)

Type locality. Japan, Honshu, Nikko National Park.

Type material. Holotype: ♂, Toyamazawa River, upper stream, Nikko National Park, altitude 1,450 m above sea-level, Okunikko, Tochigi Prefecture, Japan, June 23–24, 1988, leg. R. UENO. Holotype is deposited in the National Science Museum (Nat. Hist.), Tokyo.

Etymology. From Latin unica – unique.

Description.

Male imago. Generally color dark brown. Legs brownish yellow, with dark

colored rings (Fig. 3). Body length 2.75 mm. Wing length 2.5 mm. Total length / wing length 1.1.

Head:— Eyes bare, without dorsomedial extension. Coronals 4, outer verticals 4, inner verticals about 19, frontals 1, postorbitals 6, clypeals 14 (44.8–64 μm). Antenna as in female, with 6 flagellomeres and without plumose (Fig. 1). Flagellomeres length (μm): 48: 25.6: 22.4: 28.8: 32: 76.8. First flagellomere with 3 short setae (38.4 μm), second and third with 5 setae (38–40 μm), fourth with 4 setae (38.4 μm), fifth with 5 setae (48–57.6 μm), sixth with 4 setae (44.8–80 μm). Pedicel 44.8 μm in length and with 5 short setae (19–32 μm). AR=0.49. Last 4 maxillary palpal segments length (μm): 70.4: 131.2: 131.2: 2: 236.8. Head width / palp length: 0.79.

Thorax:— Antepronotum with 1–2 dorsal and 14 ventral antepronotals. Acrostichals 20 (16–41.6 μm), in anterior half only; dorsocentrals 9–10, prealars 17–18; supraalars 1; scutellars about 60. Distribution of setae and color of scutum are shown in Fig. 2.

Wing:— Surface with microtrichia, well visible under $\times 400$. R and R_1 with 52, R_{4+5} with 28 macrotrichia. Squama with 30 setae (56–72.6 μm) in 1–2 rows. Anal lobe a little reduced.

Legs:— With short setae, $BR_{1-3}=1.6$ –1.8. Front tibial spur length 44.8 μm , middle tibial spurs length 51.2 and 48 μm , hind tibial spurs length 70.4–57.6 μm and with comb of 14 spines.


Length (μm) and proportions of legs:

P	fe	ti	ta ₁	ta ₂	ta ₃	ta ₄	ta ₅	LR	SV	BV
P ₁	781	891	562	315	164	55	82	0.63	2.98	3.63
P ₂	1178	1274	740	438	233	82	123	0.58	3.31	3.64
P ₃	808	959	562	343	164	41	82	0.59	3.14	3.69

Hypopygium (Fig. 4):— Tergite IX with small anal point (13.2 μm) and 21 short setae (6.6–10 μm). Laterosternite IX with 10–14 setae (56–85.8 μm). Aedeagal lobe (A1) membranous, with small pores at posterior part. Inferior volsella (IVo) small, with setae in 1–2 rows (36–39.6 μm). Sternapodeme high, trapeziform. Gonocoxite simple, elongate. Gonostylus widest in middle part, surface thickly covered with short spine-like setae; apical part with lobe and two megasetae (16.5–19.8 μm), a seta (13.2 μm) near megaseta present.

Female, pupa and larva are unknown.

Remarks. SERRA-TOSIO (1989) recorded two species of *Boreoheptagyia* with female-like antennae of males. *Boreoheptagyia lurida* (GARRET) from Nearctic has the antenna with 5 flagellomeres and *B. cinctipes* (EDWARDS) has the antenna with 8 flagellomeres. *Boreoheptagyia unica* sp. nov. is evidently different from these species in having 6-flagellomere antenna and features of hypopygium.



Figs. 1–4. Male of *Boreoheptagyia unica* sp. nov.; 1, antenna; 2, scutum; 3, legs; 4, hypopygium (without proctiger); Al – aedeagal lobe, IVo – inferior volsella.

Acknowledgments

I am much indebted to Dr. T. IWAKUMA and R. UENO of the National Institute for

Environmental Studies in Tsukuba, Ibaraki, for making the material available to me. This study was partly supported by the research fund of the Japan Society for the Promotion of Science.

References

- BRUNDIN, L., 1966. Transantarctic relationships and their significance, evidenced by chironomid midges. With a monograph of the subfamilies Podonominae and Aphroteniinae and the austral Heptagyiinae. *K. Svenska Vetensk Akad. Handl.*, 11: 1-472.
- MAKARCHENKO, E. A., 1985. Chironomids of the Soviet Far East. Subfamilies Podonominae, Diamesinae and Prodiamesinae. 208 pp. DVNC AN SSSR Press, Vladivostok. (In Russian.)
- SAETHER, O. A., 1980. Glossary of chironomid morphology terminology (Diptera, Chironomidae). *Ent. scand. Suppl.*, 14: 51 pp.
- SASA, M., 1989. Chironomidae of Japan: checklist of species recorded, key to males and taxonomic notes. *Res. Rept. natn. Inst. env. Stud. Japan*, (125): 1-177.
- SERRA-TOSIO, B., 1989. Révision des espèces ouest-paléarctiques et néarctiques de *Boreoheptagyiina* BRUNDIN avec des clés pour les larves, les nymphes et les imagoes (Diptera, Chironomidae). *Spixiana*, 11: 133-173.
- TOKUNAGA, M., 1936. Chironomidae from Japan. VI. Diamesinae. *Philipp. J. Sci.*, 59: 525-552.
- 1937. Family Chironomidae (1). *Fauna Nipponica*, 10 (7-1): 1-110.
- 1939. Chironomidae from Japan (Diptera). XI. New or little-known midges, with special reference to the metamorphosis of torrential species. *Philipp. J. Sci.*, 69: 297-345.