

Karyotypes of Bagrid Catfishes, *Mystus wyckii* and *Bagroides macracanthus*, from Thailand

By

Wichian MAGTOON

Department of General Science, Faculty of Science, Srinakharinwirot
University at Bangkhen, Bangkok, Thailand

and

Ryoichi ARAI

Department of Zoology, National Science Museum, Tokyo

Abstract Chromosomes of bagrid catfishes, *Mystus wyckii* and *Bagroides macracanthus*, from Thailand, are observed. The karyotype of *Mystus wyckii* comprises 16 metacentric, 14 submetacentric, and 24 acrocentric chromosomes, and that of *Bagroides macracanthus* comprises 16 metacentric, 26 submetacentric, 2 subtelocentric, and 6 acrocentric chromosomes. Karyotypes of 30 species of Asian bagrid catfishes are shown in a table.

Karyotypes of 28 species in the Bagridae have been reported from India (SRIVASTAVA & DAS, 1969; RISHI, 1973; MANNA & PRASAD, 1974; DAS & KAR, 1977; MANNA & KHUDA-BUKHSH, 1978; KHUDA-BUKHSH *et al.*, 1980; TRIPATHY & DAS, 1980; BARAT & KHUDA-BUKHSH, 1986; SHARMA & TRIPATHI, 1986; KHUDA-BUKHSH & BARAT, 1987), China (LING, 1982; SHEN *et al.*, 1983; HONG & ZHOU, 1984; YU *et al.*, 1987), Korea (KIM *et al.*, 1982; LEE *et al.*, 1983; UENO, 1985) and Japan (FUJIOKA, 1973; UENO, 1974, 1985). However, there have been no reports on bagrid karyotypes from Southeast Asia, in spite of the fact that Southeast Asia is a very important area for the study on systematics and zoogeography of the Bagridae.

As we observed chromosomes of two species in the Bagridae, *Mystus wyckii* and *Bagroides macracanthus*, from Thailand, their karyotypes will be reported in the following lines.

Materials and Method

Three specimens of *Mystus wyckii* (BLEEKER), 14.0 to 23.5 cm SL, and three specimens of *Bagroides macracanthus* BLEEKER, 20.0 to 21.5 cm SL, were collected from Nakhon Phanom Province, northeast Thailand and used for chromosome observation. Mitotic metaphase chromosomes of the kidney tissue cells were observed. Method of chromosome preparation is similar to that of OJIMA and KURISHITA (1980).

Table 1. Frequency distributions of diploid chromosome counts in two species of the Bagridae.

Species	2n														Total
	48	49	50	51	52	53	54	55	56	57	58	59	60		
<i>Mystus wyckii</i>					1		16		4		5		3	29	
<i>Bagroides macracanthus</i>	5	1	17	1										24	




Fig. 1. Mitotic metaphase chromosomes and karyotypes of two bagrid catfishes. A and C, *Mystus wyckii*; B and D, *Bagroides macracanthus*. Each scale indicates 5 μ . M, meta-centrics; SM, submetacentrics; ST, subtelocentrics; A, acrocentrics.

Table 2. Karyotypes of the family Bagridae from Asia.

Species	2n	M	SM	ST	A	NF	Locality	Reference	
<i>Bagroides macracanthus</i>	50	16	26	2	6	92	Thailand	This paper	
<i>Coreobagrus brevicorpus</i>	44	22	14	0	8	80	Korea	KIM <i>et al.</i> , 1982	
<i>C. ichikawai</i>	56	18	14	-24-		88	Japan	UENO, 1985	
<i>Leiocassis crassilabris</i>	52	24	14	14	0	90	China	HONG & ZHOU, 1984	
<i>L. longirostris</i>	52	20	16	16	0	88	China	HONG & ZHOU, 1984	
<i>Mystus bleekeri</i>	56	20	14	10	12	90	India	SHARMA & TRIPATHI, 1986	
<i>M. cavasius</i>	58	18	16	10	14	92	India	SHARMA & TRIPATHI, 1986	
"	58	14	26	4	14	98	India	KHUDA-BUKHSH <i>et al.</i> , 1980	
"	58	18	22	0	18	98	India	TRIPATHY & DAS, 1980	
<i>M. corsula</i>	58	22	20	0	16	100	India	BARAT & KHUDA-BUKHSH, 1986	
<i>M. elongatus</i>	60	20	12	16	12	92	China	YU <i>et al.</i> , 1987	
<i>M. gulio</i>	58	30	12	2	14	100	India	MANNA & KHUDA-BUKHSH, 1978	
<i>M. macropterus</i>	60	20	12	16	12	92	China	HONG & ZHOU, 1984	
<i>M. seenghala</i>	54	28	12	8	6	94	India	SHARMA & TRIPATHI, 1986	
"	50	30			20	80	India	SRIVASTAVA & DAS, 1969	
<i>M. tengara</i>	male	54	10	-38-		6	India	} RISHI, 1973	
	female	54	9	-38-		7	India		
<i>M. vittatus</i>		54	22	20	12	0	96	India	SHARMA & TRIPATHI, 1986
"		54	22	26	6	0	102	India	TRIPATHY & DAS, 1980
"	type A	54	20	24	10	0	98	India	} MANNA & PRASAD, 1974
"	type B	58	16	10	20	12	84	India	
"		50					India	SRIVASTAVA & DAS, 1969	
<i>M. wyckii</i>		54	16	14	0	24	84	Thailand	This paper
<i>Pelteobagrus eupogon</i>		50	20	14	16	0	84	China	HONG & ZHOU, 1984
<i>Pe. fulvidraco</i>		52	24	14	10	4	90	China	HONG & ZHOU, 1984
"		52	18	26	- 8-		96	Korea	KIM <i>et al.</i> , 1982
"		52	24	24	0	4	100	Korea	LEE <i>et al.</i> , 1983
"		52	28	12	12	0	92	China	SHEN <i>et al.</i> , 1983
"		52	-22-	24	6	74	China	LING, 1982	
<i>Pe. nitidus</i>		52	20	16	16	0	88	China	HONG & ZHOU, 1984
"		52	22	20	-10-		94	Korea	KIM <i>et al.</i> , 1982
<i>Pe. nudiceps</i>		56	18	12	-26-		86	Japan	UENO, 1985
"		56	-44-		-12-		100	Japan	FUJIOKA, 1973
<i>Pe. vachellii</i>		52	22	16	14	0	90	China	HONG & ZHOU, 1984
"		52	16	20	-16-		88	Korea	UENO, 1985
<i>Pseudobagrus adiposalis</i>		50	20	14	14	2	84	China	YU <i>et al.</i> , 1987
<i>Ps. aurantiacus</i>		48	20	12	16	0	80	Japan	} UENO, 1974
"		56	24	12	-20-		92	Japan	
<i>Ps. emarginatus</i>		52	20	20	-12-		92	Korea	UENO, 1985
<i>Ps. pratti</i>		52	20	14	8	10	86	China	YU <i>et al.</i> , 1987
<i>Ps. tenuis</i>		52	22	16	14	0	90	China	HONG & ZHOU, 1984
<i>Ps. truncatus</i>		52	26	14	12	0	92	China	YU <i>et al.</i> , 1987
<i>Ps. ussuriensis</i>		52	18	16	-18-		86	Korea	KIM <i>et al.</i> , 1982
"		52	24	18	10	0	94	China	HONG and ZHOU, 1984
<i>Ps. sp.</i>		48	20	14	-14-		82	Korea	KIM <i>et al.</i> , 1982
"		48	18	14	-16-		80	Korea	UENO, 1985
<i>Rita chrysea</i>		54	28	20	6	0	102	India	DAS & KAR, 1977
<i>R. rita</i>		54	14	34	6	0	102	India	KHUDA-BUKHSH & BARAT, 1987

Classification of chromosomes is adopted from LEVAN *et al.* (1964). Metacentrics and submetacentrics are described as two-arm chromosomes, and subtelocentrics and acrocentrics as one-arm chromosomes.

Results

Mystus wyckii (Fig. 1 A and C). As shown in Table 1, the diploid chromosome number is 54. The karyotype comprises 16 metacentric, 14 submetacentric, and 24 acrocentric chromosomes. The arm number is 84.

Bagroides macracanthus (Fig. 1 B and D). The diploid chromosome number of this species is 50. The karyotype comprises 16 metacentric, 26 submetacentric, 2 subtelocentric, and 6 acrocentric chromosomes. The arm number is 92.

Discussion

Table 2 shows karyotypes of 30 species in the family Bagridae from Asia. The diploid chromosome number of *Mystus wyckii* is 54, which is common in *Mystus* and also found in *Rita* from India. That of *Bagroides macracanthus* is smaller than those of *Leiocassis*, *Mystus*, and many species of both *Pelteobagrus* and *Pseudobagrus*. The diploid chromosome number of *Mystus* ranges from 50 to 60: $2n=50-54$ from India and Thailand, $2n=56-58$ from India, $2n=60$ from China. The reason why $2n$ of *Mystus* is so variable is not known, but one of reasons may be attributable to a confused status of classification of the genus. Besides, it seems puzzling that $2n=54$ has not been observed in *Pelteobagrus* ($2n=50-56$) and *Pseudobagrus* ($2n=48-56$). To solve such problems, it is necessary to examine karyotypes of more species of the family Bagridae.

Acknowledgment

We wish to thank Mr. Thawat DONSAKUL, Srinakharinwirot University at Bangkok, Bangkok, for his help in this study.

References

- BARAT, A., & A. R. KHUDA-BUKHS, 1986. Karyomorphological studies in two species of fishes, *Lepidocephalichthys guntea* (fam: Cobitidae) and *Mystus corsula* (fam: Bagridae). *Perspectives in Cytology and Genetics*, 5: 115-118.
- DAS, R. K., & R. N. KAR, 1977. Somatic chromosome analysis of a siluroid fish, *Rita chrysea* DAY. *Caryologia*, 30: 247-253.
- FUJIOKA, Y., 1973. A comparative study of the chromosomes in Japanese freshwater fishes. I. *Bull. Fac. Educ., Yamaguchi Univ.*, 23: 191-195. (In Japanese with English summary.)
- HONG, Y., & T. ZHOU, 1984. Karyotypes of nine species of Chinese catfishes (Bagridae). *Zool. Res.*, 5 (3), suppl.: 21-28, 3 pls. (In Chinese with English abstract.)

- KHUDA-BUKHSH, A. R., & A. BARAT, 1987. Chromosomes in fifteen species of Indian teleosts (Pisces). *Caryologia*, **40**: 131-144.
- , S. K. GUPTA & S. GOSWAMI, 1980. Karyotypic studies in *Garra lamta* and *Mystus cavassius* (Pisces). *Proc. Ind. Acad. Sci.*, (Anim. Sci.), **89**: 557-562.
- KIM, D. S., E. H. PARK & J. S. KIM, 1982. Karyotypes of nine species of the Korean catfishes (Teleostomi: Siluriformes). *Korean J. Genet.*, **4**: 57-68.
- LEE, H. Y., C. H. YU, S. K. JEON & H. S. LEE, 1983. The karyotype analysis on 29 species of fresh water fish in Korea. *Bull. Inst. basic Sci., Inha Univ.*, **4**: 79-86, pls. 1-7. (In Korean with English abstract.)
- LEVAN, A., K. FREDGA & A. A. SANDBERG, 1964. Nomenclature of centromeric position on chromosomes. *Hereditas*, **52**: 201-220.
- LING, J., 1982. Studies on the karyotypes of eight species of fish. *J. Wuhan Univ.*, (Nat. Sci.), **1982** (2): 109-112, pls. 2-3. (In Chinese with English abstract.)
- MANNA, G. K., & A. R. KHUDA-BUKHSH, 1978. Karyomorphological studies in three species of teleostean fishes. *Cytologia*, **43**: 69-73.
- & R. PRASAD, 1974. Cytological evidence for two forms of *Mystus vittatus* (BLOCH) as two species. *Nucleus*, **17**: 4-8.
- OJIMA, Y., & A. KURISHITA, 1980. A new method to increase the number of mitotic cells in the kidney tissue for fish chromosome studies. *Proc. Japan Acad.*, **56B**: 610-615.
- RISHI, K. K., 1973. Somatic karyotypes of three teleosts. *Genen Phaenen*, **16**: 101-107.
- SHARMA, O. P., & N. K. TRIPATHI, 1986. Karyotypic diversity in genus *Mystus* (Bagridae: Pisces). *Cytologia*, **51**: 1-9.
- SHEN, J., Z. FAN & G. WANG, 1983. Investigation on karyotype in *Pseudobagrus fulvidraco*. *Hereditas, Beijing*, **5**: 23-24. (In Chinese.)
- SRIVASTAVA, M. D. L., & B. DAS, 1969. Somatic chromosomes of teleostean fish. *J. Hered.*, **60**: 57-58.
- TRIPATHY, N. K., & C. C. DAS, 1980. Chromosomes in three species of Asian catfish. *Copeia*, **1980**: 916-918.
- UENO, K., 1974. Chromosomal polymorphism and variant of isozymes in geographical populations of *Pseudobagrus aurantiacus*, Bagridae. *Jpn. J. Ichthyol.*, **21**: 158-164. (In Japanese with English abstract.)
- 1985. Karyotypes of bagrid catfishes from Japan and Korea. *Kaiyôkagaku, Tokyo*, **17**: 102-108. (In Japanese.)
- YU, X., T. ZHOU, K. LI, Y. LI & M. ZHOU, 1987. On the karyosystematics of cyprinid fishes and a summary of fish chromosome studies in China. *Genetica*, **72**: 225-235.

