

Phenolic Compounds in the Leaves of *Pedicularis chamissonis* in Japan

Yoshinori Murai* and Tsukasa Iwashina

Department of Botany, National Museum of Nature and Science,
Amakubo 4-1-1, Tsukuba, Ibaraki 305-0005, Japan

*E-mail: murai@kahaku.go.jp

(Received 18 May 2015; accepted 24 June 2015)

Abstract We analyzed phenolic compounds in the leaves of *Pedicularis chamissonis* which were obtained from a wide range of Japan. Acteoside and luteolin 7-*O*-glucuronide were found as major compounds in the all plants from various populations. Luteolin 7-*O*-glucoside was additionally found in the plants sampled from in the southern part of their habitats. Fujii *et al.* (2013) divided the plants into two species, i.e., *P. chamissonis* (Northern type) and *P. japonica* (Central Honshu type) based on DNA, morphological and ecological data. Our chemical data seem to support this classification from the view point of chemotaxonomy. In addition, ecological significance of the isolated compounds is also described.

Key words: chemical ecology, chemotaxonomy, flavonoids, *Pedicularis chamissonis*, *Pedicularis japonica*, phenylpropanoid.

Introduction

The genus *Pedicularis* (Orobanchaceae) consists of ca. 600 species that is mainly distributed in mountain area of Central and East Asia (Mabberley, 2008). Of their species, *P. chamissonis* Steven is a perennial herb and widely distributed in subalpine and alpine zone of Japan. The plant shows a variety of morphological features, and the diversity resulting in several intraspecific taxa (Yamazaki, 1993). Fujii *et al.* (1997) have reported that the two major clades of the species, i.e., Central Honshu clade and Northern clade by gene phylogeny using chloroplast DNA. Subsequently, they suggested that the plants should be treated Central Honshu type as *P. japonica* and Northern type as *P. chamissonis* from the evidence obtained of genetic, morphological and ecology data (Fujii *et al.*, 2013).

Phenolic compounds such as flavonoids and phenylpropanoids are known as UV protectants, antioxidants and so on. The intensity of UV radiation markedly increases at high altitude (Shibata

1992; Murai *et al.*, 2009). Plants that occupy the highland area are thought to be adaptive to several environmental stresses (e.g. intense UV-B and cold) by accumulating UV-absorbing compounds such as flavonoids and phenylpropanoids (Spitaler *et al.*, 2006; Murai *et al.*, 2009, 2015). In addition, these compounds are also used as chemical markers of plant species. Some studies have been carried out on phenolics of *Pedicularis* (Yoshitama *et al.*, 1980; Zimin and Zhongjian, 1991; Di *et al.*, 2004; Chu *et al.*, 2007), but the number of related study is limited.

In this study, we analyzed *P. chamissonis* growing in Central Japan to Hokkaido for their phenolic compounds, and describe the chemotaxonomical and ecological significance of their compounds.

Materials and Methods

Plant materials

Pedicularis chamissonis was collected from Nagano to Hokkaido (S1–S8), Japan (Fig. 1),



Fig. 1. Collection sites of *Pedicularis chamissonis*. S1: Mt. Hoken-dake. S2: Mt. Norikura-dake. S3: Mt. Arakawa-naka-dake. S4: Mt. Jizo-dake. S5: Mt. Gassan. S6: Mt. Chokai. S7: Mt. Hayachine. S8: Mt. Shari-dake.

i.e., S1: Mt. Hoken-dake, Kiso Mountains, Nagano Prefecture, S2: Mt. Norikura, Hida Mountains, Nagano Prefecture, S3: Mt. Arakawa-naka-dake, Akaishi Mountains, Shizuoka Prefecture, S4: Mt. Jizo-dake, Akaishi Mountains, Yamanashi Prefecture, S5: Mt. Gassan, Dewa Mountains, Yamagata Prefecture, S6: Mt. Chokai-san, Dewa Mountains, Yamagata Prefecture, S7: Mt. Hayachine, Iwate Prefecture, and S8: Mt. Shari-dake, Hokkaido, in summer of 2011–2014. Plant collections were carried out under the permissions of each prefecture, the Ministry of Environment, the Ministry of Agriculture, Forestry and Fisheries, and the Agency for Cultural Affairs, Japan, and Tokushu Tokai Paper Co., Ltd. and Dewa Sanzan Shrine. Voucher specimens were deposited in the Herbarium of National Museum of Nature and Science, Japan (TNS).

Extraction and separation

Fresh leaves of each individual of *P. chamissonis* (1.5 g) were extracted with MeOH. The concentrated extracts were applied to preparative paper chromatography using solvent systems: BAW (*n*-BuOH/HOAc/H₂O = 4:1:5, upper phase) and 15% HOAc. The compounds were purified by Sephadex LH-20 column chromatography (solvent system: 70% MeOH).

Quantitative HPLC analysis of phenolic compounds

Fresh leaves (0.2 g) from three individuals of each population were extracted with 4 ml MeOH. After filtration with Maisyordisc H-13-5 (Tosoh), the extracts were analyzed using a Shimadzu HPLC system with *L*-column2 ODS (5 μ m particle material, I.D. 6.0 $\times$ 150 mm, Chemicals Evaluation and Research Institute, Japan), at flow-rate: 1.0 mL min⁻¹, detection: 190–400 nm and eluents: MeCN/H₂O/H₃PO₄ (20:80:0.2), injection: 10 μ L.

Liquid chromatograph-mass spectra (LC-MS)

LC-MS were measured with Shimadzu LC-MS system using *L*-column2 ODS (3 μ m particle material, I.D. 2.1 $\times$ 100 mm, Chemicals Evaluation and Research Institute), at a flow-rate of 0.2 mL min⁻¹, eluting with HCOOH/MeCN/H₂O (1:20:79), injection: 3 μ L, ESI⁺ 4.5 kV, ESI⁻ -3.5 kV, 250°C.

Identification of compounds

In this research, one phenylpropanoid, acteoside (**1**) (Fig. 3), and two flavonoids, 7-*O*-glucoside (**2**) and 7-*O*-glucuronide (**3**) of luteolin (Fig. 4), were identified by UV spectroscopy (Mabry *et al.*, 1970), acid hydrolysis, LC-MS and HPLC comparisons with authentic standards from the leaves of *Plantago* spp. (Murai *et al.*, 2008, 2009). UV, acid hydrolysis, HPLC and LC-MS data of the isolated compounds are as follows.

Acteoside (verbascoside) (**1**). White powder. UV: λ_{max} (nm) MeOH 292, 330; + NaOMe 380; + AlCl₃ 300, 361; + AlCl₃/HCl 292, 330; + NaOAc 290, 340; + NaOAc/H₃BO₃ 295, 355.



Fig. 2. *Pedicularis chamissonis* in Mt. Norikura, Nagano Prefecture (left) and Mt. Chokai, Yamagata Prefecture (right).




Fig. 3. Chemical structure of acteoside from *Pedicularis chamissonis*.




Fig. 4. Chemical structures of flavones from *Pedicularis chamissonis*. **2**: luteolin 7-*O*-glucoside (R = glucosyl), **3**: luteolin 7-*O*-glucuronide (R = glucuronyl).

HPLC: Rt 7.4 min. LC-MS: m/z 647 $[M + Na]^+$, 623 $[M - H]^-$.

Luteolin 7-*O*-glucoside (**2**). Pale yellow powder. UV: λ_{max} (nm) MeOH 255, 267, 343; + NaOMe 267, 385 (inc.); + $AlCl_3$ 273, 426; + $AlCl_3/HCl$ 262sh, 276, 297 362, 392sh; + NaOAc 260, 400; + NaOAc/ H_3BO_3 260, 373. HPLC: Rt 8.0 min. Acid hydrolysis: luteolin and glucose. LC-MS: m/z 449 $[M + H]^+$, 447 $[M - H]^-$ (luteolin + 1 mol glucose), and 287 $[M - 162 + H]^+$, 285 $[M - 162 - H]^-$ (luteolin).

Luteolin 7-*O*-glucuronide (**3**). Pale yellow powder. UV: λ_{max} (nm) MeOH 256, 268, 348; + NaOMe 266, 389 (inc.); + $AlCl_3$ 274, 296sh, 427; + $AlCl_3/HCl$ 263sh, 272, 295, 364, 386sh; + NaOAc 260, 401; + NaOAc/ H_3BO_3 260, 373. HPLC: Rt 8.6 min. Acid hydrolysis: luteolin and glucuronic acid. LC-MS: m/z 463 $[M + H]^+$, 461 $[M - H]^+$ (luteolin + 1 mol glucuronic acid), and 287 $[M - 176 + H]^+$, 285 $[M - 176 - H]^-$ (luteolin).




Fig. 5. HPLC patterns of MeOH extracts from *Pedicularis chamissonis*. (a) Central Honshu type. (b) Northern type. 1: acteoside, 2: luteolin 7-*O*-glucoside, 3: luteolin 7-*O*-glucuronide.

Results and Discussion

The composition of phenolic compounds

One phenylpropanoid, acteoside (1) and two flavones, luteolin 7-*O*-glucoside (2) and 7-*O*-glucuronide (3) were isolated from the leaves of *P. chamissonis*. However, the presence or absence of 2 is different between the northern and southern populations (Fig. 5; Table 1). Compound 2 was exclusively produced by the plants sampled from S1 (Mt. Hoken-dake) to S5 (Mt.

Gassan), while the other populations locating in the more northern part of Japan lacked this compound. The other peaks on HPLC chromatograms (Fig. 5) were presumed as other phenylpropanoids and flavonoids from their UV spectra obtained by photo diode array HPLC detector, and they were not identified in this study. The concentration of these minor compounds was varied in low level among populations.

Table 1. Phenolic compounds and DNA type of *Pedicularis chamissonis*

Population	Phenolic compounds			DNA type*
	1	2	3	
Mt. Hoken-dake (S1)	+	+	+	CHT
Mt. Norikura-dake (S2)	+	+	+	CHT
Mt. Arakawa-naka-dake (S3)	+	+	+	CHT
Mt. Jizo-dake (S4)	+	+	+	CHT
Mt. Gassan (S5)	+	+	+	CHT/NT
Mt. Chokai (S6)	+	—	+	CHT/NT
Mt. Hayachine (S7)	+	—	+	NT
Mt. Shari-dake (S8)	+	—	+	NT

1: acteoside, 2: luteolin 7-*O*-glucoside, 3: luteolin 7-*O*-glucuronide. + : presence. — : absence.

*According to Fujii *et al.* (2013). CHT: Central Honshu type, NT: Northern type.

Flavonoids and phylogeny of *Pedicularis chamissonis*

Flavonoids are more stable chemotaxonomic marker than phenylpropanoids among phenolic compounds (Sahin *et al.*, 2007). Two chemotypes (presense or absence of **2**) were found in *P. chamissonis* of Japan. These chemotypes seem to support the classification by Fujii *et al.* (2013) that divided *P. chamissonis* in Japan into *P. japonica* (Central Honshu type) and *P. chamissonis* (Northern type). According to Fujii *et al.* (2013), two species, *P. japonica* and *P. chamissonis* are both distributed in Mt. Gassan and Mt. Chokai. In the present study, we collected the plants from their large population in each mountain and another chemotype (species) was not detected in HPLC analysis, but some plants at the other populations in those mountains might possess the opposite chemotype.

Ecological significance

In this study, acteoside and luteolin glycosides were isolated from *Pedicularis chamissonis*. Both phenolics are *ortho*-dihydroxylated phenolics that are strong antioxidants among related compounds (Rice-Evans *et al.*, 1996; Pietta, 2000; Nishibe, 2002). Interestingly, several plants growing in higher altitudes synthesize such *ortho*-dihydroxylated phenolics (Spitaler *et al.*, 2006; Murai and Iwashina, 2010; Murai *et al.*, 2009, 2015). These compounds may play an important role in protection against several environmental stresses (e.g. intense UV light and

cold) in high altitudes. Further studies, e.g. physiological experiment, using alpine plants are required for the understanding of their functions.

Acknowledgments

This work was partially supported by JSPS KAKENHI Grant No. 24570035 (Representative, T. Iwashina).

References

- Chu, H.-B., Tan, N.-H. and Zhang, Y.-M. 2007. Chemical constituents from *Pedicularis rex* C. B. Clarke. *Zeitschrift für Naturforschung* 62b: 1465–1470.
- Di, D.-L., Chen, J. and Shi Y.-P. 2004. Determination of phenylpropanoid glycosides in Chinese herbal extracts from *Pedicularis* species by HPLC. *Journal of Liquid Chromatography and Related Technologies* 27: 2235–2245.
- Fujii, N., Ueda, K., Watano, Y. and Shimizu, T. 1997. Intraspecific sequence variation of chloroplast DNA in *Pedicularis chamissonis* Steven (Scrophulariaceae) and geographic structuring of the Japanese “Alpine” plants. *Journal of Plant Research* 110: 195–207.
- Fujii, N., Ueda, K., Watano, Y. and Shimizu, T. 2013. Taxonomic revival of *Pedicularis japonica* from *P. chamissonis* (Orobanchaceae). *Acta Phytotaxonomica et Geobotanica* 63: 87–97.
- Mabberley, D. J. 2008. *Mabberley's Plant-book*, third ed. 143 pp. Cambridge University Press, Cambridge.
- Mabry, T. J., Markham, K. R. and Thomas, M. B. 1970. *The Systematic Identification of Flavonoids*. Springer, Berlin.
- Murai, Y. and Iwashina, T. 2010. Flavonol glucuronides from *Geum calthifolium* var. *nipponicum* and *Sieversia pentapetala* (Rosaceae). *Biochemical Systematics and*

- Ecology 38: 1081–1082.
- Murai, Y., Kanemoto, T. and Iwashina, T. 2008. Flavone glucuronides from *Plantago hakusanensis* endemic to Japan. *Biochemical Systematics and Ecology* 36: 815–816.
- Murai, Y., Setoguchi, H., Kitajima, J. and Iwashina, T. 2015. Altitudinal variation of flavonoid content in the leaves of *Fallopia japonica* and the needles of *Larix kaempferi* on Mt. Fuji. *Natural Product Communications* 10: 407–411.
- Murai, Y., Takemura, S., Takeda, K., Kitajima, J. and Iwashina, T. 2009. Altitudinal variation of UV-absorbing compounds in *Plantago asiatica*. *Biochemical Systematics and Ecology* 37: 378–384.
- Nishibe, S. 2002. The plant origins of herbal medicines and their quality evaluation. *Yakugaku Zasshi* 122, 363–379.
- Pietta, P.-G. 2000. Flavonoids as antioxidants. *Journal of Natural Products* 63: 1035–1042.
- Rice-Evans, C. A., Miller, N. J. and Paganga, G. 1996. Structure-antioxidant activity relationships and phenolic acids. *Free Radical Biology and Medicine* 20: 933–956.
- Sahin, F. P., Yamashita, H., Guo, Y., Terasaka, K., Kondo, T., Yamamoto, Y., Shimada, H., Fujita, M., Kawasaki, T., Sakai, E., Tanaka, T., Goda, Y. and Mizukami, H. 2007. DNA authentication of *Plantago* herb based on nucleotide sequences of 18S–28S rRNA internal transcribed spacer region. *Biological and Pharmaceutical Bulletin* 30: 1265–1270.
- Shibata, O. 1992. *Puranta* 22. pp. 12–17 (in Japanese).
- Spitaler, R., Schlorhauser, P. D., Ellmerer, E. P., Merfort, I., Bortenschlager, S., Stuppner, H. and Zidorn, C. 2006. Altitudinal variation of secondary metabolite profiles in flowering heads of *Arnica montana* cv. ARBO. *Phytochemistry* 67: 409–417.
- Yamazaki, T. 1993. *Pedicularis*. In: Iwatsuki, K., Yamazaki, T., Boufford, D. E. and Ohba, H. (eds.), *Flora of Japan*, vol. 3a. Kodansha, Tokyo. pp. 364–371.
- Yoshitama, K., Ishii, K. and Yasuda, H. 1980. A chromatographic survey of anthocyanins in the flora of Japan. *Journal of the Faculty of Science, Shinshu University* 15: 19–26.
- Zimin, L. and Zhongjian, J. 1991. Phenylpropanoid and iridoid glycosides from *Pedicularis striata*. *Phytochemistry* 30: 1341–1344.