

Silurian Favositids (Coelenterata: Tabulata) from the Gionyama Formation, Miyazaki Prefecture, Japan

Shuji Niko¹ and Tomio Adachi²

¹ Department of Environmental Studies, Faculty of Integrated Arts and Sciences, Hiroshima University,
1–7–1 Kagamiyama, Higashihiroshima, Hiroshima 739–8521, Japan

E-mail: niko@hiroshima-u.ac.jp

² 3–2–26 Kadokawa Nishisakae, Higashiusuki, Miyazaki 889–0622, Japan

E-mail: t_adachi@taupe.plala.or.jp

Abstract Five species of favositid tabulate corals are described from the Silurian Gionyama Formation in the Kuraoka area, Miyazaki Prefecture, southern Japan. They are *Favosites aokii* Sugiyama, 1940, *Favosites* sp. cf. *F. forbesi* Milne-Edwards and Haime, 1851, *Favosites* sp. cf. *F. favosiformis* Sokolov, 1951, *Thecia ichikawai* sp. nov., and *Angopora hisingeri* (Jones, 1930). Except for *F.* sp. cf. *F. forbesi* that occurs in the upper Wenlock (upper Lower Silurian) G2 and lower Ludlow (lower Upper Silurian) G3 Members, *F. aokii*, *F.* sp. cf. *F. forbesi* and *T. ichikawai* are only known from the G2 Member and the occurrence of *A. hisingeri* is restricted in the G3 Member. The most characteristic respects of *T. ichikawai* are its relatively large corallite diameters and closely to very closely spaced tabulae. The Gionyama specimens of *Angopora* represent the first record of the genus in Japan.

Key words: Wenlock to Ludlow (late Early to early Late Silurian), *Favosites*, *Thecia*, *Angopora*, Gionyama Formation

Introduction

This paper is a new instalment of the serial papers (Niko, 1998; Niko and Adachi, 1999a and b, 2000, 2002, 2004, 2008) on the tabulate coral fauna from the Silurian Gionyama Formation of the Kuraoka area, Miyazaki Prefecture, southern Japan. During our long termed project, important geologic knowledge of the Gionyama Formation was added by Kido and Sugiyama (2007) and Kido (2009), in which papers Hamada's (1959) subdivision of the formation into the G1 to G4 Members was abandoned and the Lower, Middle and Upper Members were newly proposed. Among them, the Lower and Upper Members can be respectively correlated with the G1 and G4 Members and the Middle Member is equivalent to the incorporated stratum of the G2 and G3 Members. Because our idea concerning the stratigraphy of the formation will publish elsewhere, we still use the previous nomenclature.

Total 106 specimens of favositid tabulate corals are examined in this paper. They are collected from three localities on the western flank of Mt. Gionyama (see Niko, 1998, for their geologic and geographic settings) and kept in National Museum of Nature and Sciences, Tokyo (prefixed NMNS).

Systematic Paleontology

Subclass Tabulata Milne-Edwards and Haime, 1850

Order Favositida Wedekind, 1937

Suborder Favositina Wedekind, 1937

Superfamily Favositoidea Dana, 1846

Family Favositidae Dana, 1846

Subfamily Favositinae Dana, 1846

Genus *Favosites* Lamarck, 1816

Type species: Favosites gothlandica Lamarck, 1816.

***Favosites aokii* Sugiyama, 1940**

(Figs. 1-1-5; 2-3, 4, 6)

Favosites asper [sic] *aokii* Sugiyama, 1940, p. 127, 128, pl. 22, figs. 1-5; Hamada, 1983, fig. 7.*Favosites aokii* Sugiyama; Hamada, 1982, p. 6, pl. 3, fig. 17.**Material examined:** Thirty-five coralla, NMNS PA17356-17390.

Description: Coralla massive with low bulbous growth form, cerioid; maximum sizes of fragmentary coralla indicate 43 mm in height and 108 mm in diameter (NMNS PA17369) and 93 mm in height and 88 mm in diameter (NMNS PA17385). Corallites prismatic have 3-8 sides in transverse section and relatively uniform in diameter ranging 0.4-1.9 mm, usually 1.2-1.6 mm in adult portion; calices very shallow and perpendicularly oriented to corallum surface. Thickness of intercorallite walls is variable even in a same corallum, ranges from thin to relatively thick, 0.03-0.21 mm; structurally they are differentiated into median line and darker stereoplastic layer; walls usually straight, but weak to strong waves frequently developed in thickened portions; mural pores commonly occur on corallite faces as mid-wall pores and less frequently occur near corallite angles; in addition, corner pores at corallite angles are rarely developed; usual diameters of pores are 0.13-0.19 mm; some pores closed by pore plates; septal spines rare to almost absent in thin and well-developed in thick portions of walls; high conical to needle-like, 0.08-0.20 mm in length; tabulae almost complete with somewhat variable profiles indicating nearly flat, concave proximally (sagging), concave proximally (uparched) or slightly wavy; spacing of tabulae is variable, ranging from sparse to close; there are 3-15 tabulae in 5 mm of corallite length.

Occurrence: Abundant in limestone pebbles to boulders of the upper Wenlock (upper Lower Silurian) G2 Member at locality 1.

Discussion: This species was described originally by Sugiyama (1940) as a subspecies of "*Favosites*" *aspera* d'Orbigny (1850, p. 49; = *Paleofavosites aspera* in up to date taxonomy as indicated by Twenhofel, 1914, Stel and Oekentorp, 1976, and Young and Elias, 1995), Hamada (1982) gave it independent species rank. We follow this subsequent determination because the most common mural pores in *Favosites aokii* situate on the corallite faces as the mid-wall pores, whereas "*F.*" *aspera* characteristically has the corner pores. The type stratum of *F. aokii* is the Wenlock to Ludlow limestone of the Kawauchi Formation in the Southern Kitakami Terrane, Northeast Japan.

As can be seen in figures, the Gionyama specimens show some variations in intercorallite wall thickness and spacing of the tabulae. Among ranges of their morphologies, thin to weakly thickened portions of the intercorallite walls (Figs. 1-1, 3, 4) and the moderately spaced tabulae (approximately 7-11 in 5 mm; Fig. 1-2) are quite identical with the holotype.

Favosites* sp. cf. *F. forbesi

Milne-Edwards and Haime, 1851


(Figs. 3-1-6)








Compare:*Favosites forbesi* Milne-Edwards and Haime, 1851, p. 238.

See synonymy in Stel (1978), and added:

Favosites forbesi Milne-Edwards and Haime; Chernyshev, 1937a, p. 9, 10, pl. 1, figs. 3a, b, pl. 4, figs. 6a, b; Shrock and Twenhofel, 1939, p. 254; Yang, 1948, p. 133-135, pl. 1, figs. 1a, b, 2a, b, 3a, b, 4a, b; Sokolov, 1949, p. 79, pl. 6, figs. 3, 4; Chernyshev, 1951, p. 21-23, pl. 2, figs. 1-5; Yü, 1956, p. 604, 615-617, pl. 1, figs. 1, 2; Yü, 1962, p. 51, pl. 12, figs. 1a, b, 2 a-c,

Fig. 1. *Favosites aokii* Sugiyama, 1940, thin sections. 1, 2, 4, NMNS PA17365. 1, transverse sections of corallites, $\times 10$. 2, longitudinal sections of corallites, $\times 10$. 4, partial enlargement of Fig. 1-1, showing details of mural pores and septal spines in thin walled corallites, $\times 14$. 3, NMNS PA17361, transverse sections of corallites, note intra-colonial variations of intercorallite walls thickness, $\times 10$. 5, NMNS PA17356, partial enlargement with the same specimen of Fig. 2-6, showing details of mural pores and septal spines in thick walled corallites, transverse sections, $\times 14$.





- pl. 14, figs. 1a, b, pl. 15, figs. 3 a–c; Yü *et al.*, 1963, p. 221, pl. 70, figs. 2a, b; King, 1974, p. 196, 197, pl. 38, figs. 1a, b; Lee and Noble, 1988, p. 32, 33, 36–38, figs. 3–1–21; 6–1–8.
- Favosites* aff. *forbesi* Milne-Edwards and Haime; Yabe and Hayasaka, 1915, p. 67, 68.
- Favosites forbesi kweitungensis* Grabau, 1930, p. 223–227, pl. 1, figs. 1–3, pl. 2, figs. 4, 5; Tsin, 1956, p. 627, pl. 2, figs. 3–6 [as *F. kweitingensis*]; Yü *et al.*, 1963, p. 221, 222, pl. 70, figs. 3a, b [as *F. kweitingensis*]; Yang in Yang *et al.*, 1978, p. 173, pl. 61, figs. 4a, b [as *F. kweitingensis*].
- Favosites* cf. *forbesi* Milne-Edwards and Haime; Ozaki in Shimizu *et al.*, 1934, p. 70, pl. 13, fig. 1.
- Favosites forbesi mammilatus* Chernyshev, 1936, p. 41, pl. 1, figs. 4, 5, 10.
- Favosites forbesi tuvaensis* Chernyshev, 1937a, p. 10, 11, pl. 1, figs. 4a, b; Keller, 1966, p. 122, 123, pl. 10, figs. 1a, b.
- Favosites forbesi multiperforata* Chernyshev, 1937b, p. 73, 74, text-fig. 2, pl. 2, figs. 2a, b; Barskaya and Sharikova, 1963, p. 142, 143, pl. 19, figs. 3, 4.
- Favosites (Eufavosites) forbesi* Milne-Edwards and Haime; Rukhin, 1937, p. 38–47, pl. 6, figs. 3–7, pl. 7, figs. 1, 2; Rukhin, 1938, p. 47, 48, pl. 8, figs. 1, 2, 7.
- Favosites (Eufavosites) forbesi culi* Rukhin, 1937, p. 47, 48, pl. 8, figs. 1, 2; Rukhin, 1938, p. 50, 51, pl. 9, figs. 10, 11.
- Favosites (Eufavosites) forbesi lenius* Rukhin, 1937, p. 48, pl. 8, figs. 3, 4.
- Favosites (Eufavosites) forbesi kolimaensis* Rukhin, 1938, p. 48, 49, pl. 8, figs. 4–6.
- Favosites (Eufavosites) forbesi mammilatus* Chernyshev; Rukhin, 1938, p. 49, pl. 9, figs. 1, 2.
- Favosites (Eufavosites) forbesi oblonga* Rukhin, 1938, p. 49, 50, pl. 9, figs. 5–7.
- Favosites (Eufavosites) forbesi kana* Rukhin, 1938, p. 50, pl. 8, figs. 8, 9, pl. 9, figs. 3, 8, pl. 12, fig. 7, pl. 13, fig. 1.
- Favosites forbesi muralis* Yang, 1948, p. 135, pl. 1, figs. 5a, b; Yü *et al.*, 1963, p. 222, pl. 70, figs. 5a, b.
- [non] *Favosites forbesi similis* Sokolov, 1952, p. 49, 50, pl. 18, figs. 3, 4 [= *F. similis* Sokolov, see Klaamann, 1962, p. 36].
- [non] *Favosites forbesi sugiyamai* Kamei, 1955, p. 49, 50, pl. 2, figs. 1 a–f, pl. 4, figs. 1, 2 [= *Squameofavosites sugiyamai* (Kamei), see Niko, 2006, p. 24].
- [non] *Favosites forbesi takarensis* Kamei, 1955, p. 50–52, pl. 2, figs. 3 a–c, 4a, b, 5a, b, pl. 4, figs. 4–6 [= *Squameopora takarensis* (Kamei), see Niko, 2006, p. 21].
- Favosites forbesi tankwanpei* Tsin, 1956, p. 626, 627, pl. 2, figs. 2 a–c.
- Favosites forbesi multiporoides* Yü, 1956, p. 604, 605, 617, 618, pl. 1, figs. 3, 4; Yü *et al.*, 1963, p. 222, 223, pl. 70, figs. 4a, b.
- Favosites forbesi qilianshanensis* Yü, 1962, p. 51, 52, pl. 13, figs. 1a, b, pl. 14, figs. 3a, b.
- Favosites forbesi dongkalaensis* Yang in Yang *et al.*, 1978, p. 172, pl. 61, figs. 1a, b.
- Favosites forbesi fenggangensis* Yang in Yang *et al.*, 1978, p. 172, pl. 61, figs. 2a, b.
- Material examined:** Thirty-nine coralla, NMNS PA17312–17350.
- Description:** Coralla massive with somewhat variable growth forms indicating thick tabular, discoid to bulbous, cerioid; boss-like projections on upper surface of coralla are rarely developed; maximum sizes of fragmentary coralla indicate 33 mm in height and 82 mm in diameter (NMNS PA17339) and 104 mm in height and 81 mm in diameter (NMNS PA17341). Corallites prismatic have 3–8 (rarely up to indistinct 10) sides in transverse section; diameters of corallites are relatively large for the genus, usually 1.8–2.3 mm in adult portion; adult corallites surrounded by immature ones. Intercorallite walls may be weakly thickened; apparent mural pore and septal spine are not observable in examined material; tabulae mostly complete with roughly flat profiles; spacing of tabulae is variable, ranging from sparse to very close; counts of tabulae in 5 mm length of corallites are 2–19.
- Occurrence:** Abundant in limestone pebbles to boulders of the G2 Member at locality 1 (NMNS PA17312–17344); rare in massive limestone of the lower Ludlow (lower Upper Silurian) G3 Member at locality 2 (NMNS PA17349, 17350) and locality 3 (NMNS PA17345–17348).

Fig. 2. **1, 2, 5, 7**, *Favosites* sp. cf. *F. favosiformis* Sokolov, 1951, thin sections. **1, 7**, NMNS PA17352. **1**, transverse sections of corallites, $\times 5$. **7**, longitudinal sections of corallites, $\times 5$. **2, 5**, NMNS PA17354, transverse sections of corallites, $\times 10$. **3, 4, 6**, *Favosites aokii* Sugiyama, 1940, thin sections. **3**, NMNS PA17371, longitudinal sections of corallites, note intra-colonial variations of intercorallite walls thickness, $\times 10$. **4**, NMNS PA17387, longitudinal sections of corallites, $\times 10$. **6**, NMNS PA17356, transverse sections of thick walled portions of corallites, $\times 10$.

Discussion: On the basis of their corallite diameters and arrangement of the mature and immature corallites, the Gionyama specimens compare favorably with *Favosites forbesi* Milne-Edwards and Haime, 1851, that widely known from the Silurian strata. Without information about the mural pores and septal spines, a reliable specific identification of the specimens is impossible.

***Favosites* sp. cf. *F. favosiformis* Sokolov, 1951**

(Figs. 2-1, 2, 5, 7)

Compare:

Favosites favosiformis Sokolov, 1951, p. 86, 87, pl. 33, figs. 3, 4; Klaamann, 1964, p. 64, 65, pl. 18, figs. 3–6; Stasinska, 1967, p. 79, pl. 21, figs. 3a, b.

Favosites favosiformis globosa Sokolov, 1951, p. 88–90, pl. 35, figs. 1–3.

Material examined: Five coralla, NMNS PA17351–17355.

Description: Coralla massive with high bulbous to subspherical growth forms, cerioid; largest specimen of fragmentally corallum (NMNS PA17351) indicates 34 mm in height and 38 mm in diameter. Corallites prismatic, large for the genus with 2.8–4.2 mm in usual diameters of adult portion. Intercorallite walls thin having 0.08–0.17 mm; mural pores common, occur on corallite faces; diameters of pores are 0.23–0.29 mm; septal spines sporadic, short conical; tabulae complete with roughly flat profiles; spacing of tabulae moderate; there are 2–5 tabulae in 5 mm of corallite length.

Occurrence: Rare in limestone pebbles of the G2 Member at locality 1.

Discussion: The specimens here included in *Favosites* sp. cf. *F. favosiformis* Sokolov, 1951, resemble the types from the upper Llandovery (lower Lower Silurian) of Estonia (Sokolov, 1951; Klaamann, 1964). The most diagnostic features of *F. favosiformis* are the large corallite diameters ranging from 3.0 to 4.5 mm, the thin intercorallite walls ranging 0.10 to 0.15 mm, large diameters of the mural pores attaining 0.4 mm and the short septal spines. Except for

their smaller diameters of the mural pores, dimensions of the present specimens nearly fall within the range of the specific concept. *Favosites favosiformis* was also recorded by Stasinska (1967) from the Wenlock of Norway and erratic boulders in Poland.

Family Theciidae Milne-Edwards and Haime, 1850

Discussion: Two different opinions have been proposed for a higher taxonomic position of the family Theciidae, namely Sokolov (1950) classified this family into the order Favositida, and on the other hand it was assigned to the order Sarcinulida by Hill (1981). We followed a Hill's idea in provisional report (Adachi and Niko, 1996). However, this family belongs herein to the Favositida based on similarities of intercorallite wall structures in the proximal corallites of the present two species, *Thecia ichikawai* sp. nov. and *Angopora hisingeri* (Jones, 1930), and the usual ones of *Favosites*.

Genus *Thecia* Milne-Edwards and Haime, 1849

Type species: *Porites expatiata* Lonsdale, 1839.

***Thecia ichikawai* sp. nov.**

(Fig. 4-1–7; 5-1, 2, 4)

Thecia sp., Adachi and Niko, 1996, p. 68, figs. 2-1–3.

Holotype: NMNS PA17400, from which five thin sections were made.

Paratypes: Fifteen thin sections were studied from the eight paratypes, NMNS PA17393–17396, 17399, 17402, 17408, 17409.

Other material examined: NMNS PA17397, 17398, 17401, 17403–17407, 17410–17417.

Diagnosis: Species of *Thecia* with encrusting coralla composed of multiple layers of lamellae and relatively large corallite diameters attaining 2.0 mm; thickened intercorallite walls attain approximately 0.5 mm; septal spines long to





Fig. 3. *Favosites* sp. cf. *F. forbesi* Milne-Edwards and Haime, 1851, thin sections. 1–3, NMNS PA17318. 1, transverse sections of corallites, $\times 10$. 2, transverse sections of corallites, $\times 14$. 3, longitudinal sections of corallites, $\times 10$. 4, 5, NMNS PA17315. 4, longitudinal sections of corallites, $\times 5$. 5, transverse sections of corallites, $\times 14$. 6, NMNS PA17312, longitudinal sections of corallites, $\times 5$.



exceptionally long with 0.14–0.59 mm; tabulae indicate close to very close in spacing; 11–21 tabulae in 5 mm.

Description: Coralla encrusting, tabular in total growth form and composed of multiple layers of relatively thick lamellae; each lamella has approximately 3–20 mm in thickness, mostly cerioid; largest fragment of corallum consisting two lamellae with intra-lamella lacuna (paratype, NMNS PA17396) attains 43 mm in height and 80 mm in diameter. Corallites sub-prismatic in proximal and prismatic in distal portions; in basal part of lamellae, proximal corallites prostrate and alveolitoid-like, then they indicate upward bent to form erect distal portions; transverse sections of corallites are sub-trapezoidal to fan-shaped in proximal and polygonal with 5–7 sides in distal portions of corallites; diameters of corallites are relatively large for the genus and almost uniform in each lamella, range from 1.0 to 2.0 mm; calices nearly perpendicular to surface of lamella; no apparent increase of new corallite is observable. Intercorallite walls always thin in proximal corallites at basal part of lamellae; in addition, thin walled portions rarely occur in distal ones near periphery; structurally thin walls differentiated into median dark line and stereoplasma; usual walls thickened by microgranular (?) stereoplasma and septal spines; partial confluence of septal spines exhibits astreoid-to thamnasterioid-like appearances; thus thickness of intercorallite walls variable, 0.08–0.13 mm in thin walled portions and thickened attaining approximately 0.5 mm in thamnasterioid-like portions; mural pores well-developed, have circular profiles, commonly occur on corallite faces and rarely at corallite angles; diameters of typical mural pores are 0.29–0.44 mm; septal

spines needle-like to high conical, long to exceptionally long, and variable in occurrence, range almost absent to numerous; length of protrude portions of septal spines are 0.14–0.59 mm; tabulae complete with horizontal to weakly concave proximally or incomplete in relatively rare cases; spacing of tabulae close to very close; there are 11–21 tabulae in 5 mm of corallite length.

Etymology: The specific name honors Dr. Koichiro Ichikawa, in recognition of his geological contributions of the Kurosegawa Terrane including the Palaeozoic strata of the Kuraoka area.

Occurrence: Abundant in limestone pebbles to boulders of the G2 Member at locality 1.

Discussion: *Thecia ichikawai* sp. nov. is most similar to *T. expatiata* (Lonsdale, 1839, p. 687, pl. 15, figs. 3, 3a; Oekentorp, 1970, p. 148, 149, 151, pl. 12, figs. 1, 2; Hill, 1981, figs. 351-2a-c) from the Wenlock strata in Great Britain, but the species differs by having larger corallite diameters and more closely spaced tabulae. Corallite diameters of the new species attain 2.0 mm, whereas those of *T. expatiata* range from 0.8 to 1.1 mm.

Genus *Angopora* Jones, 1936

Type species: *Laminopora hisingeri* Jones, 1930.





Angopora hisingeri (Jones, 1930)

(Figs. 5-3, 5, 6)

Laminopora hisingeri Jones, 1930, p. 35.

Angopora hisingeri (Jones); Jones, 1936, p. 18, 19, pl. 2, figs. 4–7, pl. 3, figs. 1, 2; Sokolov, 1955, pl. 2, fig. 4, pl. 27, figs. 3–6; Stasinska, 1967, p. 61, 62, pl. 10, figs. 1–3; Klaamann, 1970, p. 62–64, 65, 66, text-figs. 2-A–C, pl. 1, figs. 1–3, pl. 2, figs. 1–4, pl. 3, figs. 1, 2, pl.

Fig. 4. *Thecia ichikawai* sp. nov. thin sections. **1, 2**, paratype, NMS PA17393. **1**, transverse sections of corallites, note intra-colonial variations of intercorallite walls thickness and distribution of septal spines, $\times 10$. **2**, longitudinal sections of corallites, $\times 10$. **3–5**, holotype, NMNS PA17400. **3**, longitudinal section of corallum, $\times 5$. **4**, partial enlargement of Fig. 4-3, showing details of intercorallite wall structure and septal spines, $\times 10$. **5**, transverse sections of thick walled corallites, $\times 10$. **6**, paratype, NMNS PA17394, transverse sections of thick walled corallites, showing details of intercorallite wall structure and septal spines, $\times 14$. **7**, paratype, NMNS PA17399, transverse sections of corallites, showing details of intercorallite wall structure and septal spines, $\times 14$.



4, figs. 1–4; Wang, 1981, p. 40, pl. 24, figs. 4a, b; Scrutton, 1997, text-figs. 13-A–D.

Thecia hisingeri (Jones); Hill and Stumm, 1956, text-figs. 351-1c, d.

Thecia (Angopora) hisingeri (Jones); Hill, 1981, text-figs. 351-1a, b.

Material examined: Two coralla, NMNS PA17391, 17392.

Description: Coralla tabular, mostly cerioid; largest fragment of corallum (NMNS PA17392) attains 19 mm in height and 64 mm in diameter. Proximal corallites in basal part of corallum are prostrate and alveolitoid-like; transverse sections of proximal corallites are sub-trapezoidal to fan-shaped; more distal corallites erect, sub-polygonal having indistinctly 4–8 sided transverse sections; corallite diameters 0.7–1.5 mm; calices normal to corallum surface; increases of new corallites are probably lateral. Intercorallite walls thin in proximal corallites, where they are structurally differentiated into median dark line and stereoplasm; walls of distal corallites are thickened by microgranular (?) stereoplasma and septal spines; thickness of intercorallite walls ranges from 0.10 to 0.38 mm; mural pores abundantly occur on corallite faces and at corallite angles; diameters of typical pores having circular profiles are 0.13–0.21 mm; septal spines absent in proximal portions of corallites, but needle-like spines numerous occur in distal ones; length of protrude portions of septal spines are long, 0.12–0.29 mm; complete tabulae most common, but incomplete ones frequently developed; spacing of tabulae very close; there are approximately 7–13 tabulae in 2 mm of corallite length.

Occurrence: Rare in massive limestone of the G3 Member at locality 2.

Discussion: The general corallite morphologies, intercorallite walls, septal spines and tabulae of the Gioniyama specimens are consistent with those of the types from the Wenlock of Got-

land. *Angopora hisingeri* is also recoded from Great Britain (Jones, 1936), Norway, Estonia and Poland (Stasinska, 1967), Xinjiang in Northeast China (Wang, 1981), and Ireland (Scrutton, 1997). The present discovery from the Kuraoka area represents the first record of *Angopora* from Japan.

Acknowledgements

We are greatly indebted to the late Dr. Takashi Hamada, who provided guidance of tabulate coral taxonomy and general geology of the Gioniyama Formation. In addition, he offered helpful insights at the initial stages of this work. We thank Dr. Hisayoshi Igo for his helpful review of the manuscript, which materially improved it.

References

- Adachi, T. and Niko, S. (1996) Some Silurian tabulate corals from the Gioniyama Formation, Miyazaki Prefecture. *Chigakukennyu*, **45**: 67–73. (In Japanese.)
- Barskaya, V. F. and Sharkova, T. T. (1963) Tabulyaty i heliolitidy ludlova [Ludrovian Tabulata and Heliolitida]. In Bogdanov, A. A. (ed.), *Stratigrafiya i fauna paleozoyskikh otlozheniy khreba Tarbagatay (Ordovik, silur, devon, nizhniy karbon)* [Stratigraphy and fauna of the Paleozoic deposits of Tarbagatai Range (Ordovician, Silurian, Devonian, Lower Carboniferous)]. pp. 137–161, pls. 17–26, Gosgeoltekhizdat, Moscow. (In Russian.)
- Chernyshev [Tschernyshev], B. B. (1936) Verkhnesiluriyskie i devonskie Tabulata basseina r. Kolymy [The Upper Silurian and Devonian Tabulata of the Kolyma River Region]. In, *Materialy po izucheniyu Okhotsko-Kolymnskogo kraya* [Contributions to knowledge of the Kolyma-Indigirka region]. Ser. 1, Geol. Geomorf., no. 4, pp. 39–46, pls. 1, 2, State Trust Dalstroy, Leningrad and Moscow. (In Russian, not seen.)
- Chernyshev, B. B. (1937a) Siluriyskie i devonskie Tabulata Mongolii i Tuvy [Silurian and Devonian Tabulata of Mongolia and Tuva]. *Tr. Akad. Nauk SSSR, Mong. Kom.*, **6**: 1–31, pls. 1–4. (In Russian.)

Fig. 5. 1, 2, 4, *Thecia ichikawai* sp. nov. thin sections. 1, 4, NMNS PA17395, paratype. 1, longitudinal section of corallum, $\times 5$. 4, transverse sections of thin walled corallites, $\times 10$. 2, NMNS PA17402, paratype, transverse sections of thick walled corallites, $\times 5$. 3, 5, 6, *Angopora hisingeri* (Jones, 1930), thin sections. 3, 6, NMNS PA17392. 3, longitudinal sections of corallum, $\times 5$. 6, partial enlargement of Fig. 5-3, showing details of intercorallite walls, $\times 10$. 5, NMNS PA17391, transverse sections of corallites, $\times 10$.

- Chernyshev, B. B. (1937b) Verkhnesiluriyskie i devonskie Tabulata Novoy Zemli, Severnoy Zemli, i Taymyra [The Upper Silurian and Devonian Tabulata of Novaya Zemlya, Severnaya Zemlya and Taimyr]. *Tr. Vses. Ark-tichi Inst.*, **91**: 67–134, pls. 1–13. (In Russian with English abstract.)
- Chernyshev, B. B. (1951) Siluriskie i devonskie Tabulata i Geliolitida okrain Kuznetskogo uglenosnogo basseyna [Silurian and Devonian Tabulata and Heliolitida from the environs of the Kuznetsk coal basin]. 160 pp, 26 pls., Vses. Nauchno-Issled. Geol. Inst., Gosgeotekhnizdat, Moscow. (In Russian.)
- Grabau, A. W. (1930) Corals of the Upper Silurian *Spirifer tingi* beds of Kweichow. *Bull. Geol. Soc. China*, **9**: 223–247, pls. 1–3.
- Hamada, T. (1959) Gotlandian stratigraphy of the Outer Zone of Southwest Japan. *Jour. Geol. Soc. Japan*, **65**: 688–700. (In Japanese with English abstract.)
- Hamada, T. (1982) Middle Paleozoic. Tabulate corals. In Fujiyama, I. *et al.* (eds.), *The Student Edition. A Picture Book of Japanese Fossils*, pp. 4–7, Hokuryukan, Tokyo. (In Japanese.)
- Hamada, T. (1983) Silurian-Devonian tabulate corals from Japan, S-D-2. In, *Atlas of Japanese Fossils*. No. 39–230, Tsukiji Shokan, Tokyo. (In Japanese.)
- Hill, D. (1981) Part F, Coelenterata. Supplement 1, Rugosa and Tabulata. In Moore, R. C. *et al.* (eds.), *Treatise on Invertebrate Paleontology*, pp. F1–F762, Geol. Soc. America and Univ. Kansas, Boulder, Colorado and Lawrence, Kansas.
- Hill, D. and Stumm, E. C. (1956) Tabulata. In Moore, R. C. *et al.* (eds.), *Treatise on Invertebrate Paleontology*. Part F, Coelenterata, pp. F444–F477, Geol. Soc. America and Univ. Kansas, New York and Lawrence, Kansas.
- Jones, O. A. (1930) A revision of some Paleozoic coral genera and species: Abstr. Dissert. Approved for the Ph. D., M. Sc., and M. Litt. Degrees in the Univ. of Cambridge for the acad. Year 1928–39, p. 35, 36, Univ. Cambridge, Cambridge.
- Jones, O. A. (1936) The controlling effect of environment upon the corallum in *Favosites*: with a revision of some massive species on this basis. *Ann. Mag. Nat. Hist., Ser. 10*, **17**: 1–24, pls. 1–3.
- Kamei, T. (1955) Classification of the Fukui Formation (Silurian) on the basis of *Favosites* with description of some *Favosites*. (Study on Paleozoic rocks of Hida II). *Jour. Fac. Lib. Arts Sci., Shinshu Univ., Part 2, Nat. Sci.*, (5): 39–63, pls. 1–4.
- Kido, E. (2009) *Nanshanophyllum* and *Shensiphyllum* (Silurian Rugosa) from the Kurosegawa Terrane, Southwest Japan, and their paleobiogeographic implications. *Jour. Paleont.*, **83**: 280–292.
- Kido, E. and Sugiyama, T. (2007) Reexamination of the stratigraphy of the Silurian-Devonian and Carboniferous strata in the Gionyama area, Gokase-cho, Miyazaki Prefecture. *Proc. Nishinohon Branch, Geol. Soc. Japan*, (153): 17. (In Japanese.)
- King, Y. (1974) Subclass Tabulata. In, *Atlas of Fossils of Yunnan*, pp. 192–252 (first fascicle), pp. 109–116, 132, 133, 220–251, 336–343, 363–370, 403, 404, 509–519 (second fascicle). The People's Publishing House of Yunnan, Kunming. (In Chinese.)
- Keller, N. B. (1966) Tabulyaty Isenskoï svity [Tabulata of the Isenian Formation]. In Chtverikova N. P. *et al.* (eds.), *Stratigrafiya i fauna siluriyskikh i nizhnedevonskikh otlozheniy Nurinskogo sinlinoriya* [Stratigraphy and fauna of the Silurian and Lower Devonian deposits of the Nurin synclinorium], *Mater. Geol. Tsentr. Kazakhstana*, **6**: 102–114, pls. 7–18. (In Russian.)
- Klaamann, E. R. (1962) Tabulyaty verkhnego silura Estonii [Upper Silurian Tabulata of Estonia]. *Eesti NSV Tead. Akad. Geol. Inst., Uurim.*, **9**: 25–74, pls. 1–16. (In Russian with Estonian and English abstracts.)
- Klaamann, E. R. (1964) Pozdneordovikskie i rannesiluriyskie Favositida Estonii [Late Ordovician and Early Silurian Favositida of Estonia]. 118 pp., 26 pls., Eesti NSV Tead. Akad., Tallinn. (In Russian with Estonian and English abstracts.)
- Klaamann, E. R. (1970) Izmenchivost i taksonomicheskoe polozenie *Angopora hisingeri* (Jones) [Variability and taxonomic position of *Angopora hisingeri* (Jones)]. *Eesti NSV Tead. Akad. Toim.*, **19**: 62–68, pls. 1–4. (In Russian with Estonian and English abstracts.)
- Lamarck, J. B. P. A. de M. de (1816) Histoire naturelle des animaux sans vertèbres, présentant les caractères généraux et particuliers de ces animaux, leur distribution, leurs classes, leurs familles, leurs genres, et la citation des principales espèces qui s'y rapportent; précédé d'une introduction offrant la détermination des caractères essentiels de l'animal, sa distinction du végétal et des autres corps naturels, enfin, l'exposition des principes fondamentaux de la zoologie. Volume 2, 568 pp., Privately published, Paris. (Reissued by Culture et Civilisation, Bruxelles, 1969.)
- Lee, D. J. and Noble, J. P. A. (1988) Evaluation of coral-lite size as a criterion for species discrimination in *Favosites*. *Jour. Paleont.*, **62**: 32–40.
- Lonsdale, W. (1839) Corals. In Murchison, R. I., *The Silurian System, founded on geological researches in the counties of Salop, Hereford, Radnor, Montgomery, Caermarthen, Brecon, Pembroke, Monmouth, Gloucester, Worcester, and Stafford; with descriptions of the Coal-fields and overlying formations*. Part 1, pp. 675–694, pls. 15, 15 bis., 16, 16 bis., John Murray, London.
- Milne-Edwards, H. and Haime, J. (1849) Mémoire sur les polypiers appartenant aux groupes naturels des Zoanthaires perforés et des Zoanthaires tabulés. *Acad. Sci. Paris, C. R.*, **29**: 257–263.
- Milne-Edwards, H. and Haime, J. (1851) Monographie

- des polypiers fossiles des terrains paléozoïques, précédée d'un tableau général de la classification des polypes. *Arch. Mus. Hist. Nat., Paris*, **5**: 1–502, pls. 1–20.
- Niko, S. (1998) Silurian tabulate corals *Eofletcheria* and *Aulocystis* from the Gionyama Formation, Miyazaki Prefecture. *Bull. Natn. Sci. Mus., Tokyo, Ser. C*, **24**: 41–49.
- Niko, S. (2006) Reexamination of Devonian favositid corals described from the Fukuji Formation of Gifu Prefecture. *Bull. Natn. Sci. Mus., Tokyo, Ser. C*, **32**: 13–30.
- Niko, S. and Adachi, T. (1999a) *Gokaselites*, a new genus of Silurian tabulate coral from the Gionyama Formation, Miyazaki Prefecture. *Bull. Natn. Sci. Mus., Tokyo, Ser. C*, **25**: 45–49.
- Niko, S. and Adachi, T. (1999b) Silurian pachyporicaes (Coelenterata: Tabulata) from the Gionyama Formation, Miyazaki Prefecture. *Bull. Natn. Sci. Mus., Tokyo, Ser. C*, **25**: 111–120.
- Niko, S. and Adachi, T. (2000) Silurian multisoleniids (Coelenterata: Tabulata) from the Gionyama Formation, Miyazaki Prefecture. *Bull. Natn. Sci. Mus., Tokyo, Ser. C*, **26**: 107–119.
- Niko, S. and Adachi, T. (2002) Silurian Alveolitina (Coelenterata: Tabulata) from the Gionyama Formation, Miyazaki Prefecture. *Bull. Natn. Sci. Mus., Tokyo, Ser. C*, **28**: 9–24.
- Niko, S. and Adachi, T. (2004) Additional material of Silurian tabulate corals from the Gionyama Formation, Miyazaki Prefecture. *Bull. Natn. Sci. Mus., Tokyo, Ser. C*, **30**: 47–54.
- Niko, S. and Adachi, T. (2008) *Aulostegites nodai*, a new species of Early Silurian tabulate coral from the Gionyama Formation, Miyazaki Prefecture. *Bull. Natn. Sci. Mus., Tokyo, Ser. C*, **34**: 39–42.
- Oekentorp, K. (1970) Revision des Genus *Thecia* Edwards & Haime, 1849 (Coelenterata: Tabulata). *Abh. Neues Jahrb. Geol. Paläont.*, **137**: 103–173.
- d'Orbigny, A. (1850) Prodrome de paléontologie stratigraphique universelle des animaux mollusques et rayonnés faisant suite au cours élémentaire paléontologie et de géologie stratigraphiques. Volume 1, 349 pp., Victor Masson, Paris.
- Rukhin, L. B. (1937) Verkhne-siluriyskie Tabulata Turkestanskogo Khrebt i Khan-tengri [Upper Silurian Tabulata from the Turkestan Range and Khan-tengri], 99 pp., 18 pls., Leningradskiy gosudarstvennyy Universitet, Leningrad. (In Russian.)
- Rukin, L. B. (1938) Nizhnepaleozoiskie korally i stromatoporoidei verkhney chasti basseyna r. Kolymy. [The lower Paleozoic corals and stromatoporids of the upper part of the Kolyma River basin]. In, Materialy po izucheniyu Kolyma-Indigirskogo kraya [Contributions to knowledge of the Kolyma-Indigirka region]. Ser. 2, Geol. Geomorf., no. 10, 119 pp., 28 pls., State Trust Dalstroy, Leningrad and Moscow. (In Russian with English abstract.)
- Scrutton, C. T. (1997) The Palaeozoic corals, I: origins and relationships. *Proc. Yorkshire Geol. Soc.*, **51**: 177–208.
- Shimizu, S., Ozaki, K. and Obata, T. (1934) Gotlandian deposits of northwest Korea. *Jour. Shanghai Sci. Inst., Sect. 2*, **1**: 62–78, pls. 9–18.
- Shrock, R. R. and Twenhofel, W. H. (1939) Silurian fossils from northern Newfoundland. *Jour. Paleont.*, **13**: 241–266, pls. 27–30.
- Sokolov, B. S. (1949) Tabulata i Heliolitida [Tabulata and Heliolitida]. In, Atlas rukovodyashchikh form iskopaemykh faun SSSR, II. Siluriyskaya sistema [Atlas of the index forms of the fossil fauna USSR, II. Silurian System]. pp. 75–98, pls. 6–10, Gosgeoltekhizdat, Moscow. (In Russian.)
- Sokolov, B. S. (1950) Sistematika i istoriya razvitiya paleozoyskikh korallov Anthozoa tabulata [Systematics and history of the development of the Paleozoic corals Anthozoa Tabulat]. *Vopr. Paleont.*, **1**, 134–210. (In Russian.)
- Sokolov, B. S. (1951) Tabulyaty paleozoya evropeyskoy chasti SSSR. Chast 2. Silur Pribaltiki (Favozitidy llandoveryanskogo yarusy) [Paleozoic Tabulata of the European parts of the USSR. Part 2. Silurian of the Baltic area (Favositidae of the Llandovery stage)]. *Tr. Vses. Nauchno-Issled. Geol.-Razved. Inst., N. S.*, **52**: 1–124, pls. 1–37. (In Russian.)
- Sokolov, B. S. (1952) Tabulyaty paleozoya evropeyskoy chasti SSSR. Chast 3. Silur Pribaltiki (Favozitidy venlokskogo ludlovskogo yarusov) [Paleozoic Tabulata of the European parts of the USSR. Part 3. Silurian of the Baltic area (Favositidae of the Wenlock-Ludlow stages)]. *Tr. Vses. Nauchno-Issled. Geol.-Razved. Inst., N. S.*, **58**: 1–85, pls. 1–22. (In Russian.)
- Sokolov, B. S. (1955) Tabulyaty paleozoya evropeyskoy chasti SSSR, Vvedenie. Obshchie voprosy sistematiki i istorii razvitiya tabulyat [Paleozoic Tabulata of the European parts of the USSR, Introduction to the general study of the systematics and development of the tabulates]. *Tr. Vses. Nauchno-Issled. Geol.-Razved. Inst., N. S.*, **85**: 1–527. (In Russian.)
- Stasinska, A. (1967) Tabulata from Norway, Sweden and from the erratic boulders of Poland. *Palaeont. Polonica*, (18): 1–112, pls. 1–38.
- Stel, J. H. (1978) Environment and quantitative morphology of some Silurian tabulates from Gotland. *Scripta Geol.*, **47**: 1–75.
- Stel, J. H. and Oekentorp, K. (1976) On the solenoid growth habit of *Paleofavosites*. *Geol. Mijinbouw*, **55**: 163–174.
- Sugiyama, T. (1940) Stratigraphical and palaeontological studies of the Gotlandian deposits of the Kitakami

- Mountainland. *Tohoku Imp. Univ., Sci. Rep., Ser. 2*, **21**: 81–146, pls. 13–33.
- Twenhofel, W. H. (1914) The Anticosti Island faunas. *Canada Geol. Surv. Mus. Bull.*, (3): 1–39.
- Tsin, H. (1956) New material of Silurian fossils from the vicinity of Kueiyang, Central Kueichou. *Acta Palaeont. Sinica*, **4**: 621–639, pls. 1–4. (In Chinese with English abstract.)
- Wang, B. (1981) Tabulata and Heliolitina. In, Palaeontological Atlas of Northwestern Regions, Xinjiang Uygur. Volume 1, Late Proterozoic-Early Palaeozoic, pp. 39–73, pls. 22–36, Geological Publishing House, Beijing. (In Chinese.)
- Yabe, H. and Hayasaka, I. (1915) Palaeozoic corals from Japan, Korea and China. I. *Michelinia* and *Favosites*. *Jour. Geol. Soc. Tokyo*, **22**: 55–70.
- Yang, C. (1948) The Silurian Waseh Formation of western Yunnan and its *Favosites* faunas. *Natn. Peking Univ. Ann. Pap., Geol. Ser.*, **50**: 129–140, pl. 1.
- Yang, S., Kim, C. and Chow, X. (1978) Tabulata. In, Atlas of the Palaeontology of the Southwestern Region of China, Guizhou [Kweichow]. Volume 1, Cambrian-Devonian, pp. 161–251, pls. 56–93, Geological Publishing House, Beijing. (In Chinese.)
- Young, G. A. and Elias, R. J. (1995) Latest Ordovician to earliest Silurian colonial corals of the East-Central United States. *Bull. Amer. Paleont.*, **108**: 1–153, pls. 1–21.
- Yü, C. (1956) Some Silurian corals from the Chiuchüan Basin, western Kansu. *Acta Palaeont. Sinica*, **4**: 599–620, pls. 1, 2. (In Chinese with English abstract.)
- Yü, C. (1962) Middle Silurian coral fossils from northern Mt. Qilianshan. *Geol. Mt. Qilianshan*, **4**: 13–109, pls. 1–42. (In Chinese.)
- Yü, C., Wu, W., Zhao, J. and Zhang, G. (1963) Chinese Fossil Corals. 390 pp., 98 pls. Science Publishing House, Beijing. (In Chinese.)