

Source of the Rhyolitic Ash Layer in Basaltic Ash Layers on Izu-Ōshima

By

Tokiko TIBA

Department of Geology, National Science Museum,
3–23–1 Hyakunincho, Shinjuku-ku, Tokyo, 169

Abstract Chemical compositions of volcanic glass, plagioclase and biotite in the rhyolitic ash layer intercalated in the basaltic ash layers of the Younger Ōshima Group in Izu-Ōshima are compared with those in ash from Mukaiyama (Niijima) and Tenjōsan (Kōzushima) volcanoes erupted in A.D. 838 and 886, respectively. High ΣFeO and Na_2O contents of volcanic glass and high Mg^* values and low ΣFe contents of biotite in the rhyolitic ash from Ōshima are features closer to those from Niijima than from Kōzushima, suggesting the origin from Mukaiyama volcano in Niijima.

Introduction

Izu-Ōshima island, the northernmost island in the seven Izu islands running from east off Izu Peninsula south to Marianas. The major part of the island consists of material erupted from Ōshima volcano, an active basaltic stratovolcano. Recent products of Ōshima volcano called the Younger Ōshima Group are divided into three formations, i.e. Yuba, Nomashi and Sashikiji Formations in descending order (NAKAMURA, 1960). A basaltic ash layer N_3 in Nomashi Formation is conspicuous for intercalation of rhyolitic ash layer supplied from a volcano other than Ōshima volcano. This rhyolitic ash layer had been recognized in N_3 throughout the island (NAKAMURA, 1964), though it varies from <0.1 cm to 1 cm or more in thickness. NAKAMURA (1964) inferred the date of eruption produced N_3 member as ninth century from historical documents and pottery remains. Based on the result of mechanical analyses of the rhyolitic ash, he further referred the source of the rhyolitic ash to Kōzushima or Niijima where Tenjōsan and Mukaiyama volcanoes erupted in 838 and 886 A.D., respectively. Since then, it remains unsettled either of the above two volcanoes is the source volcano of this rhyolitic ash, for both volcanoes provided rhyolitic lavas and pyroclastics almost identical in chemical composition and constituting minerals.

In this paper, the author intended to specify the source volcanoes of this rhyolitic ash by putting chemical data of ash constituents such as volcanic glass, plagioclase and biotite together.




Fig. 1. Index map of Ōshima, Niiijima and Kōzushima.

Samples and Analytical Method

On Ōshima island, rhyolitic ash samples are collected from several exposures on northern slope outside the caldera wall. Ash samples of 838 A.D. eruption of Tenjōsan volcano in Kōzushima (ISSHIKI, 1982) and 886 eruption of Mukaiyama volcano in Niiijima (ISSHIKI, 1987) are taken from the eastern coast of the islands closer to Ōshima island (Fig. 1). Common constituents of the ash samples are volcanic glass, quartz, plagioclase, biotite and magnetite. Ash from Niiijima contains olivine, clinopyroxene and orthopyroxene provided by basaltic volcanism occurred possibly on Miyakejima.

The ash samples are washed in distilled water and dried in the room temperature. Volcanic glass fragments, plagioclase grains and biotite flakes for analysis are hand picked under the binocular microscope. Those fraction are fixed on slide glass with epoxy resin and polished. Chemical analysis of the samples was done by using Link Systems energy dispersive X-ray sepcrometer.

Chemical Composition

Volcanic glass: Major oxide analyses of volcanic glasses are given in Appendices 1–3. Rhyolitic glass JR-2 is used to inspect the validity of the analyses. The analyses are plotted on oxide- K_2O diagrams (Fig. 2) after conversion to anhydrous total = 100%. In Fig. 2, K_2O content is taken as abscissa, for K is the sole incompatible element analysed that tends to enrich in high silica fluid




Fig. 2. Plots of volcanic glass analyses on SiO₂, Al₂O₃, ΣFeO, CaO and Na₂O contents vs. K₂O diagrams.

phase till potash feldspar begins to crystallize.

As seen in Fig. 2, plots of volcanic glasses from Niiijima fall on the lower K₂O side, those from Kōzushima on the high K₂O side and those from Ōshima on intermediate area between them. As far as the major oxide contents concerned, volcanic glasses with somewhat high ΣFeO and Na₂O are found both ash samples from Niiijima and Ōshima.

Plagioclase: Chemical analyses of plagioclase grains are shown in Appendix 4 and plotted on An–Ab–Or system (Fig. 3). Plagioclase in ash from Niiijima covers the widest compositional range from An₃₄ to An₁₆. On the other hand, plagioclase from Kōzushima shows a limited compositional variation from An₂₂ to An₁₇. Some plagioclase grains from Ōshima are much sodic than those from Niiijima and Kōzushima.

Biotite: Biotite flakes in ash from Ōshima are usually more or less altered, presenting paler color than those from Niiijima and Kōzushima. This must have been caused by meteoric water preserved more abundantly in basaltic ash layers that overlay and underlay the rhyolitic ash layer. Major element analyses of biotite (Appendix 5) reflect the effect of alteration, i.e. compositions of biotite from Ōshima are characterized by low ΣFe and higher Mg contents than those from Niiijima and Kōzushima. Plots of the analyses on atomic ratios-1 Mg*




Fig. 3. Plots of plagioclase analyses on An-Ab-Or system.

diagrams (Fig. 4) demonstrate selective extraction of iron from biotite, resulting high Mg^* values and low ΣFe atomic ratios in biotite from Ōshima, though the degree of alteration is not defined. These diagrams also present higher ΣFe ratios and lower Mg^* values of those from Kōzushima than those from Niiijima.

Considerations

The ninth century eruptions on Niiijima and Kōzushima yielded rhyolitic lavas almost identical in bulk chemistry (see ISSHIKI, 1987 & 1982), together with rhyolitic ash. The present data, however, allow to regard rhyolitic ash from Tenjōsan volcano (Kōzushima) is derived from much evolved magma than those of Mukaiyama volcano (Niiijima), since rhyolitic ash from Kōzushima includes K rich volcanic glass, less calcic plagioclase and biotite with low Mg^* values. Judging from low ΣFe ratios in biotite and high ΣFeO and Na_2O contents of volcanic glass in rhyolitic ash from Ōshima, Mukaiyama volcano on Niiijima is the likely source supplied the rhyolitic ash layer in the Younger Ōshima Group in Ōshima.




Fig. 4. Plots of biotite analyses on Si, Ti, Al, Fe, Mn, Mg, Na and K atomic ratios (on anhydrous basis of O=22)-Mg* diagram. A bulk analysis of biotite phenocrysts (Mg*=53, ISSHIKI, 1987) in the rhyolite lava extruded contemporaneously with the volcanic ash studied is also plotted together for comparison.

* $100\text{Mg}/(\text{Mg} + \Sigma\text{Fe} + \text{Mn})$

Acknowledgment

The author is grateful for the kind assistance in this work provided by Mrs. A. SHIGEOKA and Dr. K. YOKOYAMA.

References

- ISSHIKI, N., 1982. Geology of the Kōzushima district. *Quadrangle series, scale 1:50,000*, Geol. Surv. Japan, 75 pp. with 8 plates. (In Japanese with English abstract.)
- , 1984. Geology of the Ō Shima district. *Quadrangle series, scale 1:50,000*, Geol. Surv. Japan, 133 pp. with 8 plates. (In Japanese with English abstract.)
- , 1987. Geology of the Nii Jima district. *Quadrangle series, scale 1:50,000*, Geol. Surv. Japan, 85 pp. with 8 plates. (In Japanese with English abstract.)
- NAKAMURA, K., 1960. Stratigraphic studies of the pyroclastics of Ōshima Volcano, Izu, deposited during the last fifteen centuries. I. Cyclic activity of "Main craters" and the absolute chronology of the pyroclastic sediments. *Sci. Pap. Coll. Gen. Education, Univ. Tokyo*, 10: 125–145.
- , 1964. Volcano-stratigraphic study of Ōshima volcano, Izu. *Bull. Earthq. Res. Inst., Univ. Tokyo*, 42: 649–728 with 10 plates.

Appendix 1. Major oxide analyses of volcanic gasses from Ōshima.

wt%	1	2	3	4	5	6	7	8	9	10	11	12	13
SiO ₂	75.77	74.86	75.56	76.13	75.98	75.26	77.07	75.52	75.47	75.09	76.75	75.53	77.03
TiO ₂	—	—	—	—	—	—	—	—	—	0.24	—	—	—
Al ₂ O ₃	12.32	12.27	12.70	12.38	12.56	12.22	12.46	12.51	12.52	12.44	12.37	12.33	12.72
ΣFeO*	0.58	0.59	0.59	0.52	0.71	0.60	0.40	0.47	0.48	0.78	0.74	0.62	0.84
MnO	—	—	—	—	—	0.23	—	—	—	—	0.26	0.29	—
CaO	0.31	0.41	0.43	0.43	0.35	0.31	0.26	0.37	0.33	0.39	0.38	0.32	0.41
Na ₂ O	4.02	3.92	4.46	4.20	4.10	4.05	4.18	4.52	4.14	3.76	4.47	4.12	4.18
K ₂ O	3.81	3.96	3.84	3.96	3.77	3.95	4.11	3.81	3.77	3.68	3.83	4.15	3.91
Cl	—	—	0.09	—	0.12	—	0.09	0.07	0.10	—	0.10	—	0.07
	96.81	96.01	97.67	97.62	97.59	96.62	98.57	97.27	96.81	96.38	98.90	97.36	99.16
O=Cl	—	—	-0.02	—	-0.03	—	-0.02	-0.02	-0.02	—	-0.02	—	-0.02
total	96.81	96.01	97.65	97.62	97.56	96.62	98.55	97.25	96.79	96.38	98.88	97.36	99.14

* total iron as FeO

Appendix 2. Major oxide analyses of volcanic gasses from Nijijima.

wt%	1	2	3	4	5	6	7	8	9	10	11
SiO ₂	76.44	75.04	77.94	75.21	76.03	74.44	75.25	74.83	77.12	78.00	75.98
TiO ₂	—	0.25	—	—	—	—	—	—	—	0.24	—
Al ₂ O ₃	12.75	12.21	12.63	12.04	12.44	12.26	12.33	12.07	12.59	12.74	12.35
ΣFeO*	0.65	0.56	0.76	0.59	0.55	0.73	0.65	0.42	0.80	0.71	0.63
MnO	—	—	—	—	—	0.26	—	—	—	—	0.26
CaO	0.55	0.59	0.50	0.54	0.22	0.47	0.41	0.48	0.44	0.41	0.35
Na ₂ O	4.44	4.14	4.61	4.03	4.26	4.29	4.01	3.97	4.48	4.06	3.89
K ₂ O	3.41	3.49	3.82	3.62	3.95	3.66	3.76	3.72	3.96	3.70	3.73
Cl	0.07	0.10	0.07	0.09	—	0.07	0.07	—	0.07	—	0.09
	98.31	96.38	100.33	96.12	97.45	96.18	96.48	95.49	99.46	99.62	97.02
O=Cl	-0.02	-0.02	-0.02	-0.02	—	-0.02	-0.02	—	-0.02	—	-0.02
total	98.29	96.36	100.31	96.10	97.45	96.16	96.46	95.49	99.44	99.62	97.00

* total iron as FeO

Appendix 3. Major oxide analyses of volcanic gasses from Kōzushima.

wt%	1	2	3	4	5	6	7	8	9	10	11	12	13
SiO ₂	75.04	74.79	76.03	74.25	75.58	75.22	74.69	75.79	74.68	75.27	75.92	75.51	74.31
TiO ₂	0.21	—	—	—	—	—	—	—	—	—	—	—	—
Al ₂ O ₃	12.28	12.38	12.36	12.41	12.51	12.42	12.24	12.54	11.65	12.52	12.48	12.63	12.43
ΣFeO	0.52	0.59	0.27	0.46	0.64	0.63	0.68	0.63	0.44	0.61	0.52	0.64	0.60
CaO	0.37	0.38	0.35	0.46	0.39	0.37	0.41	0.37	0.33	0.32	0.33	0.44	0.40
Na ₂ O	4.08	3.49	4.07	3.71	4.07	4.27	4.21	4.24	3.47	3.96	4.13	4.22	4.26
K ₂ O	3.80	3.96	4.14	3.83	4.25	4.00	3.92	3.93	4.59	3.88	4.15	3.95	3.80
Cl	—	0.07	0.07	0.06	0.03	0.06	0.09	0.07	—	0.06	0.09	0.06	0.07
	96.30	95.66	97.29	95.18	97.47	96.97	96.24	97.57	95.16	96.62	97.62	97.45	95.87
O=Cl	—	-0.02	-0.02	-0.01	-0.01	-0.01	-0.02	-0.02	—	-0.01	-0.02	-0.01	-0.02
total	96.30	95.64	97.27	95.17	97.46	96.96	96.22	97.55	95.16	96.61	97.60	97.44	95.85

* total iron as FeO

Appendix 4. Major oxide analyses of plagioclases.

wt%	Ōshima								
	1	2	3	4	5	6	7	8	9
SiO ₂	65.42	64.52	63.98	64.01	63.43	63.70	66.10	63.91	66.45
Al ₂ O ₃	22.00	22.28	21.29	23.1	22.68	22.48	21.09	22.05	20.73
ΣFeO	—	0.40	0.31	—	0.33	0.35	—	0.29	—
CaO	3.32	4.00	3.23	4.27	4.23	3.94	2.25	3.29	2.54
Na ₂ O	8.64	8.95	9.04	8.42	8.12	9.40	9.63	9.03	9.23
K ₂ O	0.81	0.62	0.83	0.66	0.67	0.62	1.03	0.76	0.94
total	100.19	100.78	98.69	100.48	99.46	100.49	100.10	99.34	99.89
Ab	78.5	77.3	79.5	75.1	74.5	78.5	83.4	79.6	82.1
An	16.7	19.2	15.7	15.7	21.4	18.1	10.8	16.0	12.4
Or	4.8	3.5	4.8	4.8	4.1	3.4	5.9	4.4	5.5

	Niijima											
	1	2	3	4	5	6	7	8	9	10	11	12
SiO ₂	64.16	64.44	60.78	63.69	64.05	61.35	64.40	63.75	63.48	64.16	63.94	62.55
Al ₂ O ₃	22.44	22.07	20.08	22.21	21.57	24.71	22.24	22.58	22.57	22.53	22.54	22.53
ΣFeO	—	—	—	—	—	—	—	—	—	—	—	—
CaO	3.70	4.00	6.08	3.71	3.25	6.95	3.62	4.13	4.17	4.11	3.97	4.96
Na ₂ O	8.93	8.68	7.77	8.59	8.88	7.44	8.96	8.08	8.10	8.91	8.64	8.40
K ₂ O	0.63	0.53	0.35	0.62	0.75	0.21	0.85	0.67	0.55	0.65	0.64	0.53
total	99.86	99.72	99.06	98.82	98.50	100.66	100.07	99.21	98.87	100.36	99.73	98.97
Ab	78.4	77.3	68.4	77.8	79.5	65.1	77.8	74.8	75.3	76.7	76.8	73.1
An	17.9	19.6	29.6	18.5	16.1	33.6	17.3	21.1	21.4	19.6	19.5	23.9
Or	3.7	3.1	2.0	3.7	4.3	1.2	4.9	4.1	3.3	3.7	3.7	3.0

	Kōzushima								
	1	2	3	4	5	6	7	8	9
SiO ₂	63.95	64.82	64.72	64.32	63.67	64.20	64.10	63.92	63.33
Al ₂ O ₃	23.20	22.70	22.19	23.13	22.76	22.19	23.13	21.90	22.65
ΣFeO	—	—	0.38	—	—	—	—	—	—
CaO	4.22	3.91	3.44	4.31	4.45	3.71	4.36	3.62	4.24
Na ₂ O	9.07	8.54	9.02	8.75	8.14	9.26	8.23	8.33	8.60
K ₂ O	0.55	0.62	0.78	0.48	0.63	0.65	0.64	0.75	0.72
total	100.99	100.59	100.53	100.99	99.65	100.01	100.46	98.52	99.54
Ab	77.1	77.0	78.9	76.4	73.9	78.8	74.4	76.9	75.3
An	19.8	19.4	16.6	20.8	22.3	17.5	21.8	18.5	20.5
Or	3.1	3.6	4.5	2.8	3.8	3.7	3.8	4.6	4.2

Appendix 5. Major oxide analyses of biotites.

wt%	Ōshima					Nijima				Kōzushima										
	1	2	3	4	5	1	2	3	4	1	2	3	4	5	6	7	8	9	10	11
SiO ₂	37.43	37.43	37.67	38.08	37.50	38.24	37.55	36.99	37.76	36.57	36.62	36.81	36.89	37.27	37.15	35.40	35.56	36.19	35.00	35.93
TiO ₂	4.00	4.04	4.06	4.1	4.28	4.02	4.29	4.09	4.03	4.05	4.00	4.15	4.15	4.27	4.27	4.17	4.08	4.35	4.30	4.39
Al ₂ O ₃	13.78	13.64	13.34	13.31	13.38	13.80	14.17	13.70	13.89	13.98	13.77	13.85	13.98	14.21	13.98	13.96	13.36	13.42	13.30	13.30
ΣFeO*	15.38	15.72	15.25	15.44	15.63	16.74	21.08	18.86	17.07	20.22	19.45	19.48	19.58	20.30	20.17	20.52	20.52	21.10	20.43	20.24
MnO	0.37	0.42	0.40	—	0.40	0.34	0.60	0.52	0.36	0.63	0.55	0.50	0.55	0.61	0.42	0.42	0.58	0.38	0.71	0.65
MgO	14.70	14.81	14.97	14.63	14.02	12.25	11.21	12.34	12.51	11.75	11.19	11.19	11.40	11.55	11.60	10.80	10.36	11.15	10.41	10.60
CaO	—	—	—	—	—	0.16	0.08	—	0.19	—	—	—	—	—	—	—	—	—	—	—
Na ₂ O	0.72	0.81	0.63	0.97	0.68	0.70	0.70	0.73	0.29	0.79	0.81	0.37	0.82	0.83	0.90	0.89	0.78	—	0.84	0.57
K ₂ O	8.55	8.49	8.76	8.86	8.60	8.92	8.75	8.35	8.36	8.65	8.69	8.63	8.61	8.72	8.93	8.74	8.62	8.75	8.98	8.65
Cl	0.10	0.08	0.07	0.08	—	0.10	0.11	0.12	0.10	0.11	0.11	0.16	0.11	0.10	0.11	0.13	0.13	0.13	0.14	0.18
O=Cl	95.03	95.44	95.15	95.47	94.49	95.27	98.54	95.70	94.56	96.75	95.19	95.14	96.09	97.86	97.53	95.03	93.99	95.47	94.11	94.66
total	—0.02	—0.02	—0.02	—0.02	—	—0.02	—0.02	—0.03	—0.02	—0.02	—0.02	—0.03	—0.02	—0.02	—0.02	—0.03	—0.03	—0.03	—0.03	—0.04
atomic ratios on anhydrous basis of O=22	95.01	95.42	95.13	95.45	94.49	95.25	98.52	95.67	94.54	96.73	95.17	95.11	96.07	97.84	97.51	95.00	93.96	95.44	94.08	94.62
Si	5.621	5.608	5.561	5.692	5.667	5.774	5.589	5.613	5.722	5.541	5.619	5.638	5.602	5.571	5.577	5.491	5.579	5.574	5.511	5.582
Ti	0.451	0.455	0.458	0.461	0.487	0.456	0.480	0.466	0.459	0.462	0.462	0.478	0.474	0.480	0.482	0.487	0.482	0.504	0.509	0.513
Al	2.440	2.409	2.360	2.344	2.383	2.456	2.486	2.449	2.480	2.496	2.492	2.502	2.502	2.504	2.474	2.552	2.470	2.438	2.467	2.463
Fe	1.931	1.969	1.913	1.930	1.976	2.067	2.623	2.394	2.163	2.563	2.496	2.495	2.487	2.537	2.532	2.663	2.692	2.717	2.690	2.630
Mn	0.047	0.053	0.050	—	—	0.440	0.076	0.067	0.046	0.081	0.071	0.065	0.071	0.077	0.053	0.056	0.077	0.049	0.095	0.086
Mg	3.291	3.308	3.348	3.260	3.159	2.757	2.488	2.792	2.826	2.654	2.561	2.556	2.580	2.573	2.597	2.498	2.422	2.561	2.443	2.452
Ca	—	—	—	—	—	—	—0.013	—	0.030	—	—	—	—	—	—	—	—	—	—	—
Na	0.209	0.235	0.184	0.282	0.198	0.205	0.202	0.213	0.086	0.233	0.240	0.110	0.243	0.241	0.263	0.268	0.238	—	0.257	0.172
K	1.639	1.624	1.676	1.690	1.658	1.718	1.662	1.618	1.617	1.673	1.700	1.686	1.668	1.663	1.711	1.730	1.725	1.719	1.803	1.716
Mg**	62.5	62.1	63.0	62.8	60.9	56.6	48.0	53.2	56.1	50.1	49.9	50.0	50.2	49.6	50.1	47.9	46.7	48.1	46.7	47.3

* total iron as FeO

** 100Mg/(Mg+ΣFe+Mn)