

Chronometric Dating of Bone from Namib IV Acheulean Site, South West Africa/Namibia

By

Myra SHACKLEY¹, Kazuhisa KOMURA², Takumi HAYASHI³,
Motoji IKEYA⁴, Shuji MATSU'URA⁵ and Nobuo UETA⁶

¹ Department of Archaeology, University of Leicester, Leicester

² Low Level Radioactivity Laboratory, Faculty of Science, Kanazawa University,
Nomi-gun, Ishikawa Prefecture

³ Institute of Plasma Physics, Nagoya University, Nagoya

⁴ Technical College, Yamaguchi University, Ube

⁵ Department of Anthropology, National Science Museum, Tokyo

⁶ Department of Biochemistry, Teikyo University School of Medicine, Tokyo

Abstract The association of fossil bone, including *Elephas recki*, with an Acheulean handaxe and cleaver industry at the ancient kill/butchery site of Namib IV is the earliest evidence for man in the Namib desert. Faunal and typological parallels with east African sites suggested a mid-Pleistocene date, which this program of uranium-series, ESR and racemization dating has corroborated with a most probable age estimate of 347^{+78}_{-48} kyr BP.

Introduction

Namib IV is an artefact concentration, with associated bone, found on a pan calcretes surface at 23°47'S 15°20'E in the dune sea of the central Namib desert. The assemblage includes a heavy-tool component of handaxes and cleavers, scattered over an area of 62,500 m². A systematic survey was made in 1979 and further faunal remains collected in 1981 (SHACKLEY, 1980, 1985). Many of the artefacts and bones were still lightly cemented on their ventral sides into deflation surfaces of lacustrine carbonates, possibly slightly altered by surface pedogenesis. The fauna consisted of mineralised and fragmented bone and tooth remains from *Elephas recki* (minimum 8 individuals), an indeterminate alcelaphine antelope (2 individuals) and a second, large-medium antelope (3 individuals) together with 1 small horse and 1 buffalo (all identified by Professor Richard KLEIN). The elephant remains are fortunately largely composed of diagnostic tooth fragments whose thin enamel, hypsodonty and tight enamel folding suggest a late stage of *Elephas recki*, probably comparable with MAGLIO's Stage 4 (MAGLIO, 1973) as represented in Olduvai Bed IV of mid-Pleistocene date. The relatively small quantities of bone and stone recovered, and the nature of the assemblage, make it unlikely that Namib IV represents a major camp site, and the material is seen as the debris of a few short-term visits at some stage in the mid-Pleistocene.

A dating program on the Namib IV bone was set about in 1982, involving chemical analysis, racemization, electron spin resonance (ESR) and uranium-series methods. The present paper reports the preliminary results of this project.

Material and Methods

Dense cortex of a bone fragment (code no. 3) from Namib IV was subjected to uranium-series dating, ESR dating and fluorine analysis. Racemization dating was applied to another compact bone specimen (code no. 1) from the same site. The analytical techniques have been described elsewhere: e.g. KOMURA & SAKANOUÉ (1967) for uranium series, IKEYA (1985) for ESR, MATSU'URA (1981) for fluorine, and MATSU'URA & UETA (1980) for racemization.

Results and Discussion

$^{230}\text{Th}/^{234}\text{U}$ Dating

The results of the radiochemical analysis and the calculated date are given in Table 1. Contamination of any significant amount of thorium, including ^{230}Th , from the environment to the sample interferes with the age determination. In this case, the absence of detectable ^{232}Th (common thorium) gives evidence for lack of such contamination. This implies that all ^{230}Th activity in the fossil bone is authigenic, thus no correction for the initial ^{230}Th that would have accompanied non-radiogenic ^{232}Th is necessary.

The bulk of the uranium now in the bone sample is believed to be due to diagenetic gains from the burial media during fossilization. Provided that no uranium loss by leaching has taken place, the ^{230}Th date of 347^{+78}_{-48} kyr BP should represent a minimum age estimate.

ESR Dating

Table 2 shows the ESR measurement result of the total dose of natural radiation (TD) to which the bone had already been exposed; together with the contents of elements involved in the age estimation, and the external gamma-ray dose rate (Dex) measured with thermoluminescence dosimeters of CaSO_4 (Tm) embedded at Namib

Table 1. Radiometric data and uranium-series age for fossil bone from Namib IV.

Sample code	U* (ppm)	Activity (dpm/g)				^{230}Th age† ($\times 10^3$ years)
		^{235}U *	^{234}U *	^{232}Th	^{230}Th *	
3	65.1	20.73	29.82	<1**	31.57	347
		± 0.41	± 0.55		± 0.39	+78
						-48

* Determined by alpha spectrometry.

** Below the detection limit of gamma-ray measurement.

† Calculated using ^{230}Th half-life of 75,200 years and ^{234}U half-life of 245,000 years.

IV for 18 months.

One should note that ESR provides only TD which increases with the lapse of time. Unless the annual dose or the radiation environment is assessed precisely, the age can not be obtained. Bone is virtually an "open" system, presenting some difficulties in evaluating the past dose rate (IKEYA, 1983, 1985). For converting the measured TD into age, here are taken the following assumptions:

- 1) a linear accumulation of uranium from the initial zero ppm to the present level (IKEYA, 1983, 1985),
- 2) a continuous fluorine accumulation at a constant rate (Recent unpublished experimental results by IKEYA suggest the reduction of ESR signal intensity by fluorination of bone. See IKEYA, 1985.),
- 3) 50% rate of radon loss out of the bone, which has been estimated from high-resolution gamma-ray spectroscopy.

The alpha-ray efficiency for defect production (k) is also a factor of the dose rate. The k -value is generally 0.1–0.2 (IKEYA, 1983). Using $k=0.15$, which has yielded reasonable ESR ages for corals (IKEYA & OHMURA, 1983), we have a ESR date of ~ 140 kyr BP for the Namib IV specimen. This calculation involves a model of ^{235}U -series disequilibrium (IKEYA, 1983, 1985; IKEYA & OHMURA, 1983). The k -value of 0.15 may be reduced, close to zero, where the high ^{235}U content of the bone is due to the localisation of this element along surface layers, cracks and crevices, and/or to the occurrence as void fillings. This shifts the ESR date towards the older side, and a tentative use of $k=0$ generates 350 kyr BP; this might be refined by a fission track map of the uranium distribution within the bone.

Table 2. Analytical data for ESR dating of Namib IV specimen.

Sample code	^{235}U (ppm)	^{232}Th (ppm)	K_2O (%)	F (%)	Dex* (mGy/y)	TD* (Gy)
3	65.1 [†]	~ 0 [†]	0.1	1.54	0.5	500

* See text.

[†] See Table 1.

As was stated previously, the age evaluation assumes a constant uptake of fluorine. However, if the fluorine content in bone can be expressed by an irreversible first-order rate equation as presented by MATSU'URA (1982), it shifts the ESR age again to the old side. Refined ages will be published elsewhere when further studies are completed.

Racemization Dating

Racemization analysis on the total hydrolysate of a fossil bone (code no. 1) from Namib IV yielded D/L amino acid ratios of 0.35 for alanine, 0.15 for valine, and 0.16 for alloisoleucine/isoleucine. BADA (1981) has shown that the Pleistocene rate constant of the isoleucine epimerization (i.e. k_{iso}) in fossil teeth from the Olduvai Gorge corresponds to an average integrated palaeotemperature of $\sim 16^\circ\text{C}$ and that this

k_{iso} value is also applicable to bones from the same region. The present day mean annual air temperature at Olduvai is $\sim 23^{\circ}\text{C}$ (HAY, 1976; BADA, 1981), while at Gobaabeb, the nearest first-order weather station to Namib IV, it is 20.5°C (GOUDIE, 1972). Provided that a difference of this magnitude had occurred in the Pleistocene and that the 2.5°C disparity represents the difference of the average temperatures to which the fossil specimens from Namib IV and Olduvai have been exposed, the Namib k_{iso} constant (yr^{-1}) can be estimated from the following equation (see BADA *et al.*, 1973, and BADA, 1981):

$$\ln \{k_{\text{iso}}/(1.3 \times 10^{-6})\} = \{(33.4 \times 10^3)/1.987\} \cdot [(-2.5)/\{289 \times (289 - 2.5)\}]$$

Using the k_{iso} value ($7.8 \times 10^{-7} \text{ yr}^{-1}$) calculated as above and the measured D-alloisoleucine/L-isoleucine ratio a date of ~ 200 kyr BP is obtained. It should, however, be noted that older bones are susceptible to contamination by secondary amino acids. Such contamination would introduce mainly L-amino acids, and in this case L-isoleucine not D-alloisoleucine. The order of the amino acid racemization observed in the Namib specimen (i.e. alanine < isoleucine < valine) is compatible with the sequence expected from elevated temperature kinetics experiments (BADA *et al.*, 1973; DUNGWORTH *et al.*, 1976), suggesting that the bone sample was not badly contaminated. The isoleucine dating thus represents a significant minimum age estimate although an *in situ* calibration of the racemization rate constants (see MASTERS, 1982; MATSU'URA & UETA, 1980) and analysis with fractionation treatment (MATSU'URA & UETA, 1980) of further specimens such as fossilized teeth would provide a better and more reliable chronology of Namib IV.

In conclusion, this joint work has been clarifying the age of the Namib IV archaeological site. The ^{230}Th date of 347_{-48}^{+78} kyr BP may be taken for the present as a most substantial and quoteable age estimate.

The suggested date accords well with the typological and faunal affinities of Namib IV, which has close parallels with many east African sites especially Olorgesailie which ISAAC (1977, 1978) considers dates to 400–500 kyr BP. Analogous Acheulean industries calibrated by K-Ar dating include Kapthurin in the central Rift Valley falling within the range 0.23–0.7 my BP (TALLON, 1978), Kilombe (site GqJhl), which postdates a trachyphonolite dated at 1.7 ± 0.05 my BP (BISHOP, 1978), and Chesowanja (site 10/4) with a maximum age of 1.42 ± 0.07 my BP (HOOKER & MILLER, 1979; GOWLETT *et al.*, 1981). The new dates confirm the provisional assignment of Namib IV to the mid-Pleistocene on typological and faunal grounds, and make it the earliest dated human activity in the Namib desert.

Acknowledgements

We wish to acknowledge that this research program was supported in part by the Grants in Aid for Specific Fields and for Fundamental Scientific Research from the Ministry of Education, Science and Culture, Japan, and by the Grant for Promotion of Sciences from Asahi Shimbun Publishing Co. Financial support for M. SHACKLEY

was provided by the Boise Fund (University of Oxford), the British Academy, the L.S.B. Leakey Foundation, the British Council, a Suzette Taylor Travelling Fellowship (Lady Margaret Hall, Oxford) and the Research Board, University of Leicester.

References

- BADA, J. L., 1981. Racemization of amino acids in fossil bones and teeth from the Olduvai Gorge region, Tanzania, East Africa. *Proceedings of the 10th Congress of the International Union of Prehistory and Protohistory*, Commission II (Mexico): 123–138.
- BADA, J. L., K. A. KVENVOLDEN and E. PETERSON, 1973. Racemization of amino acids in bones. *Nature*, **245**: 308–310.
- BISHOP, W. W., 1978. Geological framework of the Kilombe Acheulian archaeological site, Kenya. In BISHOP, W. W. (ed.) "Geological Background to Fossil Man", Edinburgh, Scottish Academic Press, 329–336.
- DUNGWORTH, G., A. W. SCHWARTZ and L. VAN DE LEEMPUT, 1976. Composition and racemization of amino acids in mammoth collagen determined by gas and liquid chromatography. *Comparative Biochemistry and Physiology*, **53B**: 473–480.
- GOUDIE, A., 1972. Climate, weathering, crust formation, dunes, and fluvial features of the Central Namib Desert, near Gobabeb, South West Africa. *Madoqua*, *Ser II*, **1**: 15–31.
- GOWLETT, J. A. J., J. W. K. HARRIS, D. WALTON and B. A. WOOD, 1981. Early archaeological sites, hominid remains and traces of fire from Chesowanja, Kenya. *Nature*, **294**: 125–129.
- HAY, R. L., 1976. Geology of the Olduvai Gorge. Berkeley, University of California Press.
- HOOKE, P. J. and J. A. MILLER, 1979. K-Ar dating of the Pleistocene fossil hominid site at Chesowanja, North Kenya. *Nature*, **282**: 710–712.
- IKEYA, M., 1983. ESR dating in anthropology. *Kagaku no Ryoiki*, **37**: 661–669. (In Japanese)
- IKEYA, M., 1985. Electron spin resonance. In RUTTER, N. W. (ed.) "Dating Methods of Pleistocene Deposits and Their Problems", Geoscience Canada, Reprint Series 2, Geological Association of Canada, 73–87.
- IKEYA, M. and K. OHMURA, 1983. Comparison of ESR ages of corals from marine terraces with ^{14}C and $^{230}\text{Th}/^{234}\text{U}$ ages. *Earth and Planetary Science Letters*, **65**: 34–38.
- ISAAC, G. LL., 1977. Olgorgesailie. Chicago, University of Chicago Press.
- ISAAC, G. LL., 1978. The Olgorgesailie Formation; stratigraphy, tectonics and the palaeogeographic context of the Middle Pleistocene archaeological sites. In BISHOP, W. W. (ed.) "Geological Background to Fossil Man", Edinburgh, Scottish Academic Press, 173–206.
- KOMURA, K. and M. SAKANOU, 1967. Studies on the dating methods for Quaternary samples by natural alpha-radioactive nuclides. *Science Reports of the Kanazawa University*, **12**: 21–66.
- MAGLIO, V. J., 1973. Origin and evolution of the Elephantidae. *Transactions of the American Philosophical Society, N. S.*, **63** (3): 1–149.
- MASTERS, P. M., 1982. An amino acid racemization chronology for Tabun. In RONEN, A. (ed.) "The Transition from Lower to Middle Palaeolithic and the Origin of Modern Man", BAR International Series 151, Oxford, B. A. R., 43–53.
- MATSU'URA, S., 1981. Determination of fluorine in fossil bone with an ion-selective electrode. *Bulletin of the National Science Museum, Tokyo, Series D (Anthropology)*, **7**: 7–16.
- MATSU'URA, S., 1982. A chronological framing for the Sangiran hominids; fundamental study by the fluorine dating method. *Bulletin of the National Science Museum, Tokyo, Series D (Anthropology)*, **8**: 1–53.
- MATSU'URA, S. and N. UETA, 1980. Fraction dependent variation of aspartic acid racemization age of fossil bone. *Nature*, **286**: 883–884.

- SHACKLEY, M., 1980. An Acheulean industry with *Elephas recki* fauna from Namib IV, South West Africa (Namibia). *Nature*, **284**: 340–341.
- SHACKLEY, M., 1985. Palaeolithic archaeology of the central Namib desert; a preliminary survey of chronology, typology and site location. *Cimbebasia* (State Museum, Windhoek), No. **6**: 1–84.
- TALLON, P. W. J., 1978. Geological setting of the hominid fossils and Acheulian artifacts from the Kapthurin Formation, Baringo District, Kenya. In BISHOP, W. W. (ed.) "Geological Background to Fossil Man", Edinburgh, Scottish Academic Press, 361–373.