

Neutron Activation Analysis of Japanese Standard Rock Samples III

By

Fumitaka WAKABAYASHI

Department of Science and Engineering,
National Science Museum, Tokyo.

Abstract

Non-destructive instrumental neutron activation analysis was carried out for determining four major and twelve trace elements in six new Japanese standards rock samples, JSd-1, JSd-2, JSd-3, JSI-1, JSI-2 and JCh-1, which were issued as a part of "sedimentary rock series" from the Geological Survey of Japan (GSJ) in 1988 and 1989.

1. Introduction

In previous reports (WAKABAYASHI, 1987 and 1988), nondestructive instrumental neutron activation analyses were carried out for eighteen Japanese geological standard rock samples (that is, fifteen "igneous rock series" samples and three "sedimentary rock series" samples) which were issued from the Geological Survey of Japan (GSJ) before 1987.

Six new "Sedimentary rock series" standards rocks, JSd-1, JSd-2, JSd-3, JSI-1, JSI-2 and JCh-1 were issued from GSJ in 1988 and 1989 (TERASHIMA *et al.*, 1989).

In this report, nondestructive instrumental neutron activation analysis results of these new sedimentary rock samples are reported.

2. Experimental

The experimental procedures are similar to those described before (WAKABAYASHI, 1987 and 1988). A brief explanation is made below.

Each samples was weighed accurately (about 10 mg) and irradiated for 10 minutes in JRR-4 reactor at a flux of 8×10^{13} neutrons/cm²·sec in the Japan Atomic Energy Research Institute (JAERI) at Tokai. Irradiated samples were cooled about 20 hours, and then produced γ -ray activities were measured by a Ge (Li) γ -ray spectrometer without any chemical treatment. For long-lived nuclides, measurements of γ -activities were carried out for several times at the intervals of 3 or more weeks.

JB-1, JG-1 and W-1 were used as standards, as well as Cr and Th solution standards in a similar way described before. The best values used as standards were summarized in the previous reports. These values were those compiled in 1986 for JB-1

Table 1. 1988 consensus values for JB-1 and JG-1 using as the standards in this work.

Element	JB-1	JG-1	Element	JB-1	JG-1
<i>Major elements (%)</i>					
Fe	6.27	1.50	K	1.19	3.30
Na	2.07	2.51	Mn	0.12	0.049
<i>Trace elements (ppm)</i>					
Ce	67	46.6	La	38	23
Co	38.7	4.0	Lu	0.31	0.46
Cs	1.19	10.2	Sc	27.4	6.5
Eu	1.52	0.76	Sm	5.0	5.1
Ga	18.1	17	Th	9.2	13.5
Hf	3.4	3.5	Zn	83	41.5

A. ANDO *et al.*, *Geochem. J.*, **23**, 143 (1989).

and JG-1 (ANDO *et al.*, 1987), and in 1972 for W-1 (FLANAGAN, 1973). As for JB-1 and JG-1, new data (1988 consensus values) were published recently (ANDO *et al.*, 1989). Most of the values were not changed from those summarized in the previous reports. But in this work, JG-1 was used as standards for elements which were not mentioned in the previous reports. So newly published 1988 consensus values for JB-1 and JG-1 used as standards in this work are summarized in Table 1. Values for W-1 are the same as those summerized before.

3. Results

Results are tabulated in Tables 2 and 3. Each values in the tables was obtained through averaging at least three data which had been measured from time to time. The quoted errors in the tables are the standard deviations.

Three samples of JSd-2 and JSI-2 were taken from the samples with same split and position numbers, respectively. Two of these three samples were irradiated at the same time, and the another was irradiated at different time. As for JSd-1 and JSI-1, two samples were taken from the samples with same split and position numbers, and irradiated at different time. All these results were averaged and are tabulated here.

On the other hand, JSd-1 and JSI-1 samples with different split and position numbers were irradiated and analyzed at the same time with this work. These results will be reported elsewhere.

Value for Cs of JSd-2 is not listed in Table 1 because of the big discrepancy between the analytical results derived from 604.5 keV γ -ray peak and 795.9 keV peak of ^{134}Cs ; 604.5 keV peak gave the result 6.1 ± 0.1 ppm, and 795.9 keV peak gave 1.19 ± 0.08 ppm. And these values are changing with time to time. This discrepancy can be attributed to unknown impurity nuclides whose γ -ray peaks are positioned at very similar energy region with ^{134}Cs 's Peaks. Extended measurements are required to confirm this explanation and to derive correct value.

Table 2. Results of neutron activation analysis of standard rock samples JSd-1, JSd-2, and JSd-3

Element	JSd-1 This work	JSd-2 This work	JSd-2 This work
<i>Major elements (%)</i>			
Fe	3.69 $\pm$ 0.03	8.68 $\pm$ 0.04	3.35 $\pm$ 0.04
Na	1.73 $\pm$ 0.01	1.87 $\pm$ 0.01	0.369 $\pm$ 0.001
K	1.84 $\pm$ 0.03	0.98 $\pm$ 0.02	1.89 $\pm$ 0.03
Mn	0.079 $\pm$ 0.002	0.103 $\pm$ 0.002	0.131 $\pm$ 0.003
<i>Trace elements (ppm)</i>			
Ce	34.9 $\pm$ 0.9	30.2 $\pm$ 0.6	50.6 $\pm$ 1.7
Co	11.5 $\pm$ 0.2	53.2 $\pm$ 0.4	14.0 $\pm$ 0.3
Cs	2.11 $\pm$ 0.07	—	28.4 $\pm$ 0.7
Eu	1.06 $\pm$ 0.02	1.09 $\pm$ 0.01	0.90 $\pm$ 0.02
Ga	19.6 $\pm$ 2.0	15.0 $\pm$ 1.3	13.2 $\pm$ 1.6
Hf	2.74 $\pm$ 0.08	1.97 $\pm$ 0.06	3.3 $\pm$ 0.2
La	17.4 $\pm$ 0.8	13.9 $\pm$ 0.5	23.7 $\pm$ 0.9
Lu	0.24 $\pm$ 0.03	0.26 $\pm$ 0.03	0.23 $\pm$ 0.03
Sc	10.9 $\pm$ 0.1	19.6 $\pm$ 0.1	11.8 $\pm$ 0.1
Sm	4.2 $\pm$ 0.3	3.1 $\pm$ 0.2	3.9 $\pm$ 0.3
Th	4.3 $\pm$ 0.2	5.0 $\pm$ 0.1	6.2 $\pm$ 0.2
Zn	75 $\pm$ 4	1206 $\pm$ 39	87 $\pm$ 6
JSd-1	Stream sediment. Stream sediment (Arkose sandstone), Irishiken river, Hitachi, Ibaraki Prefecture. split 2, position 58.		
JSd-2	Stream sediment. East side of Ibaraki Prefecture. split 5, position 16.		
JSd-3	Stream sediment. Stream sediment near Takatori Mine (tungsten), Katsuramura, Ibaraki Prefecture. split 9, position 48.		

4. Acknowledgments

The author wishes to acknowledge Dr. Atsushi ANDO of the Geological Survey of Japan, who has provided me the standard rock samples used in this work through Dr. Masako SHIMA of the National Science Museum.

A part of this work was supported by 1988 and 1989 National Universities' Program for the Common Use of JAERI Facilities, the Ministry of Education, Science and Culture (Nos. 8818 and 89017).

The author expresses his sincere gratitude to Professor Yasuo ITO, Mr. Takemi TAKANO, Mr. SAWAHATA and other personnel of the Tokai Branch, Research Center for Nuclear Science and Technology, the University of Tokyo, and reactor staff of the Japan Atomic Energy Research Institute at Tokai, for their assistance to this work.

Appreciation are also expressed to Dr. Akihiko OKADA of the Institute of Physical and Chemical Research, for the safekeeping of irradiated damples in his Institution.

Table 3. Results of neutron activation analysis of standard rock samples
JSI-1, JSI-2, and JCh-1

Element	JSI-1 This work	JSI-2 This work	JCh-1 This work
<i>Major elements (%)</i>			
Fe	4.76 $\pm$ 0.03	4.59 $\pm$ 0.03	0.274 $\pm$ 0.004
Na	1.68 $\pm$ 0.01	0.992 $\pm$ 0.001	0.0419 $\pm$ 0.0002
K	2.46 $\pm$ 0.03	2.42 $\pm$ 0.02	0.220 $\pm$ 0.004
Mn	0.051 $\pm$ 0.002	0.0688 $\pm$ 0.0010	0.018 $\pm$ 0.006
<i>Trace elements (ppm)</i>			
Ce	58.4 $\pm$ 1.2	65.6 $\pm$ 1.0	5.58 $\pm$ 0.18
Co	16.2 $\pm$ 0.2	16.0 $\pm$ 0.1	17.0 $\pm$ 0.2
Cs	7.8 $\pm$ 0.2	8.7 $\pm$ 0.1	0.30 $\pm$ 0.02
Eu	1.33 $\pm$ 0.02	1.29 $\pm$ 0.01	0.076 $\pm$ 0.02
Ga	18.4 $\pm$ 1.4	16.7 $\pm$ 1.1	1.5 $\pm$ 0.2
Hf	3.37 $\pm$ 0.10	3.53 $\pm$ 0.07	0.25 $\pm$ 0.02
La	29.0 $\pm$ 0.6	32.0 $\pm$ 0.5	1.9 $\pm$ 0.1
Lu	0.40 $\pm$ 0.03	0.36 $\pm$ 0.02	0.028 $\pm$ 0.002
Sc	16.8 $\pm$ 0.1	16.9 $\pm$ 0.1	1.03 $\pm$ 0.01
Sm	6.3 $\pm$ 0.3	6.5 $\pm$ 0.2	0.42 $\pm$ 0.04
Th	8.5 $\pm$ 0.2	10.0 $\pm$ 0.2	0.49 $\pm$ 0.02
Zn	84 $\pm$ 4	83 $\pm$ 3	7.1 $\pm$ 0.6
JSI-1 Slate. Toyama slate, Permian, Toyama-machi, Miyagi Prefecture. split 4, position 81.			
JSI-2 Slate. Toyama slate, Permian, Ogatsu-cho, Miyagi Prefecture. split 2, position 92.			
JCh-1 Chert. Ashio chert, Permian to middle Mesozoic, Ashikaga, Tochigi Prefecture. split 2, position 5.			

Finally the author is grateful to Dr. M. SHIMA for reading this manuscript and helpful discussions.

References

- ANDO, A., N. MITA, and S. TERASHIMA, 1987. 1986 values for fifteen GSJ rock reference samples, "Igneous rock series". *Geostandards Newslett.*, **11**: 159–166.
- ANDO, A., H. KAMIOKA, S. TERASHIMA and S. ITOH, 1989. 1988 values for GSJ rock reference samples, "Igneous rock series". *Geochem. J.*, **23**, 143–148.
- FLANAGAN, F. J., 1973. 1972 values for international geochemical reference samples. *Geochim. Cosmochim. Acta*, **37**: 1189–1200.
- TERASHIMA, S., 1989. Elemental composition of GSJ rock reference samples, "Sedimentary rock series". The 1989 annual meeting of the Geochemical Society of Japan, 3B10 (in Japanese).
- WAKABAYASHI, F., 1987. Determination of major and trace elements in nine Japanese geochemical standard rock samples by instrumental neutron activation analysis. *Bull. Natn. Sci. Mus., Tokyo, Ser. E*, **10**: 13–19.
- WAKABAYASHI, F., 1988. Neutron activation analysis of Japanese Standard Rock Samples II. *Bull. Natn. Sci. Mus., Tokyo, Ser.*, **11**: 9–16.