

A comparison of phytoplankton communities between the ancient Lakes Biwa and Baikal

N.A. Bondarenko^{1,*}, A. Tuji² & M. Nakanishi³

¹*Limnological Institute of Siberian Branch of Russian Academy of Sciences, Irkutsk, Russia*

²*Department of Botany, Independent Administrative Institution, National Science Museum, Amakubo, Tsukuba, Japan*

³*Research Institute for Humanity and Nature, Kyoto, Japan*

(*Author for correspondence: E-mail: evs@lin.irk.ru)

Key words: lake Biwa, lake Baikal, phytoplankton community, taxonomic structure, comparative analysis

Abstract

As a result of comparative analysis of phytoplankton population structures between the ancient lakes Biwa and Baikal, we found floristic similarities and differences; but the former were minor. Both lakes had highly original algal floras due to presence of endemic, relict and rare forms. The richest in species number in the lakes were green algae and diatoms; *Scenedesmus*, the most diverse genus in Chlorococcales and *Cyclotella*, *Stephanodiscus*, *Aulacoseira* and *Synedra* in diatoms. The chief distinction between the lakes was that desmids appeared abundantly in the pelagic zone of Lake Biwa, whereas they were observed only in shallows and bays in Lake Baikal. The Sorensen's coefficient calculated for the planktonic floras was very low. This suggests that the algal floras of these lakes are significantly diverse.

Introduction

Lake Biwa, located in the central part of Japan, is one of the oldest lakes in the world (2–5 million years). The lake being 680 km² in the surface area consists of two basins, the north basin, oligo-mesotrophic (mean depth 44 m, maximum 104 m) and the south basin, eutrophic (mean depth 3 m). It is a characteristic of Lake Biwa that there are many endemic species including six species of fishes and 20 of molluscs. However, the ecosystem structure seems to have changed since the 1960's through progressive eutrophication, chemical pollution, physical alteration of habitats and invasion of alien organisms.

Lake Baikal is located in East Siberia, in an active rift zone. The lake is the oldest, ca. 25–30 million years, and deepest, 1637 m deep, in the world. It consists of three basins. The water is low mineralised, about 100 mg l⁻¹ total ions (Votint-

sev, 1961). Baikal is not only a natural laboratory for understanding evolutionary processes of lacustrine systems but also practically important as a huge reservoir of ca. 20% of the Earth's freshwater. The lake is rich in its biodiversity including 2500 species of animals, 82% endemic (Timoshkin, 1995), and about 1000 species of plants, 30% endemic.

The purpose of the present work is to make a comparison between the phytoplankton communities of the ancient Asian lakes Biwa and Baikal from a taxonomic point of view.

Results

Lake Biwa

Taxonomic studies on phytoplankton in Lake Biwa have been developed by late Professor Neg-

oro. Negoro (1968, 1971) and Ichise & Wakabayashi (1982) described many species of planktonic algae through their regular surveys in Lake Biwa. Mori (1980) and Mori & Miura (1980, 1990) compiled algal species in the fauna and flora lists on the basis of data by Negoro. Recently, much taxonomic information on European and American phytoplankton species has been introduced (Krammer & Lange-Bertalot, 1991; Patrick & Reimer, 1966, 1975) and made it possible to reassess the taxonomic positions of phytoplankton taxa in Lake Biwa. As a result, many new records have been described from Biwa. Ito (1988) reported 39 species of chrysophytes as new records – only four taxa in Mori & Miura (1980). Tuji & Houki (2001) reported nine centric taxa in diatoms as new records and ten taxa as misidentification – 13 taxa in Mori & Miura (1980). Shiga prefecture (1995, 2000) also reported many new records.

Tuji et al. (in preparation) and Tuji (in preparation) compiled a new flora list of planktonic algae on the basis of these recent studies. To understand fully the biodiversity of the Biwa phytoplankton community further, taxonomic work is required.

After Tuji et al. (in preparation), phytoplankton in Lake Biwa consists of 574 taxa, including 276 taxa of green algae, 91 of diatoms, 89 of chrysophytes and 63 of blue-green. Most of the taxa are cosmopolitan. Only three species of Bacillariophyta, namely *Stephanodiscus suzukii* Tuji et Kociolek, *Stephanodiscus pseudosuzukii* Tuji et Kociolek and *Aulacoseira nipponica* (Skvortzow) Tuji, are regarded as endemic (Tuji & Kociolek, 2000; Tuji, in preparation). *S. suzukii*, *S. pseudosuzukii* and *A. nipponica* have been identified previously as *S. carconensis* Grunow var. *carconensis*, *S. carconensis* var. *pusilla* Grunow and *Melosira solida* Eulenstein (*Aulacoseira solida* (Eulenstein) Krammer) respectively. These endemic taxa are very similar to those from western United States, and seem to be sister taxa (Tuji & Kociolek, 2000). These endemic diatoms are observed in the north basin of Lake Biwa in winter (ca. 7 °C) and inhabit oligotrophic, cold water (Tuji & Houki, 2001).

Pediastrum biwae Negoro and its variety were described as endemic species, but Barrientos (1979) modified these taxa as synonyms of *P. simplex*

Meyen. Though other several taxa have been described also as endemic, taxonomic positions of these are uncertain. *Staurostrum planctonicum* Teiling and *Staurostrum pseudosebaldii* Wille, which have been identified as *S. dorsidentiferum* W.-G.S. West var. *ornatum* (Gontcharov et al., 1999), are very important and unique to Lake Biwa. These taxa are observed abundantly from late summer to autumn in the pelagic zone of the north basin of Lake Biwa. Though many other species of desmids are found, their habits are restricted to the littoral zone.

Phytoplankton flora in Lake Biwa has changed somewhat since the 1970's. A "red tide" by sudden propagation of *Uroglena americana* Calkins occurred in 1977 and has been observed annually ever since (Nakanishi, 1984; Nakanishi & Sekino, 1996). Further, populations of *P. biwae* (*P. simplex*) and *M. solida* (*Aulacoseira nipponica*) tended to decrease around 1985, while *Anabaena* spp. and *Microcystis* spp. increased (Nakanishi & Sekino, 1996). Recently, *S. suzukii* has been decreasing in abundance, too. *A. nipponica* and *S. suzukii* are regarded as endangered species (Tuji & Houki, 2001).

Lake Baikal

Lake Baikal phytoplankton have been intensively studied during the 20th century (Meyer, 1930; Yasnitsky, 1930; Yasnitsky & Skabichevsky, 1957; Kozhova, 1959, 1987; Antipova, 1963; 1974; Votintsev et al., 1975; Popovskaya, 1987; and others). The last systematic list of baikalian planktonic algae numbers 299 species and subspecies belonging to 111 genera, 56 families and eight phyla (Bondarenko, 1995). In species diversity, green algae are most numerous, having 112 taxa, blue-green 66, diatoms 63, and chrysophytes 37. The phytoplankton differ in species numbers between the pelagic and shallow zones of Lake Baikal, pelagic species numbering only 158. However, pelagic diatoms are the most numerous, comprising 34% of the total algal composition, but no representatives of Euglenophyta and Xanthophyta have been observed. Cryptophytes are represented by the same species both in pelagic and shallow waters. The two epiplanktonic algae *Chrysosphaera melosirae* K. Meyer of chryso-

Table 1. Phytoplankton taxonomic structure of Lakes Biwa and Baikal

Phylum, class, order	Species and subspecies number		Number of common taxa
	Lake Baikal	Lake Biwa*	
Cyanophyta	66	63	20
Chrysophyta	37	89	19
Cryptophyta	8	14	5
Dinophyta	9	19	2
Raphydophyta	—	1	—
Euglenophyta	3	16	2
Xanthophyta	1	5	—
Bacillariophyta:	63	91	21
Centrophyceae	43	32	11
Pennatophyceae	20	59	10
Chlorophyta:	112	276	47
Volvocales	4	53	4
Tetrasporales	—	7	—
Chlorococcales	68	176	38
Zygnematales	—	2	—
Ulotrichales	5	5	2
Desmidiaceae	35	32	3
Oedogoniales	—	1	—
Total	299	574	116

* for Lake Biwa were accounted also attached forms being abundant in plankton.

phytes and diatom *Synedra cyclopum* Brutschy inhabit the plankton.

In the Lake Baikal phytoplankton, 13 of the 56 families are rich in species and subspecies. They number 132 taxa, ranged lower than genus, that is 40% of the total list. The richest genera are 14 (of 111), which include 55% of the total species composition. Among these, most diverse are the 7 genera *Anabaena* Bory (13 species and subspecies), *Cyclotella* Kützing (14), *Stephanodiscus* Ehrenberg (9), *Aulacoseira* Thwaites (9), *Dinobryon* Ehrenberg (9), *Scenedesmus* Meyen (9), *Mallomonas* Perty (8). So, the Lake Baikal phytoplankton tend to concentrate species diversity into a limited number of genera, i.e. 14 round the lake and three within the pelagic zone, as well as into 13 families. At the same time, there are many genera and families, poor in species number. According to Skabichevsky (1974), species richness in a limited pool of genera and families, mentioned above, and the low species number in most genera and

families reflect complicated processes in their genesis. This trend suggests that allochthony was important in the formation of the Lake Baikal algal flora. An autochthonous origin was important in genesis of the genera *Anabaena*, *Dinobryon*, *Scenedesmus*, *Mallomonas*, *Aulacoseira*, *Cyclotella*, *Stephanodiscus*. Only among the last three, are there endemic taxa. Species of these listed genera, except *Scenedesmus* dwelling in shallows, dominate in the pelagic zone.

Most algae of Lake Baikal plankton are cosmopolitan (62%). Also there are 14% boreal forms, 3% arctic-alpine ones, and 15% of unidentified nature. Of the planktonic algae, 5% have been found to be endemic, being mostly numbered among diatoms by seven taxa, principally *Aulacoseira baicalensis* (K. Meyer) Simonsen, *Cyclotella baicalensis* Skvortzow, Meyer, *C. minuta* (Skvortzow) Antipova, *Stephanodiscus binderanus* var. *baicalensis* Popovskaya et Genkal. Except in the genus *Crateriportula* Flower, Hakansson, planktonic algae express their endemism at species and subspecies level. In Baikal phytoplankton, there are relict forms also, described as endemic but later found elsewhere; for example, *Gymnodinium coeruleum* Antipova. Among baikalian plankton, there are forms that are globally rare, namely the diatom *S. cyclopum*, the chrysophyte *Mallomonas vannigera* Asmund and the green alga *Pediastrum primum* (Printz) Hegew.

Comparative analyses

In comparing the phytoplankton communities of the ancient Asian lakes Baikal and Biwa, it is apparent both lakes have endemic, relict and rare forms. Endemic forms have been found in diatoms, dinoflagellates and chrysophytes in Baikal but only in diatoms in Biwa. The richest in species number are green algae and diatoms.

A difference between the lakes is that desmids, especially *S. planctonicum* and *S. pseudosebaldii*, appear abundantly in the open water of Biwa, while they are observed only in shallows and bays in Baikal. Volvocales have been found to similarly spread.

Though diatom numbers are higher in Lake Biwa (91 spp.) than in Baikal (63 spp.), most of small centric species, including *Stephanodiscus* spp.

and *Cyclotella* spp., are observed in the littoral zone, as are the potamoplankton in Biwa. Some of these species prefer brackish and/or eutrophic water. Another difference between the lakes is that 16 species of euglenophytes are described in Biwa but only three species in Baikal. This seems to reflect a difference in the trophic status of the lakes.

Species common to both the lakes number 116 (Table 1). The Sorensen's coefficient (Sorensen, 1948) is a low 27%, indicating that the algal floras of the lakes are more diverse than similar.

References

- Antipova, N. L., 1963. Seasonal and annual variations of phytoplankton in Lake Baikal. In Galaziy, G. I. (ed.), Proceedings of Limnological Institute. USSR Academy of Sciences publishers, Moscow, Leningrad, 2(22): 12–28.
- Antipova, N. L., 1974. Annual variations of phytoplankton in Lake Baikal in Bolshie Koty region during 1960–1970. In Kozhova, O. M. (ed.), Productivity and anthropogenic changes in Lake Baikal. Irkutsk state University Press, Irkutsk, 75–84.
- Barrientos, O. P., 1979. Revision der Gattung *Pediastrum* Meyen (Chlorophyta). Bibliotheca Phycologica, 48.
- Bondarenko, N. A., 1995. List of planktonic algae in Lake Baikal. In Timoshkin, O. A. (ed.), Guide and key to pelagic animals of Baikal. Nauka, Novosibirsk, 621–630.
- Data book of Lake Biwa plankton (phytoplankton) 1990–1993, 1995. The Shiga Prefecture Institute of Public Health Environmental Science, Shiga Prefecture, 109 pp.
- Data book of Lake Biwa plankton (phytoplankton) 1994–1998, 2000. The Shiga Prefecture Institute of Public Health Environmental Science, Shiga Prefecture, 51 pp.
- Gontcharov, A. A., F. Kasai & M. M. Watanabe, 1999. Morphology and taxonomy of two planktonic species of *Staurastrum* (Desmidiaceae, Chlorophyta) in Lake Biwa, Japan. Algological Studies 93: 79–90.
- Ichise, S. & T. Wakabayashi, 1982. Plankton of Lake Biwa. Shiga Prefecture. Institute of Public Health Environmental Science, Shiga Prefecture, 144 pp.
- Ito, H., 1988. Scale-bearing chrysophytes in the south basin of Lake Biwa, Japan. Japanese Journal of Phycology 36: 143–153.
- Kozhova, O. M., 1959. Systematic list of plankton algae of Lake Baikal and some data on the biology of their dominant forms. Issues Siberian Division of the Russian Academy of Sciences 10: 112–124.
- Kozhova, O. M., 1987. Phytoplankton of Lake Baikal: structural and functional characteristics. Archiv für Hydrobiologie Beiheft Ergebnisse Limnologie 25: 19–37.
- Lake Biwa Study Monographs, 2001. Lake Biwa Research Institute, Otsu, 7: 89 pp.
- Krammer, K., H. Lange-Bertalot, 1991. Süsswasserflora von Mitteleuropa. Bacillariophyceae, 3. Teil: Centrales, Fragilariaceae, Eunotiaceae. VEB Gustav Fischer, Jena, 2: 576 pp.
- Meyer, K. I., 1930. Introduction to algal flora of Lake Baikal. Bull. MOIP, Novaya Seriya 39: 179–396.
- Mori, S., 1980. Fourth report of the regular limnological survey of Lake Biwa (1970) III. Phytoplankton. Memories of the Faculty of Science, Kyoto University. Series D, Biology 8: 37–57.
- Mori, S. & T. Miura, 1980. List of plant and animal species living in Lake Biwa. Memories of the Faculty of Science, Kyoto University. Series D, Biology 8: 1–34.
- Mori, S. & T. Miura, 1990. List of plant and animal species living in Lake Biwa (Corrected third edition). Memories of the Faculty Science, Kyoto University. Series D, Biology 14: 13–32.
- Nakanishi, M., 1984. Phytoplankton. In Horie, S. (ed.), Lake Biwa. Dr. W. Junk Publ. Dordrecht, 281–290.
- Nakanishi, M. & T. Sekino, 1996. Recent drastic changes in Lake Biwa bio-communities, with special attention to exploitation of the littoral zone. Geo Journal 40: 63–67.
- Negoro, K., 1967. First report of the regular limnological survey of Lake Biwa (Oct.1965- Dec. 1966) II. Plankton in gerald and phytoplankton. Memories of the Faculty of Science, Kyoto University. Series D Biology 1: 41–61.
- Negoro, K., 1968. The phytoplankton of Lake Biwa. In Kitamura, S. (ed.), Flora Ohmiensis. Hoikusha, 275–330.
- Negoro, K., 1971. The plankton of Lake Biwa. In Lake Biwa quasi-national park scientific survey group (ed.), Lake Biwa quasi-national park scientific survey report, 245–278.
- Patrick, R. M. & C. W. Reimer, 1966. The diatoms of the United States exclusive of Alaska and Hawaii. Monographs Series of Academy of Natural Sciences Philadelphia 1, 688 pp.
- Patrick, R. M. & C. W. Reimer, 1975. The diatoms of the United States exclusive of Alaska and Hawaii. Entomoneidaceae, Cymbellaceae, Gomphonemaceae, Epithemaceae. Monographs Series of Academy Natural Sciences Philadelphia 2, 213 pp.
- Plankton of Lake Biwa, 1982. Shiga Prefecture Institute of Public Health Environmental Science, Shiga Prefecture, 144 pp.
- Popovskaya, G. I., 1987. Phytoplankton of the deepest lake in the world. In Gutelmakher, B. L. (ed.), Marine and Freshwater Plankton. Nauka, Leningrad, 199–210.
- Skabichevsky, A. P., 1974. The algae intraspecies evolution and taxonomy problems. In Problems of lower plant phylogeny. Nauka, Moscow, 7–19.
- Sorensen, T., 1948. A method of establishing groups of equal amplitude in plant ecology. Biologiske Skrifter 5.
- Timoshkin, O. A. (ed.), 1995. Guide and key to pelagic animals of Baikal. Nauka Novosibirsk, 693.
- Tuji, A. & A. Houki, 2001. Centric diatoms in Lake Biwa. In Lake Biwa Study Monographs. Lake Biwa Research Institute. Shiga Prefecture 7: 1–90.
- Tuji, A. & J. P. Kociolek, 2000. Morphology and taxonomy of *Stephanodiscus suzukii* sp. nov. and *Stephanodiscus pseudosuzukii* sp. nov. (Bacillariophyceae) from Lake Biwa, Japan,

- and *S. carconensis* from North America. Phycologica Research 48: 231–239.
- Votintsev, K. K., 1961. Hydrochemistry of Lake Baikal. USSR Academic Sciences publishers, Moscow, 311 pp.
- Votintsev, K. K., A. I. Mesheryakova & G. I. Popovskaya, 1975. Turnover of organic matter in Lake Baikal. Nauka, Novosibirsk, 189 pp.
- Yasnitsky V. N., 1930. Results of the observations of Baikal plankton in the region of Biological station during 1926–1928. Proc. biol. geographic. Institute of Irkutsk State University Press, Irkutsk, 4: 191–234.
- Yasnitsky, V. N. & A. P. Skabichevsky, 1957. Baikal phytoplankton. In Galaziy, G.I. (ed.), Proceedings of the Limnological Station, the Russian Academy of Sciences Publishers, Moscow, Leningrad, 15: 212–261.