

Chevkinite-(Ce) in tuff at Heguri, Boso Peninsula, Chiba Prefecture, Japan

Ritsuro Miyawaki^{1*}, Satoshi Matsubara¹, Kazumi Yokoyama¹, Koichi Momma¹,
Takashi Sano¹, Yukiyasu Tsutsumi¹, Masako Shigeoka¹ and Katsumi Nishikubo²

¹Department of Geology and Paleontology, National Museum of Nature and Science,
4-1-1 Amakubo, Tsukuba, Ibaraki 305-0005, Japan

²Fukuei, Ichikawa, Chiba 272-0137, Japan

Abstract Chevkinite-(Ce) occurs in the rhyolitic tuff of the Mineoka group from Boso Peninsula, Chiba Prefecture, Japan. It forms as black tiny crystals less than 100 μm across in association with zircon, quartz, K-feldspar and analcime. Optically, chevkinite-(Ce) is pleochroic in brown and blackish brown. The electron microprobe analysis gave the following empirical formula: $(\text{Ce}_{1.57}\text{La}_{0.83}\text{Ca}_{0.58}\text{Nd}_{0.56}\text{Pr}_{0.15}\text{Y}_{0.09}\text{Th}_{0.08}\text{Sm}_{0.05}\text{Gd}_{0.04}\text{Dy}_{0.02})_{\Sigma 3.97}(\text{Ti}_{2.74}\text{Fe}_{1.84}\text{Al}_{0.14}\text{Mg}_{0.05}\text{Nb}_{0.05}\text{Mn}_{0.02})_{\Sigma 4.84}\text{Si}_4\text{O}_{21.25}\text{F}_{0.18}$ on the basis of $\text{Si} = 4$. It is non-metamict and gives sharp X-ray diffractions. The unit cell parameters were refined from data of a single crystal on a 4-circle diffractometer; $a = 13.3949(12)$, $b = 5.7293(8)$, $c = 11.0523(10) \text{ \AA}$, $\beta = 100.636(7)^\circ$, $V = 833.62(16) \text{ \AA}^3$ and $Z = 2$. The refinement of crystal structure converged to $R1 = 0.0371$ and $wR2 = 0.1304$ for 1984 independent reflections.

Key words: chevkinite-(Ce), crystal structure, Chiba, Japan, rare-earth

Introduction

Chevkinite-(Ce) is a silicate mineral of rare earth elements (REE), iron and titanium, and is a member of the chevkinite group minerals having general formula $A_4M_5\text{Si}_4\text{O}_{22}$, where A = larger cations such as REE, Sr, Ca, Th, and M = cations with octahedral coordination such as Ti, Nb, Ta, Zr, Fe, Mg, Mn, Cr. The members of chevkinite group can be classified into two groups with their crystal structures; the chevkinite-type structure in $C2/m$ and the perrierite-type structure in $P2_1/a$. The minor difference between two structures brings very similar powder X-ray diffraction patterns and makes it difficult to identify the mineral species with crystallographic data. An elaborate single crystal diffraction study is necessary to determine the crystal structure type, whereas chevkinite-(Ce) usually occurs in the metamict state giving no or poor X-ray diffraction data.

During our geological survey of the Mineoka group from Boso Peninsula, Chiba Prefecture, Japan, we recognized black fragments in the

zircon-bearing heavy sand sample for the dating partitioned from pale green tuff between the shale strata at Heguri, Chiba Prefecture ($35^\circ 6' 13''\text{N}$, $139^\circ 55' 44''\text{E}$). Our mineralogical investigation revealed that the fragments are non-metamict crystalline grains of chevkinite-(Ce).

Occurrence

Chevkinite-(Ce) occurs as euhedral black crystals less than 100 μm across in rhyolitic tuff in association with zircon, quartz, K-feldspar and analcime (Figs. 1 and 2). Monazite-(Ce) and fine particles of sulfides, such as galena and sphalerite, can be observed in addition to the associating minerals. The rhyolitic tuff consists of quartz, K-feldspar and volcanic glass. The almost parts of the glass were altered into analcime. Under the microscope, chevkinite-(Ce) is pleochroic in brown and blackish brown. More than 10 grains of chevkinite-(Ce) could be separated in the heavy liquid from approximate 150 g sample of




Fig. 1. The appearance of pale green rhyolitic tuff (a) and the close up view of black crystal of chevkinite-(Ce) of the rhyolitic tuff (b).

the rhyolitic tuff. The detailed geological background of chevkinite-(Ce) from rhyolitic tuff is given in Tsutsumi *et al.* (2012).

Chemical Composition

The chemical composition of chevkinite-(Ce) from Heguri was determined by means of a JEOL 8800 electron microprobe analyzer (WDS mode, 15kV, 20nA, and 2 μ m beam diameter). The analytical standard materials are as follows; REEP₅O₁₄ for REE (rare earth elements), ThO₂ for Th, anatase for Ti, fayalite for Fe and Si, sillimanite for Al, forsterite for Mg, columbite for Nb, rhodonite for Mn, CaF₂ for F. The result of chemical analysis with ZAF corrections is given in Table 1. The empirical formula derived from the chemical analysis is (Ce_{1.57}La_{0.83}Ca_{0.58}Nd_{0.56}Pr_{0.15}Y_{0.09}Th_{0.08}




Fig. 2. Photomicrographs [plane polarized light (PPL) (a) and crossed polars (CPL) (b)] and back scattered electron image (c) of chevkinite-(Ce) [brown in PL and pale gray in BSE] and zircon [colorless in PL and gray in BSE] in the rhyolitic tuff.

Sm_{0.05}Gd_{0.04}Dy_{0.02})_{Σ3.97}(Ti_{2.74}Fe_{1.84}Al_{0.14}Mg_{0.05}Nb_{0.05}Mn_{0.02})_{Σ4.84}Si₄O_{21.25}F_{0.18} on the basis of Si = 4.

Table 1. Chemical composition of chevkinite-(Ce) from Heguri.

CaO	2.68	Ca	0.58
Sc ₂ O ₃	0.02	Sc	0.00
ThO ₂	1.70	Th	0.08
Y ₂ O ₃	0.81	Y	0.09
La ₂ O ₃	11.24	La	0.83
Ce ₂ O ₃	21.31	Ce	1.57
Pr ₂ O ₃	2.05	Pr	0.15
Nd ₂ O ₃	7.75	Nd	0.56
Sm ₂ O ₃	0.79	Sm	0.05
Gd ₂ O ₃	0.57	Gd	0.04
Dy ₂ O ₃	0.30	Dy	0.02
		ΣA	3.97
TiO ₂	18.14	Ti	2.74
Nb ₂ O ₅	0.50	Nb	0.05
Ta ₂ O ₅	0.00	Ta	0.00
FeO	10.96	Fe	1.84
MnO	0.14	Mn	0.02
MgO	0.15	Mg	0.05
Al ₂ O ₃	0.58	Al	0.14
		$\Sigma M (BC)$	4.84
SiO ₂	19.92	Si	4
F	0.28	F	0.18
O = F	-0.12		
Total	99.77		

X-ray crystallography

Powder X-ray diffraction data of chevkinite-(Ce) from Heguri were obtained with a 114.6 mm Gandolfi camera. A fragment of chevkinite-(Ce) which was chemically analyzed was picked up from the thin section, and mounted on a glass fiber of 10 μ m diameter. The Debye-Scherrer ring images were recorded on an imaging plate (IP) in the Gandolfi camera using Ni-filtered CuK α radiation for a few days. Two exposures for measurement were carried out with and without Si standard material for X-ray diffraction angle calibration. The data were processed with a Fuji BAS-2500 bio-imaging analyser using a computer program written by Nakamuta (1999). The reflections were indexed according to the observed single crystal X-ray diffraction data and

Table 2. The powder X-ray diffraction data of chevkinite-(Ce) from Heguri.

<i>h</i>	<i>k</i>	<i>l</i>	<i>I</i> / <i>I</i> ₀	<i>d</i> _{obs.}	<i>d</i> _{calc.}	<i>h</i>	<i>k</i>	<i>l</i>	<i>I</i> / <i>I</i> ₀	<i>d</i> _{obs.}	<i>d</i> _{calc.}	<i>h</i>	<i>k</i>	<i>l</i>	<i>I</i> / <i>I</i> ₀	<i>d</i> _{obs.}	<i>d</i> _{calc.}
0	0	1	9	10.9	10.9	3	1	3	7	2.34	2.34				9 q	1.542	
2	0	0	6	6.58	6.59	4	0	4	12	2.31	2.32	0	2	6	5	1.528	1.530
0	0	2	23	5.43	5.43	3	1	4		2.31	2.31	3	1	6	8	1.517	1.517
2	0	1	5	5.21	5.22				6 q	2.28		3	1	7	11	1.501	1.502
1	1	1	23	4.87	4.87	0	2	3			2.25	4	2	6	5	1.476	1.479
1	1	1	35	4.60	4.60	4	0	3	18 q	2.24	2.24	2	2	6	5	1.442	1.441
			15 q	4.26		6	0	1			2.23	0	4	0	12	1.432	1.431
1	1	2	4	3.92	3.92	4	2	1	33	2.17	2.17	3	3	5	5	1.415	1.415
2	0	2	5	3.86	3.85	4	2	0			2.16	8	2	1	8	1.383	1.387
0	0	3	21	3.62	3.62				5 q	2.13		7	1	4		1.383	1.385
3	1	0		3.48		4	2	2	11	2.10	2.10	6	2	4	10 q	1.376	1.377
3	1	1	21	3.47	3.47	6	0	1	8	2.07	2.08	7	3	2	10	1.340	1.343
			76 q	3.34		4	2	1			2.07	7	3	0		1.340	1.340
3	1	1	61	3.18	3.18	3	1	4	17	2.00	2.00	4	4	2	5	1.298	1.298
3	1	2	49	3.15	3.15	3	1	5	33	1.971	1.978	4	4	1	7	1.291	1.292
1	1	3	33	3.08	3.09	0	2	4			1.970	8	2	5		1.292	1.292
4	0	2	33	3.08	3.08	2	2	4			1.961	6	2	5		1.271	1.271
4	0	1	23	3.00	3.00	1	1	5	25	1.958	1.956	0	4	4	11	1.268	1.266
2	0	3	7	2.95	2.95	1	3	1	6	1.853	1.852	6	2	7	9	1.253	1.254
1	1	3	18	2.88	2.88	2	2	4	16 q	1.819	1.821	0	2	8	4	1.226	1.226
0	2	0	25	2.87	2.86	7	1	2	5	1.799	1.794	3	3	6	5	1.214	1.214
3	1	2	50	2.76	2.75	4	2	3			1.763	3	1	8		1.206	1.206
3	1	3		2.72	2.72	6	2	1	10	1.762	1.761	3	3	7	11	1.205	1.206
0	0	4	100	2.72	2.71	6	2	2	19	1.734	1.735	3	1	9		1.196	1.196
2	2	0	10	2.62	2.62	4	0	6	20	1.729	1.727	4	2	7	7	1.196	1.193
4	0	2		2.61	2.61	6	0	5			1.711	11	1	1		1.189	1.189
2	2	1	12	2.61	2.60	3	3	1	20	1.707	1.708	7	1	8	8	1.185	1.186
0	2	2	16	2.53	2.53	3	3	2			1.703	7	1	6	5	1.173	1.173
2	2	1	9	2.51	2.51	6	2	1			1.682	7	3	4	6	1.143	1.143
1	1	4	4	2.49	2.49	4	0	5	18 q	1.676	1.676	8	0	6	5	1.118	1.119
			6 q	2.46		8	0	1			1.675	4	4	6	6	1.102	1.102
2	2	2			2.43	5	1	4	15	1.662	1.662	6	4	5			1.098
5	1	1	5	2.43	2.42	3	3	3	16	1.625	1.623	3	5	1	5	1.097	1.097
5	1	0	3	2.40	2.39	6	2	2	5	1.593	1.592						

q: quartz

 $a = 13.405(6)$, $b = 5.7242(15)$, $c = 11.050(3)$ Å, $\beta = 100.69(3)^\circ$, $V = 833.2(5)$ Å³, $Z = 2$.

calculated X-ray diffraction pattern simulated from the subsequent result of crystal structure analysis. The unit-cell parameters were refined from the powder X-ray diffraction data using the computer program of Toraya (1993) and are: $a = 13.405(6)$, $b = 5.7242(15)$, $c = 11.050(3)$ Å, $\beta = 100.69(3)^\circ$, $V = 833.2(5)$ Å³ and $Z = 2$. X-ray powder diffraction data are given in Table 2.

The X-ray diffraction intensity data of the single crystal selected for structure analysis were collected using a Rigaku RASA-7R four-circle diffractometer employing graphite-monochromatized MoK α radiation from a rotating anode (50 kV and 200 mA). Experimental details of the data collection are given in Table 3. Data reduction to F_o^2 with Lorentz and polarization corrections and correction for absorption (ϕ -scan pro-

cedure) were carried out with a computer program by Prof. Kazumasa Sugiyama of Tohoku University (pers. comm.). The SHELXL-97 software package (Sheldrick, 2008) was employed to refine the crystal structure. The scattering factors for the neutral atoms and anomalous dispersion factors were taken from the International Tables for X-ray Crystallography, Volume C (1992). Full-matrix least-squares refinement was performed by refining the positional parameters, scale factor, and displacement parameters.

The single crystal X-ray diffraction data showed the chevkinite-type C -lattice with extinction rules consistent with the space group $C2/m$. The atomic positional parameters of chevkinite-(Ce) "Crystal 1" (Sokolova *et al.*, 2004) were used for the starting model. Although chevkinite-

Table 3. Crystallographic data of chevkinite-(Ce) and experimental details.

Temperature	298 K
a (Å)	13.3949(12)
b (Å)	5.7293(8)
c (Å)	11.0523(10)
β (°)	100.636(7)
V (Å ³)	833.62(16)
Space group	$C2/m$
Z	2
Formula	(Ce _{0.83} Ca _{0.15} Y _{0.02}) ₄ Fe(Fe _{0.686} Al _{0.314}) ₂ Ti ₂ Si ₄ O ₂₂
D_{calc} (g/cm ³)	4.795
μ (cm ⁻¹)	12.587
Crystal dimension (mm)	0.15 × 0.07 × 0.03
Diffractometer	Rigaku AFC-7R
Radiation	MoK α (graphite-monochromatized)
Scan mode, rate (°/min in ω)	2θ - ω , 2
2θ range (deg.)	5–70
Reflection range	$-21 \leq h \leq 21$ $0 \leq k \leq 9$ $-17 \leq l \leq 8$
No. of measured reflections	2238
No. unique reflections	1984
No. of observed reflections [$I > 2\sigma(I)$]	1750
R_{int}	0.0158
No. of variable parameters	101
$R1$ [$I > 2\sigma(I)$], $R1$ (all reflections)	0.0296, 0.0371
$wR2$ (all reflections)	0.1304
Weighting parameters, a , b	0.1, 0
Goodness of fit	1.111
Final $\Delta\rho_{\text{min}}$ (e/Å ³)	–2.976
Final $\Delta\rho_{\text{max}}$ (e/Å ³)	1.432

$$R1 = \sum ||F_o| - |F_c|| / \sum |F_o|$$

$$wR2 = \{ \sum [w(F_o^2 - F_c^2)^2] / \sum [w(F_o^2)^2] \}^{0.5}$$

$$w = 1 / [\sigma^2(F_o^2) + (aP)^2 + bP]$$

$$P = [2F_c^2 + F_o^2] / 3$$

Table 4. Final atomic coordinates and displacement parameters ($\text{\AA}^2$).

		occupancy	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{eq}
Ce1	<i>A</i> 1	Ce _{0.83} Y _{0.02} Ca _{0.15}	0.35634(2)	0	0.73373(3)	0.01315(12)
Ce2	<i>A</i> 2	Ce _{0.83} Y _{0.02} Ca _{0.15}	0.06982(2)	0	0.74050(3)	0.01681(12)
Fe1	<i>B</i> (<i>M</i> 1)	Fe	0.5	0	0.5	0.00745(18)
Fe2	<i>C</i> 1(<i>M</i> 2)	Fe _{0.686(8)} Al _{0.314}	0.25	0.25	0	0.0071(2)
Ti3	<i>C</i> 2a(<i>M</i> 2)	Ti	0	0	0	0.0095(2)
Ti4	<i>C</i> 2b(<i>M</i> 3)	Ti	0.5	0	0	0.0096(2)
Si1		Si	0.20107(10)	0.5	0.73167(11)	0.0065(2)
Si2		Si	0.35743(10)	0.5	0.54576(12)	0.0078(2)
O1		O	0.47764(19)	0.2519(5)	0.8741(2)	0.0107(4)
O2		O	0.1462(3)	0	0.9777(4)	0.0100(6)
O3		O	0.1859(3)	0	0.5981(3)	0.0126(6)
O4		O	0.3477(3)	0	0.9906(4)	0.0106(6)
O5		O	0.4264(3)	0.2720(6)	0.5943(3)	0.0214(6)
O6		O	0.2284(2)	−0.7353(5)	0.8146(2)	0.0118(4)
O7		O	0.0845(3)	0.5	0.6708(4)	0.0257(11)
O8		O	0.2685(4)	0.5	0.6282(5)	0.0356(14)

	<i>U</i> ₁₁	<i>U</i> ₂₂	<i>U</i> ₃₃	<i>U</i> ₂₃	<i>U</i> ₁₃	<i>U</i> ₁₂
Ce1	0.01331(18)	0.01482(18)	0.01094(17)	0	0.00127(12)	0
Ce2	0.01025(17)	0.0306(2)	0.00995(18)	0	0.00271(12)	0
Fe1	0.0098(4)	0.0078(4)	0.0042(3)	0	−0.0001(3)	0
Fe2	0.0078(3)	0.0096(4)	0.0041(3)	0.0000(2)	0.0010(2)	0.0002(2)
Ti3	0.0098(5)	0.0120(5)	0.0074(4)	0	0.0031(4)	0
Ti4	0.0085(5)	0.0106(5)	0.0094(5)	0	0.0006(4)	0
Si1	0.0073(5)	0.0061(5)	0.0062(5)	0	0.0014(4)	0
Si2	0.0089(5)	0.0098(5)	0.0048(5)	0	0.0013(4)	0
O1	0.0094(10)	0.0130(11)	0.0102(9)	−0.0021(9)	0.0035(8)	−0.0026(8)
O2	0.0069(13)	0.0111(14)	0.0125(14)	0	0.0034(11)	0
O3	0.0178(16)	0.0139(16)	0.0055(13)	0	0.0000(12)	0
O4	0.0077(13)	0.0089(14)	0.0146(15)	0	0.0006(11)	0
O5	0.0271(15)	0.0236(15)	0.0127(11)	0.0060(11)	0.0015(11)	0.0153(12)
O6	0.0191(11)	0.0082(10)	0.0082(10)	0.0013(8)	0.0027(8)	0.0030(8)
O7	0.0092(16)	0.054(3)	0.0129(18)	0	−0.0017(13)	0
O8	0.028(3)	0.064(4)	0.021(2)	0	0.020(2)	0

(Ce) showed a complex chemistry of isomorphous substitutions, simplified occupancy parameters were employed in the initial cycles of the refinement calculations; Ce₄Fe₃Ti₂Si₄O₂₂ as the structural formula. Although Fe and Ti have the similar scattering properties, the former was adopted to the Fe1 (*B* or *M*1) and Fe2 (*C*1 or *M*2) sites, whereas the latter was employed for the Ti3 (*C*2, or *M*3 and *M*4) sites, in reference to the previous studies on the crystal structure of chevkinite-(Ce) (Yang *et al.*, 2002, 2007; Sokolova *et al.*, 2004). In the next stage of refinement cycles, the occupancy parameters for the *A* (REE) sites were fixed as Ce_{0.83}Y_{0.02}Ca_{0.15} according to the chemical composition. In the further refinement

cycles, Al, as the representative of Mg and Al, was introduced into the octahedral Fe1, Fe2, Ti3 and Ti4 sites to refine the occupancy parameters against to the Ti or Fe, the representative of Ti, Mn and Fe. The refined occupancy parameters indicated the partial (0.3) substitution of Al(Mg) for Fe only at the Fe2 site, whereas no substitution at the other octahedral sites. In the final cycles of refinements, the occupancy parameters of Al and Fe at the Fe2 site were refined. The final positional parameters and anisotropic displacement parameters with equivalent isotropic displacement parameters are given in Table 4. Selected interatomic distances are summarized in Table 5.

ing two silicate tetrahedra, Si_2O_7 . The sheets are parallel to the ab plane, and stack alternately in the c^* direction to form a three-dimensional framework. The resulted structural formula is therefore $(\text{Ce}_{0.83}\text{Ca}_{0.15}\text{Y}_{0.02})_4\text{Fe}(\text{Fe}_{0.686}\text{Al}_{0.314})_2\text{Ti}_2\text{Si}_4\text{O}_{22}$. An approximation of minor atoms into the structural formula leads to the formula $(\text{Ce}_{0.83}\text{Ca}_{0.15}\text{Y}_{0.02})_4(\text{Fe}_{0.98}\text{Mn}_{0.02})(\text{Fe}_{0.43}\text{Ti}_{0.40}\text{Al}_{0.07}\text{Mg}_{0.03}\square_{0.07})_2(\text{Ti}_{0.97}\text{Nb}_{0.03})_2\text{Si}_4\text{O}_{22}$, suggesting a fractional vacancy at the Fe2 site. Whereas the Ti and Fe are simply allocated into Ti3, Ti4 and Fe1 sites, respectively, in this approximation, a complex mixing of Fe and Ti is probable at the octahedral sites in chevkinite-(Ce) from Heguri, as the site population in the Fe-rich chevkinite-(Ce) from China assigned by Yang *et al.* (2007).

The present chevkinite-(Ce) is the second occurrence in Japan, following to the first occurrence from alkaline dike in Cape Ashizuri, Kochi Prefecture, Shikoku, Japan (Imaoka and Nakashima, 1994). Chevkinite-(Ce) occurs in pumice fragments and air-fall ash, as well as in alkali granite and syenite pegmatites. The occurrence mode at Heguri is similar to the former type. The X-ray fluorescence whole rock analysis (Table 2 in Tsutsumi *et al.*, 2012) indicated considerable abundances of Y (101 ppm) and Ce (144 ppm) in the chevkinite-(Ce) bearing pale green rhyolitic tuff from Heguri. These rare earth elements are considered to be concentrated into the crystals of chevkinite-(Ce) and the associating monazite-(Ce). The source magma of rhyolitic tuff should be as rich in the rare earth as the tuff. It is worth to note that such a REE-rich volcano once existed within or close to Japanese Islands, whereas the position of volcano can not be identified.

Acknowledgement

The authors are grateful to the staff of

Marusho Sekisan Co. for their assistance for our geological survey in their quarry area. Prof. Zhuming Yang and Dr. Stuart J. Mills are thanked for their helpful reviews. This study was supported by the Integrated Research Project in the National Museum of Nature and Science, "Research on the Earth's surface processes and biota in and near the Sea of Japan".

References

- Imaoka, T. and Nakashima, K. (1994) Chevkinite in syenites from Cape Ashizuri, Shikoku Island, Japan. *Neues Jahrbuch für Mineralogie — Monatshefte*, **8**, 358–366.
- Nakamuta, Y. (1999) Precise analysis of a very small mineral by an X-ray diffraction method. *Journal of the Mineralogical Society of Japan*, **28**, 117–121 (in Japanese with English abstract).
- Sheldrick, G. M. (2008) A short history of SHELX. *Acta Crystallographica*, **A64**, 112–122.
- Sokolova, E., Hawthorne, F. C., Della Ventura, G., and Kartashov, P. M. (2004) Chevkinite-(Ce): Crystal structure and the effect of moderate radiation-induced damage on site-occupancy refinement. *Canadian Mineralogist*, **42**, 1013–1025.
- Toraya, H. (1993) The determination of unit-cell parameters from Bragg reflection data using a standard reference material but without a calibration curve. *Journal of Applied Crystallography*, **26**, 583–590.
- Tsutsumi, Y., Horie, K., Sano, T., Miyawaki, R., Momma, K., Matsubara, S., Shigeoka, M. and Yokoyama, K. (2012) La-ICP-MS and SHRIMP ages of zircons in chevkinite and monazite tuffs from the Boso Peninsula, Central Japan. *Bulletin of the National Museum of Nature and Science, Ser. C*, **37**, 15–32.
- Yang, Z., Fleck, M., Smith, M., Tao, K., Song, R. and Zhang, P. (2002) The crystal structure of natural Fe-rich chevkinite-(Ce). *European Journal of Mineralogy*, **14**, 969–975.
- Yang, Z., Li, H., Liu, M. and Pertlik, F. (2007) The crystal chemistry of iron in non-metamict chevkinite-(Ce): Valence state and site occupation proportions. *Journal of Rare Earths*, **25**, 238–242.