



































































































































































































































































- Frisch, A., Ohmura, Y., Ertz, D. & Thor, G. 2015. *Inoderma* and related genera in Arthoniaceae with elevated white pruinose pycnidia or sporodochia. *Lichenologist* 47: 233–256.
- Frisch, A., Thor, G., Ertz, D. & Grube, M. 2014. The Arthonialean challenge: restructuring Arthoniaceae. *Taxon* 63: 727–744.
- Frisch, A., Thor, G., Moon, K. H. & Ohmura, Y. 2017. *Arthonia incarnata* (Arthoniaceae), a rare and poorly known old-growth forest lichen new to Asia. *Nordic Journal of Botany* 35: 587–594.
- Fryday, A. M. 2005a. The genus *Porpidia* in northern and western Europe, with special emphasis on collections from the British Isles. *Lichenologist* 37: 1–35.
- Fryday, A. M. 2005b. (1656) Proposal to reject the name *Lecidea flavocoerulescens* (lichenized Ascomycota, Porpidiaceae). *Taxon* 54: 181–182.
- Fryday, A. M., Printzenn, C. & Ekman, S. 2014. *Bryobilimbia*, a new generic name for *Lecidea hypnorum* and closely related species. *Lichenologist* 46: 25–37.
- Gagarina, L. V. 2015. *Coenogonium theae* comb. et stat. nov. (Lichenized Ascomycota). *Herzogia* 28: 794–797.
- Galloway, D. J. 1988. Studies in *Pseudocyphellaria* (lichens). I. The New Zealand species. *Bulletin of the British Museum (Natural History), Botany Series* 17: 1–267.
- Galloway, D. J., Kantvilas, G. & Elix, J. A. 2001. *Pseudocyphellaria*. In: McCarthy, P. M. (ed.): *Flora of Australia*. Vol. 58A, Lichens 3. ABRS/CSIRO Australia, Melbourne, pp. 47–77.
- Gaya, E., Högnabba, F., Holguin, Á., Molnar, K., Fernández-Brime, S., Stenroos, S., Arup, U., Søchting, U., van den Boom, P., Lücking, R., Sipman, H. J. M. & Lutzoni, F. 2012. Implementing a cumulative supermatrix approach for a comprehensive phylogenetic study of the Teloschistales (Pezizomycotina, Ascomycota). *Molecular Phylogenetics and Evolution* 63: 374–387.
- Gierl, C. & Kalb, K. 1993. Die Flechtengattung *Dibaeis*. Eine Übersicht über die rosafrüchtigen Arten von *Baeomyces* sens. lat. nebst Anmerkungen zu *Phyllobaeis* gen. nov. *Herzogia* 9: 593–645.
- Giralt, M., Paz-Bermúdez, G. & Elix, J. A. 2009. New data on *Sculptolumina japonica* (Physciaceae). *Bryologist* 112: 397–403.
- Gueidan, C. & Roux, C. 2007. *Verrucaria calciseda* DC. Néotypification, description et transfert dans le genre Bagliettoa. *Bulletin de la Société Linnéenne de Provence* 58: 181–194.
- Gueidan, C., Savić, S., Thüs, H., Roux, C., Keller, C., Tibell, L., Prieto, M., Heiðmarsson, S., Breuss, O., Orange, A., Fröberg, L., Amtoft Wynns, A., Navarro-Rosinés, P., Krzewicka, B., Pykälä, J., Grube, M. & Lutzoni, F. 2009. Generic classification of the Verrucariaceae (Ascomycota) based on molecular and morphological evidence: recent progress and remaining challenges. *Taxon* 58: 184–208.
- Hafellner, J. 1979. Karschia. Revision einer Sammelgattung an der Grenze von lichenisierten und nichlichenisierten Ascomyceten. Beiheft zur Nova Hedwigia 62: 1–248.
- Hafellner, J. 1993. Über Funde von lichenicolen Pilzen und Flechten im südlichen Norwegen. *Herzogia* 9: 749–768.
- Hafellner, J. & Türk, R. 2001. Die lichenisierten Pilze Österreichs: eine Checkliste der bisher nachgewiesenen Arten mit Verbreitungsangaben. *Stapfia* 76: 3–167.
- Hafellner, J. & Türk, R. 2016. Die lichenisierten Pilze Österreichs – eine neue checkliste der bisher nachgewiesenen taxa mitangaben zu verbreitung und substratökologie. *Stapfia* 104: 1–216.
- Halonen, P., Myllys, L., Velmala, S. & Hyvärinen, H. 2009. *Gowardia* (Parmeliaceae)—a new alectorioid lichen

- genus with two species. *Bryologist* 112: 138–146.
- Harada, H. 2005. *Collema limosum* (lichenized Ascomycota; Collemataceae) as new for Japan. *Lichenology* 4: 113–115.
- Harada, H. 2008. *Gyalideopsis chibaensis* (lichenized Ascomycota, Gomphillaceae), a new gyalectoid lichen from central Japan. *Lichenology* 7: 25–29.
- Harada, H. 2010. *Leptogium chibaense* (lichenized Ascomycota, Collemataceae), a new small cyanolichen from central Japan. *Lichenology* 9: 19–22.
- Harada, H. 2011a. *Coccotrema pocillarium* (lichenized Ascomycota, Coccotremataceae) occurs in Japan. *Lichenology* 9: 39–43.
- Harada, H. 2011b. *Leptogium bosoense* (lichenized Ascomycota, Collemataceae), a new terricolous cyanolichen from Chiba-ken, central Japan. *Lichenology* 10: 29–32.
- Harada H. 2011c. Taxonomic study on the freshwater species of Verrucariaceae of Japan (1). *Verrucaria praetermissa* and *V. yoshimurae* sp. nov. *Lichenology* 10: 33–41.
- Harada, H. 2012. Taxonomic study on the freshwater species of Verrucariaceae of Japan (2). Genus *Verrucaria*. *Lichenology* 10: 97–135.
- Harada, H. 2013a. *Agonimia deguchii* (lichenized Ascomycota, Verrucariaceae), a new saxicolous species from central Japan. *Hikobia* 16: 307–310.
- Harada, H. 2013b. The lichen genus *Thelidium* (Verrucariaceae) in Japan. *Lichenology* 11: 53–66.
- Harada, H. 2014. *Cresporhaphis chibaensis* sp. nov. (lichenized Ascomycota, Trichosphaeriaceae) from Chiba-ken, central Japan. *Lichenology* 12: 31–36.
- Harada, H. 2015a. *Leptogium rivale* (lichenized Ascomycota, Collemataceae), an aquatic freshwater species, newly found in Japan. *Lichenology* 13: 45–51.
- Harada, H. 2015b. *Thelocarpon epibolum* (Ascomycota, Thelocarpaceae) newly found in Japan. *Lichenology* 13: 53–57.
- Harada, H. 2015c. Saxicolous and corticolous species of *Porina* (lichenized Ascomycota, Porinaceae) of Japan (part 1). *Lichenology* 14: 1–20.
- Harada, H. 2016a. Saxicolous species of the genus *Strigula* (lichenized Ascomycota, Strigulaceae) in Japan. *Lichenology* 14: 73–89.
- Harada, H. 2016b. Saxicolous and corticolous species of *Porina* (Lichenized Ascomycota, Porinaceae) of Japan (Part 2). *Lichenology* 14: 91–118.
- Harada, H. 2016c. *Gyalidea izuensis* sp. nov. (lichenized Ascomycota, Solorinellaceae), a semi-aquatic freshwater species from Shizuoka-ken, central Japan. *Lichenology* 15: 85–90.
- Harada, H. 2017a. Taxonomic notes on pyrenocarpus lichens in Japan (8). *Trypetheliopsis yoshimurae* (lichenized Ascomycota, Monoblastiaceae) sp. nov. *Lichenology* 16: 15–21.
- Harada, H. 2017b. *Leptogium kiyosumiense* (lichenized Ascomycota, Collemataceae), a new species of the Mallotium-group from Chiba-ken, Central Japan. *Lichenology* 16: 23–30.
- Harada, H. 2017c. Taxonomic notes on pyrenocarpus lichens in Japan (7). *Anisomeridium throwerae* (Monoblastiaceae) new to Japan. *Lichenology* 16: 69–73.
- Harada, H. 2017d. *Sarcogyne regularis* (lichenized Ascomycota; Acarosporaceae) new to Japan. *Lichenology* 16: 81–85.

- Harada, H. 2017e. *Caloplaca cerina* var. *chloroleuca* (lichenized Ascomycota; Teloschistaceae) new to Japan. Lichenology 16: 87–90.
- Harada, H. & Kawakami, S. 2011. *Gyalideopsis lunata* (Lichenized Ascomycota, Gomphillaceae), a new gyalectoid lichen from Gifu-ken, central Japan. Lichenology 10: 23–28.
- Harada, H. & Sakata, A. 2016. *Gyalidea oosumiensis* sp. nov. (lichenized Ascomycota, Solorinellaceae) from Kagoshima-ken, Kyushu, southwestern Japan. Lichenology 15: 1–5.
- Harada, H. & Sato, H. 2008. Contributions to the lichen flora of the Yamaru area, Okinawa, Ryukyu Is., Southwestern Japan (1). Genus *Porina*. Lichenology 7: 131–142.
- Harada, H. & Yoshikawa, H. 2017. *Psoroma tenue* var. *boreale* (lichenized Ascomycota; Pannariaceae) new to Japan. Lichenology 16: 75–80.
- Harada, H., Kinoshita, Y. & Sato H. 2007. *Dibaeis soreliata* (lichenized Ascomycota; Icmadophilaceae) as new for Japan. Lichenology 6: 11–14.
- Harada, H., Okamoto, T. & Yoshimura, I. 2004. A checklist of lichens and lichen-allies of Japan. Lichenology 2: 47–165.
- Harada, H., Wang, L. -S. & Yoshimura, I. 2012a. Lichen flora of Japan (2). Genus *Bryoria* (Parmeliaceae). Lichenology 10: 147–168.
- Harada, H., Yamamoto, Y. & Yoshimura, I. 2012b. *Myelochroa xantholepis* (lichenized Ascomycota, Parmeliaceae), newly found in Japan. Lichenology 11: 13–15.
- Harada, H., Yoshikawa, H., Izumi, H. & Sakata, A. 2016. Taxonomic notes on pyrenocarpous lichens in Japan (6). *Polymeridium proponens* (Trypetheliaceae) new to Japan. Lichenology 15: 91–97.
- Hawksworth, D. L. 1972. Regional studies in *Alectoria* (Lichenes) II. The British species. Lichenologist 5: 181–261.
- Hawksworth, D. L. 1975. Notes on British lichenicolous fungi, I. Kew Bulletin 30: 183–203.
- Hawksworth, D. L. 2004. Rediscovery of the original material of Osbeck's *Lichen chinensis* and the re-instatement of the name *Parmotrema perlatus* (Parmeliaceae). Herzogia 17: 37–44.
- Hawksworth, D. L. & Cole, M. S. 2003. A first checklist of lichenicolous fungi from China. Mycosystema 22: 359–363.
- Higashi, A., Yoshikawa, H., Watanuki, O. & Harada, H. 2017. Marine and maritime lichens of Japan (1). *Buellia yoshimurae* sp. nov. Lichenology 16: 1–13.
- Hue, A. 1899. Lichenes extra-europaei a pluribus collectoribus ad Museum Parisiensi missi. Nouvelles Archives du Muséum d'Histoire Naturelle de Paris, sér. 4, 1: 27–220.
- Hue, A. 1901. Lichenes extra-europaei a pluribus collectoribus ad Museum Parisiensi missi. Nouvelles Archives du Muséum d'Histoire Naturelle de Paris, sér. 4, 3: 21–196, 18 pls.
- Inoue, M. 1981. A taxonomic study of the Japanese species of *Fuscidea* (lichens). Hikobia Supplement 1: 161–176.
- Inoue, M. 1982. The genera *Lecidea*, *Lecidella* and *Huilia* in Japan I. *Lecidea*. Journal of Science of the Hiroshima University, Series B, Division 2 (Botany) 18: 1–55.
- Inoue, M. 2001. Taxonomic notes on *Rhizocarpon* growing in solfatara fields in Japan. Bulletin of the National Science Museum, Tokyo, Series B (Botany) 27: 11–21.
- Inoue, M. 2010. Notes on four lecideoid lichens new to Japan. Memoirs of the Faculty of Education and Human

- Studies. Akita University (Natural Science) 65: 17–21.
- Inoue, M. & Kanda, H. 2009. Distribution patterns of Japanese lecideoid lichens occurring in subalpine or alpine regions, with special reference to the taxa being distributed in both Japan and Svalbard. *Memoirs of the Faculty of Education and Human Studies Akita University (Natural Science)* 64: 1–8.
- Inoue, M., Kashiwadani, H. & Moon, K. H. 2007. Alpine lecideoid lichens from southern part of Mts. Akaishi, central Japan. *Memoirs of the Faculty of Education and Human Studies Akita University (Natural Science)* 62: 9–17.
- Inoue, M., Kashiwadani, H., Miyawaki, H., Nakanishi, M. & Ohmura, Y. 2010. Common names of Japanese lichen genera. *Lichen* 16: 9–13. (In Japanese).
- Inumaru, S. 1943. *Studia Lichenum Japoniae. III. Peltigeraceae Japoniae III. Acta phytotaxonomica et geobotanica* (Kyoto). *Acta Phytotaxonomica et Geobotanica* 12: 1–16.
- Jørgensen, P. M. 1975. Contributions to a monograph of the Mallotium-hairy *Leptogium* species. *Herzogia* 3: 433–460.
- Jørgensen, P. M. 2007. Lichinaceae. In: *Nordic lichen flora. Volume 3. Cyanolichens*. Uddevalla: Nordic Lichen Society.
- Jørgensen, P. M. 2008. *Vahliella*, a new lichen genus. *Lichenologist*, 40: 221–225.
- Jørgensen, P. M. 2010. New discoveries in the lichen genus *Staurolemma* Körber. *Nova Hedwigia* 90: 153–159.
- Jørgensen, P. M. & Kashiwadani, H. 2001. Notes on the lichen genus *Parmeliella* in Japan. *Journal of Japanese Botany* 76: 288–296.
- Joshi, Y., Thüs, H. & Hur, J. S. 2011. *Caloplaca aequata* is a synonym of *C. cinnabarina* (Teloschistaceae). *Lichenologist* 43: 141–146.
- Kalb, K., Staiger, B., Elix, J. A., Lange, U. & Lumbsch, H. T. 2008. A new circumscription of the genus *Ramboldia* (Lecanoraceae, Ascomycota) based on morphological and molecular evidence. *Nova Hedwigia* 86: 23–42.
- Kantvilas, G. & Elix, J. A. 1994. *Ramboldia*, a new genus in the lichen family Lecanoraceae. *Bryologist* 97: 296–304.
- Kashiwadani, H. 2000. *Lichenes minus cogniti exsiccati, fasc. VII (nos. 151–175)*. National Science Museum, Tokyo. 4 pp.
- Kashiwadani, H. & Moon, K. H. 2007. The genus *Ramalina* (Ascomycota, Ramalinaceae) in Indonesia. *Bibliotheca Lichenologica* 96: 145–156.
- Kashiwadani, H. & Moon, K. H. 2017. Two new species of *Ramalina* (Ascomycotina: Ramalinaceae) from the Nansei Islands, Japan. *Journal of Japanese Botany* 92: 27–33.
- Kashiwadani, H., Aptroot, A. & Moon, K. H. 2009a. New pyrenocarpous lichens in Japan, with the resurrection of the genus *Trypetheliopsis* for *Musaespora*. *Bibliotheca Lichenologica* 99: 247–258.
- Kashiwadani, H., Kobayashi, K. & Moon, K. H. 2009b. New or noteworthy species of lichens from Shiga Prefecture, central Japan. *Journal of Japanese Botany* 84: 255–258.
- Kashiwadani, H., Nakanishi, M. & Moon, K. H. 2014. Two new species of *Graphis* and *Halegrapha* (Graphidaceae, Ostropales) from southern Japan. *Journal of Japanese Botany* 89: 12–16.
- Kashiwadani, H., Aptroot A., Nakanishi, M., Moon, K. H., Inoue, M. & Sasaki, O. 2015. New or noteworthy species of lichens found in Akita Prefecture, Japan. *Journal of Japanese Botany* 90: 227–234.

- Kawakami, H., Takahashi, K., Mizoguchi, Y., Yamamoto, A. & Yamamoto, Y. 2012. Lichens in Kiifudoki-no-oka, Wakayama city, Honshu, Japan. *Nanki Seibutu* 54: 8–12.
- Kawakami, S., Tojo, B. & Harada, H. 2011. Contributions to the lichen flora of Gifu Prefecture, central Japan. (1). *Gyalecta jenensis* (Gyalectaceae) as new for Japan. *Lichenology* 10: 43–46.
- Kawakami, S., Tojo, B., Harada, H. & Okamoto, T. 2012a. *Caloplaca decipiens* (Teloschistaceae) as new for Japan. *Lichenology* 11: 17–20.
- Kawakami, S., Saito, Y., Hibino, H., Tojo, B. & Harada, H. 2012b. Contributions to the lichen flora of Gifu Prefecture, central Japan. (5). *Gyalecta kibiensis* (Gyalectaceae) as new for Gifu. *Lichenology* 11: 31–34.
- Knoph, J. -G. 2004. *Cornutispora herteliana*, a new lichenicolous coelomycete from Japan. *Bibliotheca Lichenologica* 88: 345–348.
- Knudsen, K. & Ryan, B. D. 2007. *Strangospora*. In: Nash, T. H. III, Gries, C. & Bungartz, F.: Lichen Flora of the Greater Sonoran Desert Region. Volume 3. Lichens Unlimited, Arizona State University, Tempe, pp. 299–301.
- Kondratyuk, S. Y. & Kärnefelt, I. 2003. Revision of three natural groups of xanthorioid lichens (Teloschistaceae, Ascomycota). *Ukrayins'kyi Botanichnyi Zhurnal* 60: 427–437.
- Kondratyuk, S. Y., Lőkös, L., Kim, J., Jeong, M. H., Kondratiuk, A., Oh, S. O. & Hur, J. S. 2014. *Kashiwadia* gen. nov. (Physciaceae, lichen-forming Ascomycota), proved by phylogenetic analysis of the Eastern Asian Physciaceae. *Acta Botanica Hungarica* 56: 369–378.
- Kondratyuk, S. Y., Lőkös, L., Upreti, D. K., Nayaka, S., Mishra, G. K., Ravera, S., Jeong, M. H., Jang, S. H., Park, J. S. & Hur, J. S. 2017a. New monophyletic branches of the Teloschistaceae (lichen-forming ascomycota) proved by three gene phylogeny. *Acta Botanica Hungarica* 59: 71–136
- Kondratyuk, S. Y., Lőkös, L., Halda, J. P., Roux, C., Upreti, D. K., Schumm, F., Mishra, G. K., Nayaka, S., Farkas, E., Park, J. S., Lee, B. G., Liu, D., Woo, J. J. & Hur, J. S. 2017b. New and noteworthy lichen-forming and lichenicolous fungi 6. *Acta Botanica Hungarica* 59: 137–260.
- Kraichak, E., Parnmen, S., Lücking, R., Rivas Plata, E., Aptroot, A., Cáceres, M. E. S., Ertz, D., Mangold, A., Mercado-Díaz, J. A., Papong, K., Van den Broeck, D., Weerakoon, G. & Lumbsch, H. T. 2014. Revisiting the phylogeny of Ocellulariaceae, the second largest tribe within Graphidaceae (lichenized Ascomycota: Ostropales). *Phytotaxa* 189: 52–81.
- Knudsen, K. & Kocourková, J. 2008. A study of lichenicolous species of *Polysporina* (Acarosporaceae). *Mycotaxon* 105: 149–164.
- Kurokawa, S. 1962. A monograph of the genus *Anaptychia*. Beihefte zur Nova Hedwigia No. 6. J. Cramer. 115 pp.
- Kurokawa, S. (ed.) 2003. Checklist of Japanese lichens. 128 pp. National Science Museum, Tokyo.
- Kurokawa, S. 2006. Phytogeographical elements of the lichen flora of Japan. *Journal of the Hattori Botanical Laboratory* 100: 721–738.
- Kurokawa, S. & Kashiwadani, H. 1988. Indices to taxa distributed under lichenes rariores et critici exsiccati. National Science Museum, Tokyo. 114 pp.
- Kurokawa, S. & Kashiwadani, H. 2006. Checklist of Japanese lichens and allied fungi. National Science Museum Monographs No. 33, National Science Museum, Tokyo. 157 pp.
- Kurokawa, S. & Nakanishi, S. 1971. Lichens of the Hidaka mountains, Hokkaido. *Memoir of the National Science Museum, Tokyo* 4: 59–70.

- Lai, M. J., Qian, Z. G. & Xu, L. 2007. Synopsis of the cetrarioid lichen genera and species (Parmeliaceae, lichenized Ascomycotina) in China. *Journal of the National Taiwan Museum* 60: 45–61.
- Lai, M. J., Chen, X. L., Qian, Z. G., Xu, L. & Ahti, T. 2009. Cetrarioid lichen genera and species in NE China. *Annales Botanici Fennici* 46: 365–380.
- Lamb, I. M. 1977. A conspectus of the lichen genus *Stereocaulon* (Schreb.) Hoffm. *Journal of the Hattori Botanical Laboratory* 43: 191–355.
- Laundon, J. R. 1992. *Lepraria* in the British Isles. *Lichenologist* 24: 315–350.
- Laundon, J. R. 2008. Some synonyms in *Chrysothrix* and *Lepraria*. *Lichenologist* 40: 411–414.
- Lendemer, J. C. 2013. A monograph of the crustose members of the genus *Lepraria* Ach. s. str. (Stereocaulaceae, Lichenized Ascomycetes) in North America north of Mexico. *Opuscula Philolichenum* 12: 27–141.
- Lendemer, J. C. & Harris, R. C. 2016. Studies in lichens and lichenicolous fungi – No. 20: Further notes on species from the eastern North America. *Opuscula Philolichenum* 15: 105–131.
- Lendemer, J. C. & Harris, R. C. 2017. Nomenclatural changes for North American members of the *Variolaria*-group necessitated by the recognition of *Lepra* (Pertusariales). *Bryologist* 120: 183–190.
- Lendemer, J. C. & Hodkinson, B. P. 2013. A radical shift in the taxonomy of *Lepraria* s.l.: molecular and morphological studies shed new light on the evolution of asexuality and lichen growth form diversification. *Mycologia* 105: 994–1018.
- Lendemer, J. C. & Knudsen, K. 2007. Changes and additions to the North American lichen Mycota—VI. *Proceedings of the Academy of Natural Sciences of Philadelphia* 156: 55–57.
- Lendemer, J. C., Sheard, J. W., Thor, G. & Tønsberg, T. 2012. *Rinodina chrysidata*, a new species from far eastern Asia and the Appalachian Mountains of North America. *Lichenologist* 44: 179–187.
- Louwhoff, S. H. J. J. 2009. Peltigeraceae. *Flora of Australia* 57: 428–446.
- Lücking, R. 2008. Follicolous lichenized fungi. *Flora Neotropica* 103. 866 pp.
- Lücking, R., Archer, A. W. & Aptroot, A. 2009. A world-wide key to the genus *Graphis* (Ostropales: Graphidaceae). *Lichenologist* 41: 363–452.
- Lücking, R., Mangold, A. & Lumbsch, H. T. 2016. A worldwide key to species of the genera *Myriotrema* and *Glaucotrema* (lichenized Ascomycota: Graphidaceae), with a nomenclatural checklist of species published in *Myriotrema*. *Herzogia* 29: 493–513.
- Lücking, R., Sérusiaux, E. & Vězda, A. 2005. Phylogeny and systematics of the lichen family Gomphillaceae (Ostropales) inferred from cladistic analysis of phenotype data. *Lichenologist* 37: 123–170.
- Lücking, R., Chaves, J. L., Sipman, H. J. M., Umaña, L. & Aptroot, A. 2008. A first assessment of the Ticolichen biodiversity inventory in Costa Rica: The genus *Graphis*, with notes on the genus *Hemithecium* (Ascomycota: Ostropales: Graphidaceae). *Fieldiana, Botany, new series* 46: 1–126.
- Lücking, R., Dal-Forno, M., Lawrey, J. D., Bungartz, F., Holgado Rojas, M. E., Hernandez, M. J. E., Marcelli, M. P., Moncada, B., Morales, E. A., Nelsen, M. P., Paz, E., Salcedo, L., Spielmann, A. A., Wilk, K., Will-Wolf, S. & Yanez, A. 2013. Ten new species of lichenized Basidiomycota in the genera *Dictyonema* and *Cora* (Agaricales: Hygrophoraceae), with a key to all accepted genera and species in the *Dictyonema* clade. *Phytotaxa* 139: 1–38.
- Lumbsch, H. T. 1989. Die holarktischen Vertreter der Flechtengattung *Diploschistes* (Thelotre mataceae). *Journal of the Hattori Botanical Laboratory* 66: 133–196.

- Magain, N., Miadlikowska, J., Mueller, O., Gajdeczka, M., Truong, C., Salamov, A. A., Dubchak, I., Grigoriev, I. V., Goffinet, B., Sérusiaux, E. & Lutzoni, F. 2017. Conserved genomic collinearity as a source of broadly applicable, fast evolving, markers to resolve species complexes: A case study using the lichen-forming genus *Peltigera* section *Polydactylon*. *Molecular Phylogenetics and Evolution* 117: 10–29.
- Magnusson, A. H. 1937. Additional notes on Acarosporaceae. *Meddelelser från Göteborgs Botaniska Trädgård* 12: 87–103.
- Mangold, A., Elix, J. A. & Lumbsch, H. T. 2009. Thelotremales. *Flora of Australia* 57: 195–420.
- Marbach, B. 2000. Corticole und lignicole Arten der Flechtengattung *Buellia* sensu lato in den Subtropen und Tropen. *Bibliotheca Lichenologica*, 74, J. Cramer, Berlin, Stuttgart. 384 pp.
- May, T. W. 2017. Report of the nomenclature committee for fungi-20. *IMA Fungus* 8: 189–203.
- Mayrhofer, H. 1984. Die saxicolen Arten der Flechtengattung *Rinodina* und *Rinodinella* in der alten Welt. *Journal of the Hattori Botanical Laboratory* 55: 327–493.
- Mayrhofer, H. 1987. Monographie der Flechtengattung *Thelenella*. *Bibliotheca Lichenologica* 26: 1–106.
- McCarthy, P. M. 1995. A reappraisal of *Clathroporina* Müll. Arg. (Trichotheliaceae). *Lichenologist* 27: 321–350.
- McCune, B. 2009. *Hypogymnia* (Parmeliaceae) species new to Japan and Taiwan. *Bryologist* 112: 823–826.
- McCune, B. 2011. *Hypogymnia irregularis* (Ascomycota: Parmeliaceae) —a new species from Asia. *Mycotaxon* 115: 485–494.
- McCune, B. & Wang, L. S. 2014. The lichen genus *Hypogymnia* in southwest China, *Mycosphere* 5: 27–76.
- Miyoshi, M. 1901. Ueber die Sporocarpenevacuation und darauf erfolgendes Sporenausstreuen bei einer Flechte. *Journal of the College of Science, Imperial University, Tokyo* 15: 367–370.
- Miyoshi, M. 1932. *Saishin Jikken Shokubutugaku*, 3rd ed. Fuzanbo, Tokyo. pp. 668+index 28 pp. (In Japanese).
- Moncada, B., Lücking, R. & Betancourt-Macuase, L. 2013. Phylogeny of the Lobariaceae (lichenized Ascomycota: Peltigerales), with a reappraisal of the genus *Lobariella*. *Lichenologist* 45: 203–263.
- Mongkolsuk, P., Meesim, S., Poengsungnoen, V., Buaruang, K., Schumm, F. & Kalb, K. 2015. The lichen family Physciaceae in Thailand—II. Contributions to the genus *Heterodermia* sensu lato. *Phytotaxa* 235: 1–66.
- Moon, K. H. & Kashiwadani, H. 2009. A new species of *Karoowia* (Parmeliaceae, Ascomycota) from Kyushu, Japan. *Journal of Japanese Botany* 84: 279–282.
- Moon, K. H., Ahn, C. & Kashiwadani, H. 2015. Revision of the lichen genus *Myelochroa* (Ascomycotina: Parmeliaceae) in Korea. *Journal of Species Research* 4: 23–32.
- Moon, K. H., Kashiwadani, H. & Kobayashi, K. 2013a. A new species of *Myelochroa* (Parmeliaceae, Lecanorales) from Shiga Prefecture, Japan. *Journal of Japanese Botany* 88: 140–143.
- Moon, K. H., Nakanishi, M. & Kashiwadani, H. 2008. Notes on species of Graphidaceae (Ascomycotina) from eastern Asia with three new species. *Memoirs of the National Science Museum, Tokyo* 45: 85–91.
- Moon, K. H., Ohmura, Y., Kashiwadani, H. & Yoshida, K. 2013b. Materials for the distribution of lichens in Japan (20) *Flavopunctelia flaventior* and *F. soledica*. *Journal of Japanese Botany* 88: 258–260.
- Moreno, P. P. & Egera, J. M. 1992. El genero *Lichinella*. *Cryptogamie, Bryologie, Lichénologie* 13: 237–259.
- Muggia, L., Nelson, P., Wheeler, T., Yakovchenko, L. S., Tønsberg, T. & Spribille, T. 2011. Convergent evolution of a symbiotic duet: the case of the lichen genus *Polychidium* (Peltigerales, Ascomycota). *American Journal of Botany* 98: 1647–1656.
- Müller, J. 1891. Lichenes Miyoshiani in Japonia a cl. Miyoshi lecti et a cl. professore Yatabe communicati.

- Nuovo Giornale Botanico Italiano 23: 120–202.
- Müller, J. 1892. Lichenes Yatabeani, in Japonia lecti et a cl. prof. Yatabe missi. Nuovo Giornale Botanico Italiano 24: 189–202.
- Nakanishi, M., Kashiwadani, H. & Moon, K. H. 2003. Taxonomical notes on Japanese Graphidaceae (Ascomycotina), including some new combinations. Bulletin of the National Science Museum, Tokyo, Series B (Botany) 29: 83–90.
- Navarro-Rosinés, P., Roux, C. & Gueidan, C. 2007. La genroj *Verrucula* kaj *Verruculopsis* (Verrucariaceae, Verrucariales). Bulletin de la Societe Linneenne de Provence 58: 133–180.
- Nordin, A., Savić, S. & Tibell, L. 2010. Phylogeny and taxonomy of *Aspicilia* and Megasperaceae. Mycologia 102: 1339–1349.
- Nordin, A. & Tibell, L. 2005. Additional species in *Tetramelas*. Lichenologist 37: 491–498.
- Nylander, W. 1890. Lichenes Japoniae. 122 pp. Paul Schmidt, Paris.
- Obermayer, E. 1997. Lichenotheca Graecensis. Fasc. 4 (Nos 61–80). Fritschiana 8: 1–6.
- Ohmura, Y., 2001. Taxonomic study of the genus *Usnea* (lichenized Ascomycetes) in Japan and Taiwan. Journal of the Hattori Botanical Laboratory 90:1–96.
- Ohmura, Y. 2012. A synopsis of the lichen genus *Usnea* (Parmeliaceae, Ascomycota) in Taiwan. Memoir of the National Museum of Nature and Science 48: 91–137.
- Ohmura, Y. & Frisch, A. 2016. *Sporodophoron primorskiense* (Arthoniaceae, lichenized Ascomycota) new to Japan, as the second locality in the world. Journal of Japanese Botany 91: 69–73.
- Ohmura, Y. & Mayrhofer, H. 2016. *Protothelenella sphinctrinoides* (Protothelenellaceae) new to Japan and new chemical features for several species in the genus. Herzogia 29: 137–142.
- Ohmura, Y., Moon, K. H. & Kashiwadani, H. 2008. Morphology and molecular phylogeny of *Ramalina pollinaria*, *R. sekika* and *R. yasudae* (Ramalinaceae, lichenized Ascomycotina). Journal of Japanese Botany 83: 156–164.
- Ohmura, Y., Yakovchenko, L. & Zhurbenko, M.P. 2014b. *Carbonea vitellinaria* new to Japan, with a key to lichenicolous fungi growing on species of *Candelariella*. Mycosphere 5: 607–611.
- Ohmura, Y., Yoshida, K. & Kashiwadani, H. 2013. Lectotypification of *Cetraria kurokawae* Shibuichi & K. Yoshida. Bulletin of the National Museum of Nature and Science, Series B (Botany) 39: 147–149.
- Ohmura, Y., Frisch, A., Lendemer, J. C. & Nakanishi, M. 2016b. Lichens of Mikura-jima Island, Izu Islands, Central Japan. Bulletin of the National Museum of Nature and Science, Series B (Botany) 42: 1–18.
- Ohmura, Y., Mizobuchi, A., Handa, S. & Lücking, R. 2016a. *Coenogonium moniliforme* (Coenogoniaceae, lichenized Ascomycota) new to Japan, with taxonomic notes of the photobiont in culture. Journal of Japanese Botany 91: 74–78.
- Ohmura, Y., Thor, G., Frisch, A., Kashiwadani, H. & Moon, K. H. 2014a. Increase of lichen diversity in the Imperial Palace Grounds, Tokyo, Japan. Memoirs of the National Science Museum, Tokyo 49: 193–217.
- Orange, A. 2013. Four new species of *Verrucaria* (Verrucariaceae, lichenized Ascomycota) from freshwater habitats in Europe. Lichenologist 45: 305–322.
- Orange, A., Wolseley, P., Karunaratne, V. & Bombuwala, K. 2001. Two leprarioid lichens new to Sri Lanka. Bibliotheca Lichenologica 78: 327–333.
- Otálora, M. A. & Wedin, M. 2013. *Collema fasciculare* belongs in Arctomiaceae. Lichenologist 45: 295–304.

- Otálora, M. A., Jørgensen, P. M. & Wedin, M. 2014. A revised generic classification of the jelly lichens, Collemataceae. *Fungal diversity* 64: 275–293.
- Parnmen, S., Lücking, R. & Lumbsch, H. T. 2012. Phylogenetic classification at generic level in the absence of distinct phylogenetic patterns of phenotypical variation: A case study in Graphidaceae (Ascomycota). *PLOS ONE* 7(12): e51392.
- Parnmen, S., Cáceres, M. E. S., Lücking, R. & Lumbsch, H. T. 2013. *Myriochapsa* and *Nitidochapsa*, two new genera in Graphidaceae (Ascomycota: Ostropales) for chroodiscoid species in the *Ocellularia* clade. *Bryologist* 116: 127–133.
- Papong, K., Lücking, R., Thammathaworn, A. & Boonpragob, K. 2009. Four new taxa of *Chroodiscus* (thelotremoid Graphidaceae) from Southeast Asia. *Bryologist* 112: 152–163.
- Paukov, A. G., Nordin, A., Roux, C., Moon, K. H. & Davydov, E. 2017. Lectotypification and synonymization of some *Aspicilia* species (Megasporaceae, Ascomycota) described by A. Hue from Korea and Japan. *Phytotaxa* 291: 94–98.
- Pišút, I., Lackovičová, A. & Lisická, E. 1993. Súpis Lišajníkov Slovenska. *Biologia, Bratislava* 48 (Supplement 1): 53–98.
- Prieto, M. & Wedin, M. 2017. Phylogeny, taxonomy and diversification events in the Caliciaceae. *Fungal Diversity* 82: 221–238.
- Printzen, C. 1995. Die Flechtengattung *Biatora* in Europa. *Bibliotheca Lichenologica* 60: 1–275.
- Printzen, C. 2004. *Biatora*. In: Nash, T. H., III, Ryan, B. D., Diederich, P., Gries, C. & Bungartz, F. (eds.): *Lichen flora of the Greater Sonoran Desert Region*, vol. 2. *Lichens Unlimited*, Arizona State University, Tempe, Arizona, pp. 37–39.
- Printzen, C. 2014. A molecular phylogeny of the lichen genus *Biatora* including some morphologically similar species. *Lichenologist* 46: 441–453.
- Räsänen, V. 1940. Lichenes ab A. Yasuda et aliis in Japonia collecti (II). *Journal of Japanese Botany* 16: 139–153.
- Resl, P., Schneider, K., Westberg, M., Printzen, C., Palice, Z., Thor, G., Fryday, A., Mayrhofer, H. & Spribille, T. 2015. Diagnostics for a troubled backbone: testing topological hypotheses of trapelioid lichenized fungi in a large-scale phylogeny of Ostropomycetidae (Lecanoromycetes). *Fungal diversity* 73: 239–258.
- Rivas Plata, E., Lücking, R., Sipman, H. J., Mangold, A., Klaus, K. A. L. B. & Lumbsch, H. T. 2010. A world-wide key to the thelotremoid Graphidaceae, excluding the *Ocellularia*-*Myriotrema*-*Stegobolus* clade. *Lichenologist* 42: 139–185.
- Rivas Plata, E., Lücking, R., Aptroot, A., Sipman, H. J. M., Chaves, J. L., Umaña, L. & Lizano, D. 2006. A first assessment of the Ticolichen biodiversity inventory in Costa Rica: the genus *Coenogonium* (Ostropales: Coenogoniaceae), with a world-wide key and checklist and a phenotype-based cladistic analysis. *Fungal Diversity* 23: 255–321.
- Roux, C., Bertrand, M. & Nordin, A. 2016. *Aspicilia serenensis* Cl. Roux et M. Bertrand sp. nov., espèce nouvelle de lichen (groupe d’*A. calcarea*, Megasporaceae). *Bulletin de la Societe Linneenne de Provence* 67: 165–182.
- Sakata, A., Harada, H. & Sato, H. 2009. *Cresponea* (lichenized Ascomycota, Roccellaceae) of Japan. *Lichenology* 8: 91–115.

- Sakata, A., Harada, H. & Yoshikawa, H. 2016. Taxonomic study on the lichen family Roccellaceae (Ascomycota, Arthoniales) of Japan (4). *Sclerophyton elegans*, new to Japan. *Lichenology* 14: 119–127.
- Sakata, A., Harada, H. & Yoshikawa, H. 2017. Taxonomic study on the lichen family Roccellaceae (Arthoniales) of Japan (5). Two new corticolous species of *Mazosia*. *Lichenology* 16: 31–47.
- Sakata, A., Harada, H., Yamamoto, Y. & Hara, K. 2014. The taxonomic study on the lichen family Roccellaceae (Ascomycota, Arthoniales) of Japan (2). *Graphidastra japonica* sp. nov. *Lichenology* 12: 51–60.
- Sato, M. 1943. Index Plantarum Nipponicarum IV. Lichenes. 128 pp. National Science Museum, Tokyo.
- Sato, M. 1959–65. Catalogus Lichenum Japonicorum (ed. 2). 56 pp. Mito.
- Sato, M. 1972. Lichen flora of Amami and Ryukyu Islands, Japan (I). *Bulletin of the College of General Education, Ibaraki University* 3: 25–40.
- Schmitt, I., Otte, J., Parnmen, S., Sadowska-Deś, A., Lücking, R. & Lumbsch, T. 2012. A new circumscription of the genus *Varicellaria* (Pertusariales, Ascomycota). *MycKeys* 4: 23–36.
- Schmull, M., Miadlikowska, J., Pelzer, M., Stocker-Wörgötter, E., Hofstetter, V., Fraker, E., Hodgkinson, B. P., Reeb, V., Kukwa, M., Lumbsch, H. T., Kauff, F. & Lutzoni, F. 2011. Phylogenetic affiliations of members of the heterogeneous lichen-forming fungi of the genus *Lecidea* sensu Zahlbruckner (Lecanoromycetes, Ascomycota). *Mycologia* 103: 983–1003.
- Sérusiaux, E., Lücking, R. & Sparrius, L. B. 2008. *Opegrapha viridistellata* (Roccellaceae), a new foliicolous lichen species from the Paleotropics. *Mycotaxon* 104: 223–227.
- Sheard, J. W., Ezhkin, A. K., Galanina, I. A., Himelbrant, D., Kuznetsova, E., Shimizu, A., Stepanchikova, I., Thor, G., Tønsberg, T., Yakovchenko, L. S. & Spribille, T. 2017. The lichen genus *Rinodina* (Physciaceae, Caliciales) in north-eastern Asia. *Lichenologist* 49: 617–672.
- Shiba, T., Moon, K. H. & Kashiwadani, H. 2008. Lichens of Akkeshi and its adjacent areas, northeastern Hokkaido, Japan. *Bulletin of the National Science Museum, Series B (Botany)* 34: 1–16.
- Sparrius, L. B. 2004. A monograph of *Enterographa* and *Sclerophyton*. *Bibliotheca Lichenologica* 89: 1–141.
- Spribille, T., Klug, B. & Mayrhofer, H. 2011. A phylogenetic analysis of the boreal lichen *Mycoblastus sanguinarius* (Mycoblastaceae, lichenized Ascomycota) reveals cryptic clades correlated with fatty acid profiles. *Molecular Phylogenetics and Evolution* 59: 603–614.
- Spribille, T., Tønsberg, T., Stabentheiner, E. & Muggia, L. 2014. Reassessing evolutionary relationships in the filamentous cyanolichen genus *Spilonema* (Peltigerales, Lecanoromycetes). *Lichenologist* 46: 373–388.
- Spribille, T., Resl, P., Ahti, T., Pérez-Ortega, S., Tønsberg, T., Mayrhofer, H. & Lumbsch, H. T. 2014. Molecular systematics of the wood-inhabiting, lichen-forming genus *Xylographa* (Baeomycetales, Ostropomycetidae) with eight new species. *Symbolae Botanicae Upsalienses* 37: 1–87.
- Staiger, B. 2002. Die Flechtenfamilie Graphidaceae. *Bibliotheca Lichenologica* 85: 1–526.
- Staiger, B. & Kalb, K. 1995. *Haematomma*. *Bibliotheca Lichenologica* 59: 1–198.
- Stevens, G. N. 1987. The lichen genus *Ramalina* in Australia. *Bulletin of the British Museum (Natural History), Botany Series* 16: 107–223.
- Takahashi, K., Anzai, T., Kinoshita, Y. & Harada, H. 2012. *Sticta praetextata* and *S. sublimbata* (lichenized Ascomycota; Lobariaceae) new to Japan. *Lichenology* 11: 1–7.
- Tehler, A. 1993. The genus *Schismatomma* (Arthoniales, Euascomycetidae). *Nordic Journal of Botany* 13: 462–462.

- Thell, A., Randle, T., Saag, A. & Kärnefelt, I. 2005. A new circumscription of the lichen genus *Nephromopsis* (Parmeliaceae, lichenized Ascomycetes). *Mycological Progress* 4: 303–316.
- Thell, A., Högnabba, F., Elix, J. A., Feuerer, T., Kärnefelt, I., Myllys, L., Randle, T., Saag, A., Stenroos, S., Ahti, T. and Seaward, M. R. D. 2009. Phylogeny of the cetrarioid core (Parmeliaceae) based on five genetic markers. *Lichenologist* 41: 489–511.
- Tibell, L. & Thor, G. 2003. Calicioid lichens and fungi of Japan. *Journal of the Hattori Botanical Laboratory* 94: 205–259.
- Tibell, L., Frisch, A. & Thor, G. 2014. Additions to the calicioid flora of Japan and Korea, with the descriptions of two new species. *Annales Botanici Fennici* 51: 189–194.
- Timdal, E. 2008. Studies on *Phyllopsora* (Ramalinaceae) in Peru. *Lichenologist* 40: 337–362.
- Timdal, E. & Tønsberg, T. 2012. *Cladonia straminea*, the correct name for *C. metacorallifera*. *Graphis Scripta* 24: 33–35.
- Tojo, B., Kawakami, S., Harada, H. & Okamoto, T. 2011. Contributions to the lichen flora of Gifu Prefecture, Central Japan. (2). *Caloplaca cinnabarina* (Teloschistaceae) as new for Japan. *Lichenology* 10: 47–51.
- Tokizawa, M., Ohmura, Y., Moon, K. H. & Takeshita, S. 2015. *Sarcogyne endopetrophila* (Acarosporaceae, lichenized Ascomycota), a new species from Japan. *Journal of Japanese Botany* 90: 46–51.
- Triebel, D. 1989. Lecideicole Ascomyceten. Eine Revision der obligat lichenicolen Ascomyceten auf lecideoiden Flechten. *Bibliotheca Lichenologica* 35: 1–278.
- Vainio [“Wainio”], E. A. 1918. Lichenes ab A. Yasuda in Japonia collecti. *Botanical Magazine, Tokyo* 32: 154–163.
- Vainio [“Wainio”], E. A. 1921. Lichenes ab A. Yasuda in Japonia collecti. *Continuatio I*. *Botanical Magazine, Tokyo* 35: 45–79.
- Vitikainen, O., Ahti, T., Kuusinen, M., Lommi, S. & Ulvinen, T. 1997. Checklist of lichens and allied fungi of Finland. *Norrinia* 6: 1–123.
- Watanuki, O., Hara, K., Harada, H., Komine, M. & Fuji, S. 2017a. *Buellia numerosa* and *B. subnumerosa*, two new species of the lichen genus *Buellia* (Caliciaceae) from Japan. *Bryologist* 120: 25–36.
- Watanuki, O., Harada, H., Hara, K., Komine, M., Yamamoto, Y. & Fuji, S. 2017b. Taxonomic study of *Buellia* s. lat. (Lichenized Ascomycota, Caliciales, Caliciaceae) in Japan (2). *Buellia erubescens* newly found in Japan. *Lichenology* 16: 49–57.
- Wei, X., Schmitt, I., Hodgkinson, B. P., Flakus, A., Kukwa, M., Divakar, P. K., Kirika, P. M., Otte, J., Meiser, A. & Lumbsch, H. T. 2017. Circumscription of the genus *Lepra*, a recently resurrected genus to accommodate the “*Variolaria*”-group of *Pertusaria* sensu lato (Pertusariales, Ascomycota). *PLOS ONE* 12: e0180284.
- Wirth, V. 1987. *Die Flechten Baden-Württembergs*. Eugen Ulmer, GmbH & Co., Stuttgart. 528 pp.
- Yamamoto, Y., Takahagi, T., Bando, M. & Kawakami, H. 2017. Lichen data of Osaka Prefecture I. Lichens of the Nagai Park (Osaka City) and four noteworthy species including two as new for Japan. *Bulletin of the Osaka Museum of Natural History* 71: 11–16.
- Yamamoto, Y., Kurogi, S., Hara, K., Komine, M., Kawakami, H. & Kalb, K. 2013. *Ramboldia haematites* (Fée) Kalb *et al.* new to Asia. *Lichenology* 11: 67–71.
- Yanaga, K., Sotome, K., Suhara, H. & Maekawa, N. 2015. A new species of *Lepidostroma* (Agaricomycetes, Lepidostromataceae) from Japan. *Mycoscience* 56: 1–9.

- Yao, Y. -J. & Spooner, B. M. 1999. Roesleriaceae, a new family of Ascomycota, and a new species of *Roeslerina*. Kew Bulletin 54: 683–693.
- Yasuda, A. 1911. Shokubutsugaku Kakuron. Inka-bu. Hakubunkan, Tokyo. 1222 pp. (In Japanese)
- Yoshida, K. & Kashiwadani, H. 2001. Typification of taxa of *Hypogymnia* (Parmeliaceae) described from Japan and Sakhalin. Bulletin of National Science Museum, Tokyo, Series B (Botany) 27: 35–46.
- Yoshikawa, H. & Harada, H. 2012. *Pilophorus cereolus*, newly found in Japan. Lichenology 11: 9–11.
- Yoshimura, I. 1962. Enumeration of Stictaceae in Shikoku. Acta Phytotaxonomica et Geobotanica 19: 92–96.
- Yoshimura, I. 1974. Lichen flora of Japan in colour. 349 pp. Hoikusha, Osaka.
- Yoshimura, I. & Harada, H. 2004. Taxonomic notes on *Psilolechia lucida* as new for Japan. Lichenology 3: 41–46.
- Zhao, X., Leavitt, S. D., Zhao, Z. T., Zhang, L. L., Arup, U., Grube, M., Pérez-Ortega, S., Printzen, C., Śliwa, L., Kraichak, E., Divakar, P. K., Crespo, A. & Lumbsch, H. T. 2016 [2015]. Towards a revised generic classification of lecanoroid lichens (Lecanoraceae, Ascomycota) based on molecular, morphological and chemical evidence. Fungal Diversity 78: 293–304.
- Zhurbenko, M. P., Braun, U., Heuchert, B. & Kobzeva, A. A. 2015a. New lichenicolous hyphomycetes from Eurasia. Herzogia 28: 584–598.
- Zhurbenko, M. P., Ezhkin, A. K., Skirina, I. F. & Ohmura, Y. 2017. *Dactylospora anziaae*, a new lichenicolous ascomycete on *Anzia* from East Asia. Folia Cryptogamica Estonica 54: 13–16.
- Zhurbenko, M. P., Frisch, A., Ohmura, Y. & Thor, G. 2015b. Lichenicolous fungi from Japan and Korea: new species, new records and a first synopsis for Japan. Herzogia 28: 762–789.

**National Museum of Nature and Science Monographs**  
(Nos. 1-35 were published as “National Science Museum Monographs”)

- No. 1\*. Early Cretaceous marine and brackish-water Gastropoda from Japan. By Tomoki Kase, 199 pp., 31 pls., 1984.
- No. 2\*. A taxonomic study on the subfamily Herminiinae of Japan (Lepidoptera, Noctuidae). By Mamoru Owada, 208 pp., 1987.
- No. 3\*. Small mammal fossils and correlation of continental deposits, Safford and Duncan Basins, Arizona, USA. By Yukimitsu Tomida, 141 pp., 1987.
- No. 4. Late Miocene floras in northeast Honshu, Japan. By Kazuhiko Uemura, 174 pp., 11 pls., 1988.
- No. 5. A revisional study of the spider family Thomisidae (Arachnida, Araneae) of Japan. By Hirotugu Ono, 252 pp., 1988.
- No. 6. The taxonomic study of Japanese dictyostelid cellular slime molds. By Hiromitsu Hagiwara, 131 pp., 1989.
- No. 7. A systematic study of the Japanese Chiroptera. By Mizuko Yoshiyuki, 242 pp., 1989.
- No. 8. Rodent and lagomorph families of Asian origins and diversification: Proceedings of Workshop WC-2 29th International Geological Congress, Kyoto, Japan. Edited by Yukimitsu Tomida, Chuankuei Li, and Takeshi Setoguchi, 195 pp., 1994.
- No. 9. A microevolutional history of the Japanese people as viewed from dental morphology. By Hirofumi Matsumura, 130 pp., 1995.
- No. 10. Studies on the human skeletal remains from Jiangnan, China. Edited by Bin Yamaguchi and Huan Xianghong, 108 pp., 3 pls., 1995.
- No. 11. Annotated checklist of the inshore fishes of the Ogasawara Islands. By John E. Randall, Hitoshi Ida, Kenji Kato, Richard L. Pyle, and John L. Earle, 74 pp., 19 pls., 1997.
- No. 12. Deep-sea fauna and pollutants in Suruga Bay. By Tsunemi Kubodera and Masaaki Machida, *et al.*, 336 pp., 12 pls., 1997.
- No. 13. Polychaetous annelids from Sagami Bay and Sagami Sea collected by the Emperor Showa of Japan and deposited at the Showa Memorial Institute, National Science Museum, Tokyo. Families Polynoidae and Acoetidae. By Minoru Imajima, 131 pp., 1997.
- No. 14. Advance in vertebrate paleontology and geochronology. Edited by Yukimitsu Tomida, Lawrence J. Flynn, and Louis L. Jacobs, 292 pp., 1998.
- No. 15. Proceedings of the Second Gondwanan Dinosaur Symposium. Edited by Yukimitsu Tomida, Thomas H. Rich, and Patricia Vickers-Rich, 296 pp., 1999.
- No. 16. Onuphidae (Annelida, Polychaeta) from Japan, excluding the genus *Onuphis*. By Minoru Imajima, 115 pp., 1999.
- No. 17. Description of a new species of Anhangueridae (Pterodactyloidea) with comments on the pterosaur fauna from the Santana Formation (Aptian-Albian), northeastern Brazil. By Alexander W. A. Kellner and Yukimitsu Tomida, 135 pp., 2000.
- No. 18. Proceedings of the First and Second Symposia on Collection Building and Natural History Studies in Asia. Edited by Keiichi Matsuura, 188 pp., 2000.

- No. 19. A taxonomic revision of the marine species of *Cladophora* (Chlorophyta) along the coasts of Japan and the Russian Far-east. By Christiaan van den Hoek and Mitsuo Chihara, 242 pp., 2000.
- No. 20. Deep-sea fauna and pollutants in Tosa Bay. Edited by Toshihiko Fujita, Hiroshi Saito and Masatsune Takeda, 380 pp., 2001.
- No. 21. Marine Fauna of the Shallow Waters around Hainan Island, South China Sea. Edited by Keiichi Matsuura, 126 pp., 2001.
- No. 22. Proceedings of the Third and Fourth Symposia on Collection Buildings and Natural History Studies in Asia and the Pacific Rim. Edited by Tsunemi Kubodera, Masanobu Higuchi, and Ritsuro Miyawaki, 193 pp., 2002.
- No. 23. Polychaetous Annelids from Sagami Bay and Sagami Sea Collected by the Emperor Showa of Japan and Deposited at the Showa Memorial Institute, National Science Museum, Tokyo (II). Orders included within the Phyllodocida, Amphinomida, Spintherida and Eunicida. By Minoru Imajima, 221 pp., 2003.
- No. 24. Proceedings of the Fifth and Sixth Symposia on Collection Building and Natural History Studies in Asia and the Pacific Rim. Edited by Shinobu Akiyama, Ritsuro Miyawaki, Tsunemi Kubodera, and Masanobu Higuchi 292 pp., 2004.
- No. 25. Revision of scydmaenid beetles of the genus *Syndicus* Motschulsky (Coleoptera, Scydmaenidae). By Pawel Jaloszynski, 108 pp., 2004.
- No. 26. A new specimen of *Apatosaurus ajax* (Sauropoda: Diplodocidae) from the Morrison Formation (Upper Jurassic) of Wyoming, USA. By Paul Upchurch, Yukimitsu Tomida, and Paul M. Barrett, 108 pp., 10 pls., 2004.
- No. 27. Leaf-rolling Sawflies of the *Pamphilius vafer* complex (Hymenoptera, Pamphiliidae). By Akihiko Shinohara, 116 pp., 2005.
- No. 28. Types of Japanese Bird. By Hiroyuki Morioka, Edward C. Dickinson, Takashi Hiraoka, Desmond Allen and Takeshi Yamasaki, 154 pp., 2005.
- No. 29. Deep-sea fauna and pollutants in Nansei Islands. Edited by Kazunori Hasegawa, Gento Shinohara and Masatsune Takeda, 476 pp., 2005.
- No. 30. Phenology and Growth Habits of Tropical Trees: Long-term Observations in the Bogor and Cibodas Botanic Garden, Indonesia. Edited by Hiroaki Hatta and Dedy Darnaedi, 436 pp., 2005.
- No. 31. The Cretaceous System in the Makarov Area, Southern Sakhalin, Russian Far East. By Yasunari Shigeta and Haruyoshi Maeda, 136 pp., 2005.
- No. 32. Revision of the Palearctic Species of the Myrmecophilous Genus *Pella* (Coleoptera, Staphylinidae, Aleocharinae). By Munetoshi Maruyama, 207 pp., 2006.
- No. 33. Checklist of Japanese lichens and allied fungi. By Syo Kurokawa and Hiroyuki Kashiwadani, 157 pp., 2006.
- No. 34. Proceedings of the Seventh and Eighth Symposia on Collection Building and Edited by Yukimitsu Tomida, Tsunemi Kubodera, Shinobu Akiyama and Taiju Kitayama, 294 pp., 2006.
- No. 35. Anatomy of a Japanese Tomistomine Crocodylian, *Toyotamaphimeia machikanensis* (Kamei et Matsumoto, 1965), from the Middle Pleistocene of Osaka Prefecture: The Reassessment of its Phylogenetic Status within Crocodylia. By Yoshitsugu Kobayashi, Yukimitsu Tomida, Tadao Kamei

and Taro Eguchi, 121 pp., 2006.

- No. 36. A systematic study of the genus *Siriella* (Crustacea: Mysida) from the Pacific and Indian oceans, with descriptions of fifteen new species. Edited by Masaaki Murano and Kouki Fukuoka, 173 pp., 2008.
- No. 37. Chromosome atlas of flowering plants in Japan. Edited by Tsunehiko Nishikawa, 706 pp., 2008.
- No. 38. The Lower Triassic System in the Abrek Bay area, South Primorye, Russia. Edited by Yasunari Shigeta, Yuri D. Zakharov, Haruyoshi Maeda and Alexander M. Popov, 218 pp., 2009.
- No. 39. Deep-sea Fauna and Pollutants off Pacific Coast of Northern Japan. Edited by Toshihiko Fujita, 755 pp., 2009.
- No. 40. Joint Haeckel and Ehrenberg Project: Reexamination of the Haeckel and Ehrenberg Microfossil Collections as a Historical and Scientific Legacy. Edited by Yoshihiro Tanimura and Yoshiaki Aita, 106 pp., 2009.
- No. 41. Middle and Late Miocene Marine Bivalves from the Northern Kanto Region, Central Japan. Edited by Yukito Kurihara, 87 pp., 2010.
- No. 42. Chemical Compositions of Electrum Grains in Ore and Placer Deposits in the Japanese Islands. By Kazumi Yokoyama, Shogo Takeuchi, Izumi Nakai, Yukiyasu Tsutsumi, Takashi Sano, Masako Shigeoka, Ritsuro Miyawaki and Satoshi Matsubara, 80 pp., 2011.
- No. 43. Digeneans (Trematoda) Parasitic in Freshwater Fishes (Osteichthyes) of the Lake Biwa Basin in Shiga Prefecture, Central Honshu, Japan. By Takeshi Shimazu, Misako Urabe and Mark J. Grygier, 105 pp., 2011.
- No. 44. Deep-sea Fauna of the Sea of Japan. Edited by Toshihiko Fujita, 291 pp., 2014.
- No. 45. Olenekian (Early Triassic) Stratigraphy and Fossil Assemblages in Northeastern Vietnam. Edited by Yasunari Shigeta, Toshifumi Komatsu, Takumi Maekawa and Huyen Dang Tran, 309 pp., 2014.
- No. 46. The Subfamily Steninae Macleay, 1825 (Coleoptera: Staphylinidae) of Japan Part 1. *Dianous* and *Stenus* (*S. Comma* Group to *S. Guttalis* Group). By Shun-Ichiro Naomi, Shûhei Nomura and Volker Puthz, 339 pp., 2017.
- No. 47. Bio-anthropological Studies of Early Holocene Hunter-Gatherer Sites at Huiyaotian and Liyupo in Guangxi, China. Edited by Hirofumi Matsumura, Hsiao-chun Hung, Li Zhen and Kenichi Shinoda, 228 pp., 2017.
- No. 48. Flavonoids of the Plants Composing Flora of Japan. By Tsukasa Iwashina, 647 pp., 2018.
- No. 49. Checklist of Lichens and Allied Fungi of Japan. Edited by Yoshihito Ohmura and Hiroyuki Kashiwadani, 143 pp., 2018.

(\* out of print)

All inquiries concerning the Monographs should be addressed to:

Library

National Museum of Nature and Science

Amakubo 4-1-1, Tsukuba-city

Ibaraki 305-0005

Japan

