

Cytotaxonomical Studies of Orchidaceae from Vanuatu and its Adjacent Regions: II

Goro KOKUBUGATA* and Tomohisa YUKAWA*

Comprehensive observations of chromosomal characters of orchid taxa in Melanesia are needed because there are few cytotaxonomical reports on Melanesian orchids such as Jones *et al.* (1982), Lim (1985a, b, c), Ishida *et al.* (1992), and Kokubugata and Yukawa (1998). In this study, we investigate the chromosomal characters of 7 taxa from Vanuatu and adjacent Pacific islands.

Materials and Methods

Table 1 shows the materials used in this study. Further data for each collection are provided in Hashimoto *et al.* (1998), Konishi *et al.* (1998), and Yukawa (1998). Voucher specimens and permanent slides are deposited in TNS.

Root tips were harvested from plants in cultivation and pretreated in 2mM 8-hydroxyquinoline at 20 °C for 2 hr. Subsequently, they were fixed in 45% acetic acid at 4°C for 10 sec and then macerated in a mixture of 1N hydrochloric acid and 45% acetic acid (2:1) at 60°C for 10 sec. Following these treatments, they were put on slides and stained in 1% aceto-orcein for 2 hr. The squash method was applied to investigate chromosomes at mitotic metaphase.

Results and Discussion

1. *Acanthephippium splendidum* J. J. Sm., $2n = 46$ (Figs. 1A and 2).

This is a new count for the genus *Acanthephippium*. The prevailing chromosome number of this genus is $2n = 48$ (e.g., Tanaka 1965, Hsu 1976), but $2n = 40$ (e.g., Vij and Shekhar 1983) and $2n = 42$ (e.g., Li and Chen 1989) were also observed. It is likely that aneuploid series play an important role on the evolution of this genus.

2. *Spathoglottis unguiculata* (Labill.) Rchb. f., $2n = 40$ (Figs. 1B and 3).

This is the first cytological record for the species. Previously, $2n = 40$ and 60 were counted for this genus.

3. *Spathoglottis pacifica* Rchb. f., $2n = 40$ (Figs. 1C and 4).

This species also has not been cytologically studied and the present count represents the first record.

4. *Spathoglottis petri* Rchb. f., $2n = 40$ (Figs. 1D and 5).

This is also the first cytological record for the species. Based on our results and previous studies, $2n = 40$ is a sole somatic chromosome number of *Spathoglottis* except for a single triploid record ($2n =$

*Tsukuba Botanical Garden, National Science Museum, Tsukuba, 305-0005 Japan.




Fig. 1. Orcein-stained chromosomes at mitotic metaphase of seven species of Orchidaceae. A. *Acanthephippium splendidum*. B. *Spathoglottis unguiculata*. C. *Spathoglottis pacifica*. D. *Spathoglottis petri*. E. *Calanthe triplicata*. F. *Calanthe ventrilabrum*. G. *Glossorhyncha macdonaldii*. Bar = 10 μ m.



Fig. 2. *Acanthephippium splendidum* in cultivation, × ca. 0.5. Sugimura 97-4307.



Fig. 3. *Spathoglottis unguiculata* in cultivation, × ca. 0.6. Hashimoto 9710720.




Fig. 4. *Spathoglottis pacifica* in cultivation, × ca. 0.4. Yukawa 97-2161.




Fig. 5. *Spathoglottis petri* in cultivation, × ca. 0.5. Yukawa 97-2232.



Fig. 6. *Calanthe triplicata* in cultivation, × ca. 0.7. Yukawa 97-2303.



Fig. 7. *Calanthe ventralbrum* in cultivation, × ca. 0.2. Yukawa 97-2095.



Fig. 8. *Glossorhyncha macdonaldii* in cultivation, × ca. 1.2. Yukawa 97-2067.

Table. 1. Chromosome numbers of seven species of Orchidaceae observed

Species	Locality	Voucher	Chrom. no. (2n)
<i>Acanthephippium splendidum</i>	Fiji: Viti Levu	Sugimura 97-4307	46
<i>Spathoglottis unguiculata</i>	Vanuatu: Espiritu Santo	Hashimoto 9710720	40
<i>Spathoglottis pacifica</i>	Vanuatu: Espiritu Santo	Yukawa 97-2161	40
<i>Spathoglottis petri</i>	Vanuatu: Efate	Yukawa 97-2232	40
<i>Calanthe triplicata</i>	New Caledonia: Grande Terre	Yukawa 97-2303	40
<i>Calanthe ventilabrum</i>	Vanuatu: Espiritu Santo	Yukawa 97-2095	40
<i>Glossorrhyncha macdonaldii</i>	Vanuatu: Espiritu Santo	Yukawa 97-2067	40

60) for *S. plicata* (Teoh 1980).

5. *Calanthe triplicata* (Willemet) Ames, 2n = 40 (Figs. 1E and 6).

Both vegetative and reproductive parts of Melanesian material are much larger than plants from other regions. Several authors recognized it as a distinct variety, *C. triplicata* var. *angraeciflora* (Schltr.) N. Hallé. We thus suspected poliploidy of these plants, but the number is identical to previous records of *C. triplicata* from various regions.

6. *Calanthe ventilabrum* Rchb. f., 2n = 40 (Figs. 1F and 7)

We confirmed the count of Ishida *et al.* (1992) where they treated this species as *C. langei* F. Muell., a later synonym. This species belongs to section *Styloglossum* and all the counts hitherto for this section show 2n = 40.

7. *Glossorrhyncha macdonaldii* Schltr., 2n = 40 (Figs. 1G and 8).

This is the second cytological record for the genus *Glossorrhyncha*. The chromosome number of this species was consistent with that of *G. chlorantha* van Royen from Papua New Guinea (Lim 1985c). Dressler (1993) disposed this genus into tribe Epidendreae subtribe Glomerinae in which 2n = 38 and 46 have been recorded for *Agrostophyllum*.

Summary

Mitotic chromosomes of seven taxa from Vanuatu and its adjacent regions were examined by the standard aceto-orcein staining method. *Acanthephippium splendidum* showed the chromosome number of 2n = 46; and three species of *Spathoglottis*, two species of *Calanthe*, and a single species of *Glossorrhyncha* showed that of 2n = 40.

Acknowledgements

We are indebted to Mr. Kazuhiro Suzuki for skillful cultivation of plant material for the present study. Financial support for this work was partially provided by the International Scientific Research Program, Nos. 08041165 and 12575013 (Representative: Tsukasa Iwashina), Japanese Ministry of Education, Science, Sports, Culture and Technology.

References

- Dressler, R. L., 1993. *Phylogeny and Classification of the Orchid Family*. Dioscorides Press, Portland.
- Hashimoto, T., S. Matsumoto, T. Yukawa, T. Konishi, K. Sugimura and T. Iwashina, 1998. A list of herbarium and live specimens of Grande Terre (New Caledonia) and Viti Levu (Fiji), collected in 1996 and 1997. *Ann. Tsukuba Bot. Gard.* 17: 105-131.
- Hsu, C., 1976. Cytological studies on the economically promising wild orchids found on Taiwan. *Proc. Natl. Sci. Council (Taiwan)* 9: 61-78.
- Ishida, G., T. Sera and K. Hashimoto, 1992. Chromosome numbers of some orchids from New Caledonia. *Bull. Hiroshima Bot. Gard.* 14: 47-50.
- Jones, K., K. Y. Lim and P. J. Cribb, 1982. The chromosomes of orchids VII *Dendrobium*. *Kew Bull.* 37: 221-227.
- Kokubugata, G. and T. Yukawa, 1998. Cytotaxonomical studies of the Orchidaceae from Vanuatu and its adjacent regions: I. *Ann. Tsukuba Bot. Gard.* 17: 69-74.
- Konishi, T., T. Hashimoto, T. Yukawa, S. Matsumoto, R. Hirayama, S. Chanel and T. Iwashina, 1998. A list of live specimens of Vanuatu, collected in 1996 and 1997. *Ann. Tsukuba Bot. Gard.* 17: 23-50.
- Li, X.-L. and R.-Y. Chen, 1989. Studies on chromosomes of Orchidaceae in China I. Chromosome numbers of some orchids in China. *In: D. Hong (ed.), Plant Chromosome Research 1987. Organizing Committee Sino-Japanese Symposium on Plant Chromosomes, Beijing.* pp. 301-307.
- Lim, K. -Y., 1985a. The chromosomes of orchids at Kew-1-*Bulbophyllum*. *Amer. Orchid Soc. Bull.* 54: 190-191.
- , 1985b. The chromosomes of orchids at Kew-2-*Dendrobium*. *Amer. Orchid Soc. Bull.* 54: 1122-1123.
- , 1985c. The chromosomes of orchids at Kew-3-Miscellaneous species. *Amer. Orchid Soc. Bull.* 54: 1234-1235.
- Tanaka, R., 1965. Chromosome numbers of some species of Orchidaceae from Japan and its neighbouring areas. *J. Jpn. Bot.* 40: 65-77.
- Teoh, S. B., 1980. Cytological studies in Malayan members of the *Phaius* tribe (Orchidaceae). II. Meiotic and B'-chromosomes. *Caryologia* 33: 483-493.
- Vij, S. P. and N. Shekhar, 1983. *In IOPB chromosome number reports LXXXI.* *Taxon* 32: 668.
- Yukawa, T., 1998. Enumeration of herbarium specimens of Orchidaceae collected in the 1997 Vanuatu Expedition. *Ann. Tsukuba Bot. Gard.* 17: 63-67.