

Comparison of Chromosome Number and Karyotype of Two *Lepidozamia* Species (Zamiaceae, Cycadales)

Goro KOKUBUGATA*, Katsuhiko KONDO** and Lou M. RANDALL***

國府方吾郎*・近藤勝彦**・Lou M. Randall***: *Lepidozamia* 属 (ザミア科, ソテツ目)
2 種における染色体数及び核型の比較研究

Lepidozamia (Zamiaceae, Cycadales; Stevenson 1992) is endemic to Australia, and consists of *L. hopei* Regel and *L. peroffskyana* Regel. Two species of *Lepidozamia* are enable to be recognized with the width of leaflet; *i. e.* the former has wider leaflet than the latter (Jones 1993).

The chromosome number and karyotype of *L. hopei* and *L. peroffskyana* was previously reported by Marchant (1968) and Moretti (1990). In certain cycad-taxa, however, there are some mistakes of their karyotype identification for the previous reports, and it is necessary to reinvestigate their of the chromosomal characters (Kokubugata *et al.* 1999). The aim of the present study is investigation of the chromosome number and karyotype of *L. hopei* and *L. peroffskyana* by the standard aceto-orcein staining method for the cytotaxonomical accumulations of *Lepidozamia*.

Materials and Methods

The taxonomical treatment follows Jones (1993). The plants investigated in the present study were collected in the each habit and exported under a CITES permit before cultivation in the greenhouse of the Tsukuba Botanical Garden (Table 1).

Root tips were harvested and pretreated in 4 mM 8-hydroxyquinoline at 4°C for 10 hour. They were fixed in acetic ethanol (3:1) at 4°C for 24 hour, and were transferred and stored in 70% ethanol at -20°C.

Fixed Root tips were macerated in a mixture of 1N HCl and 45% acetic acid (2:1) at 60°C for 10 sec., put on glass slides, and then stained in 2% aceto-orcein at room temperature for 4 hour. Orcein-stained chromosomes were classified according to centromeric position defined using arm ratios (Levan *et al.* 1964).

Table 1. *Lepidozamia* species used in this study

Species	Origin	Accession no.
<i>L. hopei</i>	Australia, State of Queensland: El Arish	TBG126733
<i>L. perffoskyana</i>	Australia, State of Queensland: Caloundra	TBG122893

* Tsukuba Botanical Garden, National Science Museum, Tsukuba, 305-0005. 国立科学博物館 筑波研究資料センター 筑波実験植物園.

** Laboratory Plant Chromosome and Gene Stock, Faculty of Science, Hiroshima University, Higashi-Hiroshima 739-8526. 広島大学 理学部附属植物遺伝子保管実験施設 〒739-8526 広島県東広島市鏡山1-4-3.

***Rockhampton Botanic Gardens, PO Box 243, Rockhampton 4700, State of Queensland, Australia.



Fig. 1. Chromosome complement of two *Lepidozamia* species at mitotic metaphase. A. *L. hopei*. B. *L. peroffskyana*. Bar indicates 10 μ m.



Fig. 2. Somatic chromosome of two *Lepidozamia* species at mitotic metaphase. A. *L. hopei*. B. *L. peroffskyana*. Bar indicates 10 μ m.

Table 2. Arm ratio and chromosome length in the complement of two *Lepidozamia* species at mitotic metaphase

Chrom. No.	Long arm (μm)		Long arm (μm)		Long arm (μm)		Arm ratio		Chromosomal classification	
	H	P	H	P	H	P	H	P	H	P
1	15.1	23.0	13.0	19.0	28.1	42.0	1.2	1.2	m	m
2	13.8	22.5	13.5	17.8	27.3	40.3	1.0	1.3	m	m
3	13.2	20.0	13.0	18.8	26.2	38.8	1.0	1.1	m	m
4	14.5	20.0	11.6	18.0	26.1	38.0	1.3	1.1	m	m
5	13.0	19.0	13.0	15.5	26.0	34.5	1.0	1.2	m	m
6	12.6	17.0	11.5	15.2	24.1	32.2	1.1	1.1	m	m
7	9.1	15.0	8.8	12.0	17.9	27.0	1.0	1.3	m	m
8	8.2	13.0	7.8	11.4	16.0	24.4	1.1	1.1	m	m
9	13.5	19.5	5.0	4.5	18.5	24.0	2.7	4.3	sm	st
10	12.8	18.5	5.1	5.0	17.9	23.5	2.5	3.7	sm	st
11	14.0	17.5	3.8	5.1	17.8	22.6	3.7	3.4	st	st
12	12.3	15.5	3.5	5.8	15.8	21.3	3.5	2.7	st	sm
13	12.2	16.9	3.4	3.8	15.6	20.7	3.6	4.4	st	st
14	12.1	17.0	3.0	3.5	15.1	20.5	4.0	4.9	st	st
15	14.9	20.5	0.6	0.5	15.5	21.0	24.8	41.0	t	t
16	14.5	20.0	0.7	0.5	15.2	20.5	20.7	40.0	t	t
17	10.0	12.8	4.2	4.2	14.2	17.0	2.4	3.0	sm	sm
18	10.0	12.7	3.6	4.1	13.6	16.8	2.8	3.1	sm	st

H: *L. hopei*. P: *L. peroffskeyana*. m: median-centromeric chromosome. sm: submedian-centromeric chromosome. st: subterminal-centromeric chromosome. t: terminal-centromeric chromosome.

Results and Discussion

Lepidozamia hopei and *L. peroffskeyana* showed the chromosome number of $2n=18$ (Fig. 1), which was consistent with those of the previous reports (Marchant 1964; Moretti 1990).

The karyotype of *L. hopei* consisted of eight median-, four submedian-, four subterminal- and two terminal-centromeric chromosomes (Figs. 2A and 3A, Table 2). On the other hand, those of *L. peroffskeyana* consisted of eight median-, two submedian-, six subterminal- and two terminal-centromeric chromosomes (Figs. 2B and 3B, Table 2). The arm ratio value of four submedian-centromeric chromosomes of *L. hopei* and two submedian-centromeric chromosomes of *L. peroffskeyana* were very close to 3.1, the lowest extreme of the arm ratio range of the subterminal-centromeric chromosome according to Levan *et al.* (1964). Thus, the karyotypes of two *Lepidozamia* species in the present investigation were very similar to each other, and was very similar to the karyotype of eight median-, four submedian- (or subterminal-) and two terminal-centromeric chromosomes of two *Lepidozamia* species reported by Marchant (1968) and Moretti (1990).

It has been reported that members of Cycadales show the stable karyotype within the each genus (Marchant 1968; Moretti 1990), excepting *Zamia* (Norstog 1980; Vovides 1983; Moretti and Sabato 1984) and *Bowenia* (Kokubugata *et al.* 1999). The present investigation indicated that the two *Lepidozamia* species are the same chromosome number, $2n=18$ and similar karyotype. These results




Fig. 3. Ideogram of two *Lepidozamia* species A. *L. hopei*. B. *L. paroffskyana*. Solid areas show long arm. Stippled areas show short arm.

intimate that there must be no karyomorphological arrangements in the speciation between two *Lepidozamia* species.

Acknowledgments

We thank Dr. K. Hill, National Herbarium of New South Wales, for his interest and helpful suggestion. This research was partly supported by the Sasakawa Scientific Research Grant from The Japan Science Society.

Summary

Somatic chromosomes at mitotic metaphase of *Lepidozamia hopei* and *L. paroffskyana* were investigated by the standard aceto-orcein staining method. They showed the chromosome number of $2n=18$ and the karyotype consisted of eight median-, eight subterminal- (or submedian-) and two terminal-centromeric chromosomes.

摘 要

Lepidozamia 属 2 種, *L. hopei* と *L. peroffskyana* の体細胞染色体をアセトオルセイン染色法により観察した。*Lepidozamia* 2 種は, ともに染色体数が $2n=18$ であり, 核型は 8 個の中部動原体型染色体, 8 個の次端部 (或いは次中部) 動原体型染色体, 2 個の端部動原体型染色体から構成されていた。

References

- Jones, D. L., 1993. Cycads of the world, Smithsonian Institution Press, Washington, D. C.
- Kokubugata G., K. Kondo, G. W. Wilson, L. M. Randall, A. van der Schans and D. K. Morris, 1999. Comparison of karyotype and rDNA-distribution in somatic chromosomes of *Bowenia* species (Stangeriaceae, Cycadales). Aust. Pl. Syst. Bot. **13**: in press.
- Levan, A., K. Fredga and A. A. Sandberg, 1964. Nomenclature for centromeric position on chromosomes. Hereditas **52**: 201-220.
- Marchant, C. J., 1968. Chromosome patterns and nuclear phenomena in the cycad families Stangeriaceae and Zamiaceae. Chromosoma (Berl.) **24**: 100-134.
- Moretti, A., 1990. Cytotaxonomy of cycads. Mem. New York Bot. Gard. **57**: 114-122.
- , and S. Sabato, 1984. Karyotype evolution by centromeric fission in *Zamia* (Cycadales). Pl. Syst. Evol. **146**: 215-223.
- Norstog, K., 1980. Chromosome numbers in *Zamia* (Cycadales). Caryologia **33**: 419-428.
- Stevenson, D. W., 1992. A formal classification of the extant cycads. Brittonia **44**: 220-223.
- Vovides, A., 1983. Systematic studies on the Mexican Zamiaceae. I. Chromosome numbers and karyotypes. Amer. J. Bot. **70**: 1002-1006.