

U–Pb age of the *Sphenoceras schmidt* Zone (middle Campanian, Cretaceous) in Hokkaido, northern Japan

Yasunari Shigeta¹ and Yukiyasu Tsutsumi¹

¹ Department of Geology and Paleontology, National Museum of Nature and Science,
4–1–1 Amakubo, Tsukuba, Ibaraki 305–0005, Japan

Abstract Radiometric age dating of zircons ($^{238}\text{U}/^{206}\text{Pb}$ ratios, using the LA-ICP-MS method) obtained from a tuff sample just below the *Sphenoceras schmidt*-bearing beds of the Yezo Group in the Nakagawa area, Hokkaido, northern Japan, yielded an age of 80.2 ± 0.8 Ma, 95% conf., which suggests an early middle Campanian (Late Cretaceous) age. This result indicates that the age of the *S. schmidt* Zone is the same in both southwestern and northern Japan. Thus, *Sphenoceras schmidt* should be regarded as an ideal zonal-index fossil for the lower middle Campanian of the Northwest Pacific region.

Key words: Campanian, Cretaceous, *Sphenoceras schmidt*, U–Pb age, zircon

Introduction

The inoceramid bivalve *Sphenoceras schmidt* (Michael, 1899) occurs abundantly in the Upper Cretaceous of southwestern Japan (Noda *et al.*, 1996; Misaki and Maeda, 2009). Because its characteristic shell sculpture, e.g. divergent ribs, is easy to recognize, it has been regarded as an ideal zonal-index fossil (Toshimitsu *et al.*, 1995). Kodama (1990) correlated the *S. schmidt* Zone in the Izumi Group with the uppermost part of polarity chron C33r, and Tsutsumi *et al.* (2014) reported an age of 79.7 ± 0.7 Ma (95% conf.) for a tuff in beds containing *S. schmidt* in the Himenoura Group. In consideration of this evidence, Shigeta *et al.* (2016) correlated the *S. schmidt* Zone with the lower middle Campanian based on the three-subdivision scheme for the Campanian.

Sphenoceras schmidt is also common in the Upper Cretaceous of the Yezo Group in Hokkaido, northern Japan (Nagao and Matsumoto, 1940) and Sakhalin, Far Eastern Russia (Zonova *et al.*, 1993). Kodama *et al.* (2000) recognized 13 magnetostratigraphic zones in the Upper Cretaceous in the Naiba area, southern Sakhalin, and concluded that the *S. schmidt*-bearing beds correspond to the bottom of polarity chron C32r (= middle upper Campanian). In contrast, Tamaki *et al.* (2008) correlated the *S. schmidt*-bearing beds in the Urakawa area, Hokkaido with the upper part of polarity chron C33r and lower part of

C33n (= lower middle Campanian). Thus, the magnetostratigraphic correlation of the *S. schmidt* Zone in the Yezo Group is still a matter of debate and the radiometric age of the zone had never been determined until now.

In this paper, we calibrate the international correlation of the *Sphenoceras schmidt* Zone with the zircon-based geochronology of the Yezo Group in the Nakagawa area, Hokkaido, and establish a more precise global correlation of the zone.

Notes on stratigraphy

Even though the Yezo Group in the Nakagawa area is complexly folded and faulted, a continuous Upper Cretaceous succession is well exposed along the Abeshinai River and its tributaries (e.g. Matsumoto, 1942; Takahashi *et al.*, 2003). Within the group, the Osoushinai Formation yields numerous relatively well-preserved Campanian megafossils from various horizons.

The 600-m thick Osoushinai Formation consists mainly of dark gray to gray siltstone, sandy siltstone and muddy sandstone, intercalated with 1–100 cm thick white, vitric tuff beds (Takahashi *et al.*, 2003; Fig. 1). The lower and middle parts contain the inoceramid *Sphenoceras naumanni* (Yokoyama, 1890) and ammonoids *Yokoyamaoceras ishikawai* (Jimbo, 1894), *Gaudryceras tenuiliratum* Yabe, 1903 and *Polyptychoceras pseudogaultinum*




Fig. 1. Index map showing distribution of the Yezo Group (black areas) in Hokkaido (A), locality (B, C) and stratigraphic horizon (D) of the tuff bed that provided the U–Pb zircon age in the Nakagawa area, Hokkaido, northern Japan.

(Yokoyama, 1890) (Fig. 1D). Characterizing the upper part are gigantic specimens of the inoceramid *S. schmidtii*, which occur abundantly in calcareous concretions as well as in the host rock (Matsuda and Ubukata, 1999; Takahashi *et al.*, 2003), together with the ammonoids *Urakawites* Matsumoto, 1955, *Teshioites* Matsumoto, 1955 and *Canadoceras* Spath, 1922 (Matsumoto, 1954, 1955).

Material and methods

Material: Radiometric age analysis was conducted on zircons extracted from a tuff sample taken from a 100-cm thick white, vitric tuff bed positioned immediately below the *Sphenoceras orientalis*-bearing beds, which occur about 18m below the *S. schmidtii*-bearing beds in an exposure of the Osoushinai Formation along the Abeshinai

River (44°40'23.90"N, 142°1'15.55"E; Fig. 1).

Methods: The zircon grains were extracted by standard techniques: crushing, heavy liquid separation and handpicking. Then, the zircon grains were mounted in an epoxy disc with the FC1 zircon standard ($^{206}\text{Pb}/^{238}\text{U} = 0.1859$; Paces and Miller, 1993), and the SRM610 glass standard and polished until the center of each grain was exposed. The backscattered electron and cathodoluminescence images of the zircon grains were used for selection of the sites for analysis. U–Pb dating of the sample was carried out using the LA-ICP-MS method, which consists of a NWR213, a 213 nm wave length Nd-YAG laser ablation system (Electro Scientific Industries) and an Agilent 7700x quadrupole ICP-MS (Agilent Technologies), installed at the National Museum of Nature and Science at Tsukuba, Japan. The experimental conditions and procedures for the measure-

Table 1. LA-ICP-MS analyzed data and calculated ages of the tuff sample from the upper part of the Osoushinai Formation. Errors are 1 sigma. Pb_c and Pb* indicate the common and radiogenic portions, respectively.

Labels	²⁰⁶ Pb _c ⁽¹⁾ (%)	U (ppm)	Th (ppm)	Th/U	²³⁸ U/ ²⁰⁶ Pb* ⁽¹⁾	²⁰⁷ Pb*/ ²⁰⁶ Pb* ⁽¹⁾	²³⁸ U/ ²⁰⁶ Pb* age ⁽¹⁾ (Ma)	²³⁸ U/ ²⁰⁶ Pb* age ⁽²⁾ (Ma)
Abe_01	1.64	183	72	0.40	76.30 ± 2.50	0.0505 ± 0.0138	83.9 ± 2.7	83.7 ± 2.7
Abe_02	0.54	406	126	0.32	81.01 ± 1.81	0.0418 ± 0.0064	79.1 ± 1.8	79.6 ± 1.7
Abe_03	0.48	209	111	0.55	80.38 ± 2.41	0.0474 ± 0.0116	79.7 ± 2.4	79.8 ± 2.3
Abe_04	0.11	207	84	0.41	84.89 ± 2.47	0.0497 ± 0.0115	75.5 ± 2.2	75.3 ± 2.2
Abe_05	0.68	290	186	0.66	81.98 ± 2.55	0.0455 ± 0.0102	78.2 ± 2.4	78.4 ± 2.4
Abe_06	0.77	173	74	0.44	80.42 ± 2.47	0.0430 ± 0.0135	79.7 ± 2.4	80.1 ± 2.5
Abe_07	0.29	492	254	0.53	79.85 ± 1.77	0.0479 ± 0.0070	80.2 ± 1.8	80.2 ± 1.7
Abe_08	0.77	117	66	0.58	78.96 ± 3.48	0.0469 ± 0.0216	81.1 ± 3.6	81.2 ± 3.3
Abe_09	2.01	170	111	0.67	86.84 ± 3.05	0.0477 ± 0.0172	73.8 ± 2.6	73.8 ± 2.4
Abe_10	0.94	490	262	0.55	78.26 ± 1.68	0.0425 ± 0.0084	81.8 ± 1.8	82.4 ± 1.7
Abe_11	0.13	353	209	0.61	77.93 ± 1.87	0.0535 ± 0.0090	82.2 ± 2.0	81.6 ± 1.9
Abe_12	0.81	172	95	0.57	80.24 ± 2.64	0.0474 ± 0.0125	79.8 ± 2.6	79.9 ± 2.6
Abe_13	2.64	137	77	0.58	79.78 ± 3.36	0.0378 ± 0.0215	80.3 ± 3.4	81.3 ± 3.2
Abe_14	0.79	196	94	0.49	82.22 ± 2.77	0.0437 ± 0.0132	77.9 ± 2.6	78.3 ± 2.6
Abe_15	0.73	364	147	0.42	79.86 ± 1.92	0.0442 ± 0.0077	80.2 ± 1.9	80.6 ± 1.9
Abe_16	0.17	474	158	0.34	73.89 ± 1.55	0.0410 ± 0.0060	86.7 ± 1.8	86.7 ± 1.7
Abe_17	0.13	503	455	0.93	81.53 ± 1.73	0.0520 ± 0.0095	78.6 ± 1.7	78.1 ± 1.5
Abe_18	1.89	171	102	0.61	80.39 ± 2.85	0.0525 ± 0.0176	79.7 ± 2.8	79.1 ± 2.6
Abe_19	0.23	312	157	0.52	79.30 ± 2.42	0.0488 ± 0.0098	80.8 ± 2.5	80.6 ± 2.4
Abe_20	1.02	375	299	0.82	79.10 ± 2.20	0.0381 ± 0.0115	81.0 ± 2.2	81.7 ± 2.0
Abe_21	0.01	139	69	0.51	78.72 ± 3.43	0.0431 ± 0.0218	81.4 ± 3.5	81.3 ± 3.1
Abe_22	0.00	258	118	0.47	78.71 ± 2.09	0.0507 ± 0.0061	81.4 ± 2.1	80.9 ± 2.2
Abe_23	0.60	663	269	0.42	80.39 ± 1.58	0.0499 ± 0.0060	79.7 ± 1.6	79.3 ± 1.5
Abe_24	0.00	117	62	0.54	77.36 ± 3.24	0.0540 ± 0.0109	82.8 ± 3.5	82.0 ± 3.6
Abe_25	0.25	190	138	0.75	87.08 ± 3.21	0.0505 ± 0.0177	73.6 ± 2.7	73.2 ± 2.4
Abe_26	1.08	99	50	0.52	76.07 ± 3.76	0.0623 ± 0.0325	84.2 ± 4.1	82.5 ± 4.1
Abe_27	1.09	137	76	0.57	79.24 ± 3.51	0.0573 ± 0.0213	80.8 ± 3.6	79.8 ± 3.4
Abe_28	2.14	335	174	0.53	78.39 ± 1.95	0.0255 ± 0.0093	81.7 ± 2.0	83.4 ± 1.9
Abe_29	1.60	226	125	0.57	82.76 ± 2.43	0.0378 ± 0.0117	77.4 ± 2.3	78.3 ± 2.2
Abe_30	0.00	99	52	0.54	73.16 ± 3.02	0.0500 ± 0.0126	87.5 ± 3.6	87.2 ± 3.8
Abe_31	2.62	100	50	0.52	83.43 ± 4.85	0.0441 ± 0.0358	76.8 ± 4.4	77.1 ± 3.7
Abe_32	0.33	139	74	0.55	79.67 ± 3.32	0.0504 ± 0.0188	80.4 ± 3.3	80.0 ± 3.1

(1) Common Pb corrected by assuming ²⁰⁶Pb/²³⁸U–²⁰⁸Pb/²³²Th age-concordance(2) Common Pb corrected by assuming ²⁰⁶Pb/²³⁸U–²⁰⁷Pb/²³⁵U age-concordance

ments were based on the methods described in Tsutsumi *et al.* (2012). The spot size of the laser was 25 μm. Corrections for common Pb was made on the basis of the measured ²⁰⁷Pb/²⁰⁶Pb ratio (²⁰⁷Pb correction) or ²⁰⁸Pb/²⁰⁶Pb and Th/U ratios (²⁰⁸Pb correction) (e.g. Williams, 1998) as well as the model for common Pb compositions proposed by Stacey and Kramers (1975). In this paper, we adopt the ²⁰⁷Pb correction for age discussion because it is more effective in calculating the Phanerozoic ²³⁸U–²⁰⁶Pb* age than the ²⁰⁸Pb correction (e.g. Williams, 1998). The pooled ages presented in this study were calculated using Isoplot/Ex software (Ludwig, 2003). The uncertainties in the mean ²³⁸U–²⁰⁶Pb* ages represent 95% confidence intervals (95% conf.). ²⁰⁶Pb* indicates radiometric ²⁰⁶Pb.

Results

Analyzed data from zircons; common ²⁰⁶Pb, U, and Th concentrations, Th/U, ²³⁸U/²⁰⁶Pb*, and ²⁰⁷Pb*/²⁰⁶Pb* ratios, and radiometric ²³⁸U/²⁰⁶Pb* ages are listed in Table 1. All errors state 1 sigma level. All zircons in the samples show rhythmic oscillatory and/or sector zoning on cathodoluminescence images, which is commonly observed in igneous zircons (e.g. Corfu *et al.*, 2003), and their higher Th/U ratios (>0.1) also support their igneous origin (Williams and Claesson, 1987; Schiøtte *et al.*, 1988; Kinny *et al.*, 1990; Hoskin and Black, 2000). Figure 2 shows a Tera-Wasserberg concordia diagram and an age distribution plot for all analyzed spots of the sample. All zircon U–Pb ages cluster in the range 73–87 Ma. The weighted mean age, after the rejection of three data points, yield 80.2 ± 0.8 Ma (MSWD = 0.8; 95% conf.), which is thought to




Fig. 2. U–Pb zircon ages of tuff sample collected from the upper part of the Osoushinai Formation. A, cathodoluminescence image (CL) of zircon grains from the sample. Circles on the grains represent spots analyzed by LA-ICP-MS. Spots are 25 μm across. B, Tera-Wasserburg U–Pb concordia diagram of zircons. C, ^{238}U – ^{206}Pb age distribution plot of tuff sample. $^{207}\text{Pb}^*$ and $^{206}\text{Pb}^*$ indicate radiometric ^{207}Pb and ^{206}Pb , respectively.

represent the magmatism/deposition age of the tuff sample.

Discussion

Zircon geochronology in this study reveals that the U–Pb age of the tuff just below the *Sphenoceramus schmidtii*-bearing beds is 80.2 ± 0.8 Ma (95% conf.), which infers an early middle Campanian age. This result confirms the validity of the work of Tamaki *et al.* (2008), who correlated the beds in the Urakawa area, Hokkaido with the upper part of polarity chron C33r and lower part of C33n (= lower middle Campanian). In contrast, our result differs from the work of Kodama *et al.* (2000), who correlated the *S. schmidtii*-bearing beds in the Naiba area, Sakhalin with the bottom of polarity chron C32r (= middle upper Campanian). We question the accuracy of this magnetostratigraphic correlation primarily because strata just below the *S.*

schmidtii-bearing beds are not well exposed in the Naiba area (Shigeta *et al.*, 2015, 2016).

The *Sphenoceramus schmidtii* Zone in southwestern Japan was shown to be lower middle Campanian by both magnetostratigraphy and zircon geochronology (Kodama, 1990; Tsutsumi *et al.*, 2014), thus suggesting that the age of the zone in southwestern Japan is the same as in Hokkaido. Therefore, *Sphenoceramus schmidtii* should be regarded as an ideal zonal-index fossil for the lower middle Campanian of the Northwest Pacific region.

Sphenoceramus schmidtii also occurs in the Upper Cretaceous of the Pacific coastal region of North America (e.g. Usher, 1952; Matsumoto, 1960; Jones, 1963). Ward *et al.* (1983, 2012) indicated that the *S. schmidtii* Zone of the Great Valley Sequence in northern California should be placed near the top of polarity chron C34n (= Santonian) and lowest part of polarity chron C33r (= lowest Campanian), which differs greatly from that of the

Northwest Pacific region. We question the accuracy of this conclusion because the magnetostratigraphy of northern California has only been studied in very short stratigraphic sequences. Further magnetostratigraphic studies of a continuous sequence spanning the Santonian to Maastrichtian as well as radiometric age determinations are needed settle this problem.

Acknowledgments

We thank Yasuyuki Tsujino (Tikushima Prefectural Museum, Tokushima) for valuable comments on the first draft. We are very grateful to Toshihiro Sakai (Asahikawa, Hokkaido) for his help during field work. Thanks are extended to Jim Jenks (West Jordan, Utah) for his helpful suggestions and improvement of the English text. This research was financially supported by the National Museum of Nature and Science project, Chemical Stratigraphy and Dating as a Clue for Understanding the History of the Earth and Life.

References

- Corfu, F., Hanchar, J. M., Hoskin, P. W. O. and Kinny, P. (2003) An atlas of zircon textures. In: Hanchar, J. M. and Hoskin, P. W. O. (Eds.), *Zircon: Reviews in Mineralogy and Geochemistry*. Mineralogical Society of America, Washington D.C., pp. 278–286.
- Hoskin, P. W. and Black, L. P. (2000) Metamorphic zircon formation by solid-state recrystallization of protolith igneous zircon. *Journal of Metamorphic Geology*, **18**: 423–439.
- Jimbo, K. (1894) Beiträge zur Kenntniss der Fauna der Kreideformation von Hokkaido. *Palaeontologische Abhandlungen (new series)*, **2**: 1–48.
- Jones, D. (1963) Upper Cretaceous (Campanian and Maastrichtian) ammonites from southern Alaska. *U. S. Geological Survey Professional Paper*, **432**: 1–53.
- Kinny, P. D., Wijbrans, J. R., Froude, D. O., Williams, I. S. and Compston, W. (1990) Age constraints on the geological evolution of the Narryer Gneiss Complex, Western Australia. *Australian Journal of Earth Sciences*, **37**: 51–69.
- Kodama, K. (1990) Magnetostratigraphy of the Izumi Group along the Median Tectonic Line in Shikoku and Awaji Islands, Southwest Japan. *Journal of the Geological Society of Japan*, **96**: 265–278. (in Japanese with English abstract)
- Kodama, K., Maeda, H., Shigeta, Y., Kase, T. and Takeuchi, T. (2000) Magnetostratigraphy of Upper Cretaceous strata in South Sakhalin, Russian Far East. *Cretaceous Research*, **21**: 469–478.
- Ludwig, K. R. (2003) User's manual for Isoplot 3.00. A Geochronological Toolkit for Microsoft Excel. Berkeley Geochronology Center Special Publication No. 4. 70 pp. Berkeley Geochronology Center, Berkeley.
- Matsuda, M. and Ubukata, T. (1999) Variation of shell sculpture in a Cretaceous bivalve *Sphenoceras* (Inoceramidae). *Geoscience Reports of Shizuoka University*, **26**: 1–15. (In Japanese with English abstract)
- Matsumoto [= Matumoto], T. (1942) Fundamentals in the Cretaceous stratigraphy of Japan, Part 1. *Memoirs of the Faculty of Science, Kyushu Imperial University, Series D, Geology*, **1**: 129–280.
- Matsumoto, T. (1954) The Cretaceous System in the Japanese Islands. 324 pp. Japan Society for the Promotion of Science, Tokyo.
- Matsumoto, T. (1955) The bituberculate pachydiscids from Hokkaido and Saghalien. *Memoirs of the Faculty of Science, Kyushu University, Series D, Geology*, **5**: 153–184.
- Matsumoto, T. (1960) Upper Cretaceous ammonites of California. Part 3. *Memoirs of the Faculty of Science, Kyushu University, Series D, Geology*, Special vol. **2**: 1–204.
- Michael, R. (1899) Ueber Kreidefossilien von der Insel Sachalin. *Jahrbuch der Königlich Preussischen Geologischen Landesanstalt und Bergakademie zu Berlin*, **18**: 153–164.
- Misaki, A. and Maeda, H. (2009) Lithostratigraphy and biostratigraphy of the Campanian–Maastrichtian Toyajo Formation in Wakayama, southwestern Japan. *Cretaceous Research*, **30**: 1398–1414.
- Nagao, T. and Matsumoto, T. (1940) A monograph of the Cretaceous *Inoceramus* of Japan, Part II. *Journal of the Faculty of Science, Hokkaido University, Series IV, Geology and Mineralogy*, **6**: 1–64.
- Noda, M., Ohtsuka, M., Kano, M. and Toshimitsu, S. (1996) The Cretaceous inoceramids from the Mifune and Hime-noura groups in Kyushu. *Geological Society of Oita, Special issue*, **2**: 1–63. (In Japanese)
- Paces, J. B. and Miller, J. D. (1993) Precise U–Pb ages of Duluth Complex and related mafic intrusions, northeastern Minnesota: geochronological insights to physical, petrogenetic, paleomagnetic, and tectonomagmatic processes associated with the 1.1 Ga midcontinent rift system. *Journal of Geophysical Research*, **98**: 13997–14013.
- Schiøtte, L., Compston, W. and Bridgwater, D. (1988) Late Archaean ages for the deposition of clastic sediments belonging to the Malene supracrustals, southern West Greenland: evidence from an ion probe U–Pb zircon study. *Earth and Planetary Science Letters*, **87**: 45–58.
- Shigeta, Y., Izukura, M., Nishimura, T. and Tsutsumi, Y. (2016) Middle and late Campanian (late Cretaceous) ammonoids from the Urakawa area, Hokkaido, northern Japan. *Paleontological Research*, **20**: 322–366.
- Shigeta, Y., Nishimura, T. and Nifuku, K. (2015) Middle and late Maastrichtian (latest Cretaceous) ammonoids from the Akkeshi Bay area, eastern Hokkaido, northern Japan and their biostratigraphic implications. *Paleontological Research*, **19**: 107–127.
- Spath, L. F. (1922) On the Senonian ammonite fauna of Pondoland. *Transactions of the Royal Society of South Africa*, **10**: 113–148.

- Stacey, J. S. and Kramers, J. D. (1975) Approximation of terrestrial lead isotope evolution by a two-stage model. *Earth and Planetary Science Letters*, **26**: 207–221.
- Takahashi, A., Hirano, H. and Sato, T. (2003) Stratigraphy and fossil assemblage of the Upper Cretaceous in the Teshionakagawa area, Hokkaido. *Journal of the Geological Society of Japan*, **109**: 77–95. (In Japanese with English abstract)
- Tamaki, M., Oshimbe, S. and Itoh Y. (2008) A large latitudinal displacement of a part of Cretaceous forearc basin in Hokkaido, Japan: paleomagnetism of the Yezo Supergroup in the Urakawa area. *Journal of the Geological Society of Japan*, **114**: 207–217
- Toshimitsu, S., Matsumoto, T., Noda, M., Nishida, T. and Maiya, S. (1995) Towards an integrated mega-, micro- and magneto-stratigraphy of the Upper Cretaceous in Japan. *Journal of the Geological Society of Japan*, **101**: 19–29. (In Japanese with English abstract)
- Tsutsumi, Y., Horie, K., Sano, T., Miyawaki, R., Momma, K., Matsubara, S., Shigeoka, M. and Yokoyama, K. (2012) LA-ICP-MS and SHRIMP ages of zircons in chevkinite and monazite tuffs from the Boso Peninsula, Central Japan. *Bulletin of the National Museum of Nature and Science, Series C*, **38**: 15–32.
- Tsutsumi, Y., Miyake, Y. and Komatsu, T. (2014) LA-ICP-MS zircon U-Pb dating of acidic tuffs from the Upper Cretaceous Himenoura Group, Koshiki-jima area, southwest Japan. *Abstract with Programs, the 2014 Annual Meeting of the Palaeontological Society of Japan*. p. 25. (In Japanese)
- Usher, J. L. (1952) Ammonite faunas of the Upper Cretaceous rocks of Vancouver Island, British Columbia. *Geological Survey of Canada Bulletin*, **21**: 1–182.
- Ward, P., Verosub, K. L., and Haggart, J. W. (1983) Marine magnetic anomaly 33–34 identified in the Upper Cretaceous of the Great Valley Sequence of California. *Geology*, **11**: 90–93.
- Ward, P. D., Haggart, J. W., Mitchell, R., Kirschvink, J. L. and Tobin, T. (2012) Integration of macrofossil biostratigraphy and magnetostratigraphy for the Pacific Coast Upper Cretaceous (Campanian–Maastrichtian) of North America and implications for correlation with the Western Interior and Tethys. *Geological Society of America Bulletin*, **124**: 957–974.
- Williams, I. S. (1998) U–Th–Pb geochronology by ion microprobe. In: McKibben M. A., Shanks W. C. P. and Ridley W. I. (Eds.), *Applications of Microanalytical Techniques to Understanding Mineralizing Processes*. Reviews in Economic Geology **7**, Society of Economic Geologists, Littleton, CO., pp. 1–35
- Williams, I. S. and Claesson, S. (1987) Isotopic evidence for the Precambrian provenance and Caledonian metamorphism of high grade paragneisses from the Seve Nappes, Scandinavian Caledonides. *Contributions to Mineralogy and Petrology*, **97**: 205–217.
- Yabe, H. (1903) Cretaceous Cephalopoda from the Hokkaido. Part 1: *Lytoceras*, *Gaudryceras*, and *Tetragonites*. *Journal of the College of Science, Imperial University of Tokyo*, **18**: 1–55.
- Yokoyama, M. (1890) Versteinerungen aus der japanischen Kreide. *Palaeontographica*, **36**: 159–202.
- Zonova, T. D., Kazintsova, L. I. and Yazykova, E. A. (1993) Atlas of Index Fossils in the Cretaceous Fauna of Sakhalin. 327 pp. Nedra, St. Petersburg. (In Russian)