

Flavonol Glycosides from the Whitish Flowers of *Primula alpicola* and *P. sikkimensis* var. *hopeana* in Bhutan

Takayuki Mizuno^{1,*}, Rinchen Yangzom², Hari Prasad Devkota³,
Yoshinori Murai¹, Rinchen Dorji², Kencho Dorji², Choki Wangmo²,
Choki Gyeltshen² and Tsukasa Iwashina¹

¹ Department of Botany, National Museum of Nature and Science,
4–1–1 Amakubo, Tsukuba, Ibaraki 305–0005, Japan

² National Biodiversity Centre, Ministry of Agriculture and Forests, Serbithang, Thimpu, 873, Bhutan

³ Graduate School of Pharmaceutical Sciences, Kumamoto University,
5–1 Oe-honmachi, Kumamoto 862–0973, Japan

*E-mail address: tmizuno@kahaku.go.jp

(Received 11 August 2020; accepted 23 September 2020)

Abstract Five flavonols were isolated from the whitish flowers of *Primula alpicola* and *P. sikkimensis* var. *hopeana* which belong to the section *Sikkimensis* in Bhutan. Flavonoid composition of two *Primula* species were qualitatively the same and they were identified as quercetin 3-*O*-xylosyl-(1→2)-[rhamnosyl-(1→6)-glucoside] (**1**), quercetin 3-*O*-rutinoside (**2**), quercetin 3-*O*-glucoside (**3**), kaempferol 3-*O*-rutinoside (**4**) and isorhamnetin 3-*O*-rutinoside (**5**). Compound **1** which might characterize in these species was reported from the family Primulaceae for the first time. Although gossypetin has been reported as principle yellow pigment in *Primula* species, it was not found in the whitish flowers of two *Primula* taxa.

Key words: flavonol, flower color expression, *Primula alpicola*, *Primula sikkimensis* var. *hopeana*, quercetin, section *Sikkimensis*.

Introduction

The genus *Primula* is widely distributed in the Northern Hemisphere and consists of ca. 500 species. In this genus, Himalaya-Hengduan and its adjacent regions have been considered to represent the modern diversification center (Richards, 1993). The species of section *Sikkimensis* having campanulate flowers are mainly distributed in the Himalayas and western China. Their flowers are colorful from purple, violet, yellow to white (Aitken *et al.*, 1999).

Some flavonoids have been reported from the flowers of *Primula* species (e.g. Karl *et al.*, 1981; Saito *et al.*, 1990; Hashimoto *et al.*, 2015; Apel *et al.*, 2017). In section *Sikkimensis*, flower fla-

vonoids have been reported to understand the color expression and their systematics (Harborne, 1968, 1969). According to Harborne (1968, 1969), yellow pigment of four *Primula* species, i.e. *P. alpicola* (W.W.Smith) Stapf, *P. florindae* Kingdon-Ward, *P. sikkimensis* Hook.f. and *P. waltonii* Watt ex I.B.Balfour has been identified as gossypetin (3,5,7,8,3',4'-hexahydroxyflavone) but not carotenoid (Harborne, 1968). Thus, it was suggested that carotenoids are replaced to gossypetin in their *Primula* flowers as major yellow pigment (Harborne, 1969). Furthermore, gossypetin was found at least in *P. velis* L., *P. vulgaris* Hudson and *P. elatior* (L.) Hill of the two sections *Sikkimensis* and *Primula* (Harborne, 1969). Pigments of cyanic flower species in the section *Sikkimensis* were isolated from the flowers of *P. secundiflora* Franch and *P. ioessa* W.W.Sm., and

anthocyanins, malvidin and delphinidin, were identified, respectively (Harborne, 1968).

In this study, we isolated and identified the flavonols from two *Primula* taxa, *P. alpicola* and *P. sikkimensis* var. *hopeana* (Balfour f. & Cooper) W.W.Smith & Fletcher having whitish flowers in the section Sikkimensis. *Primula alpicola* is growing in stream sides of 3600–4100 m elev. in southeastern Tibet, Bhutan and northeastern India, and has campanulate white, yellow or purple flower (Aitken *et al.*, 1999). On the other hand, *P. sikkimensis* var. *hopeana* having campanulate ivory white flowers is mainly growing in wet meadows or scrub of 3600–4700 m elev. in southeastern Tibet, Bhutan and eastern Nepal (Aitken *et al.*, 1999). Moreover, we performed aglycone analysis using crude extracts for the elucidation of the flower color expression and discussed the white color expression of *Primula* species of the section Sikkimensis.

Materials and Methods

Plant materials

Fresh flowers of *Primula alpicola* (Fig. 1A) were collected between Zizingoma–Omta Tsho and tent site of Omta Tsho, Thampe La zone, Bhutan on 30 June 2017. On the other hand, *P. sikkimensis* var. *hopeana* (Fig. 1B) were collected between Yakuchu–Wangzhi, Black Mountains, Bhutan on 31 July 2018. Species were identified by one of the authors (R. Yangzom) and the voucher specimens were deposited in the Herbarium of National Biodiversity Centre (THIM), Bhutan. Flowers were immediately dried in collection sites.

Extraction and isolation

Dry flowers of *P. alpicola* (15.1 g) and *P. sikkimensis* var. *hopeana* (10.0 g) were extracted with MeOH for an overnight at room temperature. After filtration and concentration, the extracts were applied to column chromatography on Diaion HP-20SS (Mitsubishi Chemical Co., Japan) eluted with 200 mL 0.5% TFA, 200 mL 10% MeCN containing 0.5% TFA and 200 mL

25% MeCN containing 0.5% TFA. Separated fractions were subjected to Sephadex LH-20 column chromatography eluted with 70% MeOH. The fractions were purified by preparative HPLC with Shimadzu HPLC systems equipped with a SPD-20A UV–Vis detector using Inertsil ODS-4 (I.D. 10 × 250 mm, GL Science Inc., Japan), at a flow-rate of 1.5–3.0 mL min^{−1}; injection of 350–400 μL; eluent of 15 or 25% MeCN containing 5% HCOOH.

Identification of the flavonoids

Isolated flavonols were identified by UV–Vis spectral survey (500–220 nm) according to Mabry *et al.* (1970), LC-MS, acid hydrolysis, and direct TLC (BAW, 15% HOAc and BEW) and HPLC comparisons with authentic samples. In addition, **1** was performed sugar analysis by HPLC, and NMR measurement.

HPLC analysis was performed with Shimadzu HPLC systems equipped with a SPD-20A UV–Vis detector using InertSustain C18 (I.D. 4.6 × 250 mm, GL Science Inc.), at a flow-rate of 1.0 mL min^{−1}. Water containing 5% HCOOH (solvent A) and 90% MeCN containing 5% HCOOH (solvent B) were used as the mobile phase under gradient elution condition. The gradient solution program was as follows: a linear gradient from 10 to 50% solvent B for 35 min, 50% solvent B for 10 min.

Liquid chromatograph-mass spectra (LC-MS) was performed with Shimadzu LC-MS systems equipped with a SPD-20A UV–Vis detector using Inertsil ODS-4 (I.D. 2.1 × 100 mm, GL Science Inc.), at a flow-rate of 0.2 mL min^{−1}, and the eluent with 12% MeCN containing 5% HCOOH. ESI⁺ 4.5 kV and ESI[−] 3.5 kV, 250°C.

NMR spectra (¹H NMR, ¹³C NMR, ¹H-¹H COSY, TOCSY, HMQC and HMBC) were recorded on a Bruker AV-600 in pyridine-*d*₅ at 600 MHz (¹H NMR) and 150 MHz (¹³C NMR). Chemical shifts (δ) were given in ppm.

Sugar identification was performed to hydrolysate of **1** according to Tanaka *et al.* (2007) and Mizuno *et al.* (2015). After acid hydrolysis by 0.5 mM HCl, sugars were separated from the




Fig. 1. Habitat of *Primula alpicola* (A, Tent site of Omta Tsho, Thampe La zone, Bhutan, 30 June 2017) and *P. sikkimensis* var. *hopeana* (B, Between Yakuchu and Wangzhi, Black Mountains, Bhutan, 31 July 2018).

aglycone with Sep-pack C18 and the neutralized with WAX column (Waters Corp., Milford, MA). The residues were dissolved in pyridine (50 μ L) containing L-cysteine methyl ester hydrochloride

(10 mg mL^{-1}) and heated at 60°C for 60 min. The solution was added to 50 μ L pyridine containing *o*-tolyl isothiocyanate (10 mg mL^{-1}), and then heated at 60°C for 60 min. The reaction solution

and authentic sugars were analyzed using column, Cosmosil 5C18-AR-II (4.6 × 250 mm, Nacalai Tesque Inc., Japan). Elution was performed with 25% MeCN containing 0.2% H₃PO₄ at a flow rate of 0.8 mL min⁻¹, and detection at 250 nm.

Aglycone analysis

Crude extracts (0.1 mL) were added into 12% aq. HCl and treated for 30 min at 100°C for acid hydrolysis. After cooling, the solutions were diluted with water and were applied to Sep-pack C18 cartridge eluted with 2 mL H₂O, 2 mL 10% MeCN containing 0.5% TFA and 2 mL MeOH. The organic fractions were evaporated and applied to HPLC analysis with Shimadzu HPLC system equipped with PDA detector using Inertsil Sustain C18 (I.D. 4.6 × 250 mm), at a flow-rate of 1.0 mL min⁻¹. The gradient solution program of the mobile phase was as follows: a linear gradient from 20 to 100% solvent B for 16 min, 100% solvent B for 6 min.

Authentic samples

Origins of the authentic samples were as follows; quercetin 3-*O*-glucoside, quercetin 3-*O*-rutinoside and kaempferol 3-*O*-rutinoside from the aerial parts and fruits of *Osyris alba* L. (Santalaceae) (Iwashina *et al.*, 2008), isorhamnetin 3-*O*-rutinoside from the leaves of *Asarum caulescens* Maxim. (Aristolochiaceae) (Iwashina *et al.*, 2005) (TNS collection number **87-1**), quercetin from the flowers of *Astrophytum ornatum* (DC.) Web. (Cactaceae) (Iwashina *et al.*, 1988) (TNS collection number **81-3**), kaempferol from hydrolysate of kaempferol 3-*O*-glucoside from *Camptotheca acuminata* Decne. leaves (Cornaceae) (Iwashina and Hatta, 1998) (TNS collection number **77-14**), and isorhamnetin from hydroyrsate of isorhamnetin 3-*O*-rhamnosylglucoside from *Epiphyllum hybrida* flowers (Cactaceae) (Iwashina, unpublished data).

Quercetin 3-*O*-xylosyl-(1→2)-[rhamnosyl-(1→6)-glucoside] (**1**, Fig. 2). TLC: R_f 0.50 (BAW), 0.71 (15%HOAc), 0.58 (BEW); Color UV (365 nm): dark purple, UV/NH₃: yellow.

HPLC: *t*R (min) 15.5. UV: λ_{max} (nm) MeOH 258, 357; + NaOMe 278, 331, 414 (inc.); + AlCl₃ 273, 412; + AlCl₃/HCl 270, 297, 361, 398; + NaOAc 271, 389; + NaOAc/H₃BO₃ 262, 379; LC-MS: *m/z* 743 [M + H]⁺, 741 [M - H]⁻; Sugar analysis: *t*R (min): 16.1, 18.6 and 27.2 min (authentic samples: D-glucose, 16.1 min, D-xylose, 18.6, L-rhamnose, 27.2 min).

Quercetin 3-*O*-rutinoside (**2**, rutin, Fig. 3). TLC: R_f 0.59 (BAW), 0.48 (15%HOAc), 0.66 (BEW); Color UV (365 nm): dark purple, UV/NH₃: yellow. HPLC: *t*R (min) 18.1. UV: λ_{max} (nm) MeOH 257, 357; + NaOMe 275, 414 (inc.); + AlCl₃








Fig. 2. Quercetin 3-*O*-xylosyl-(1→2)-[rhamnosyl-(1→6)-glucoside] (**1**). A = Chemical structure. B = Significant HMBC correlations were shown as the greyed lines.

Fig. 3. Quercetin 3-*O*-rutinoside (**2**, rutin).Fig. 5. Kaempferol 3-*O*-rutinoside (**4**, nicotiflorin).Fig. 4. Quercetin 3-*O*-glucoside (**3**, isoquercitrin).Fig. 6. Isorhamnetin 3-*O*-rutinoside (**5**, narcissin).

273, 420; + AlCl₃/HCl 268, 297, 362, 398; + NaOAc 262, 409; + NaOAc/H₃BO₃ 271, 306, 399; LC-MS: *m/z* 611 [M + H]⁺, 609 [M - H]⁻.

Quercetin 3-*O*-glucoside (**3**, isoquercitrin, Fig. 4). TLC: R_f 0.67 (BAW), 0.28 (15%HOAc), 0.73 (BEW); Color UV (365 nm): dark purple, UV/NH₃: yellow. HPLC: *t*R (min) 19.1. UV: λ_{max} (nm) MeOH 257, 358; + NaOMe 275, 325, 408 (inc.); + AlCl₃ 267, 369, 402; + AlCl₃/HCl 267, 297, 360, 403; + NaOAc 269, 372; + NaOAc/H₃BO₃ 262, 379; LC-MS: *m/z* 465 [M + H]⁺, 463 [M - H]⁻.

Kaempferol 3-*O*-rutinoside (**4**, nicotiflorin, Fig. 5). TLC: R_f 0.70 (BAW), 0.54 (15%HOAc),

0.76 (BEW); Color UV (365 nm): dark purple, UV/NH₃: dark greenish yellow. HPLC: *t*R (min) 20.5. UV: λ_{max} (nm) MeOH 265, 351; + NaOMe 277, 322, 397 (inc.); + AlCl₃ 268, 303, 351, 406sh.; + AlCl₃/HCl 271, 301, 348, 387; + NaOAc 270, 306, 364; + NaOAc/H₃BO₃ 265, 354; LC-MS: *m/z* 595 [M + H]⁺, 593 [M - H]⁻.

Isorhamnetin 3-*O*-rutinoside (**5**, narcissin, Fig. 6). TLC: R_f 0.68 (BAW), 0.54 (15%HOAc), 0.73 (BEW); Color UV (365 nm): dark purple, UV/

NH₃: yellow. HPLC: *t*R (min) 21.1. UV: λ_{max} (nm) MeOH 255, 356; + NaOMe 276, 324, 405 (inc.); + AlCl₃ 266, 301, 358, 411sh.; + AlCl₃/HCl 266, 302, 358, 402sh.; + NaOAc 270, 371; + NaOAc/H₃BO₃ 255, 360; LC-MS: *m/z* 623 [M – H][–].

Results and Discussion

HPLC chromatograms of flower extracts of *Primula alpicola* and *P. sikkimensis* var. *hopeana* were shown in Fig. 7. Flavonoid composition was qualitatively the same to each other. Of five flavonoids isolated from two *Primula* species, **1** was mainly identified by NMR, together with LC-MS, acid hydrolysis and UV-Vis spectra. On the other hand, **2–5** were characterized by LC-MS, UV-Vis spectra, and direct HPLC comparisons with authentic samples.

LC-MS of **1** showed the molecular ion peaks at *m/z* 743 [M + H]⁺ and 741 [M – H][–] and frag-

ment ion peaks, *m/z* 611 [M – 132 + H]⁺, 465 [M – 132 – 146 + H]⁺ and 303 [M – 132 – 146 – 162 + H]⁺, showing the attachment of each 1 mol of hexose, rhamnose and pentose to pentahydroxyflavone. Quercetin, D-glucose, L-rhamnose and D-xylose were liberated by acid hydrolysis of **1**. In NMR analysis of **1**, four aromatic proton signals, δ_{H} 8.33 (1H, *d*, H-2'), 8.23 (1H, *dd*, H-6'), 7.39 (1H, *d*, H-5'), 6.67 (1H, *d*, H-6) and 6.60 (1H, *d*, H-8), appeared (Table 1). The β -pyranose structure of glucose was shown by the coupling constant (*J* = 7.7 Hz) of anomeric proton signal (Markham and Geiger, 1994). HMBC correlation between Glc-1 proton and δ_{C} 134.7 corresponding to C-3 of quercetin showed that glucose is attached to 3-position of quercetin. The α -pyranose structure of rhamnose was shown by the ³*J*_{H,C} correlation of δ_{H} 5.24 (1H, *brs*, Rham-1) with δ_{C} 69.6 (Rham-5) and occurrence as broad-singlet of anomeric proton signal. HMBC correlation between Rham-1 proton and




Fig. 7. HPLC analysis of the flowers of *Primura alpicola* (A) and *P. sikkimensis* var. *hopeana* (B). **1** = quercetin 3-*O*-xylosyl-(1→2)-[rhamnosyl-(1→6)-*O*-glucoside] (15.5 min), **2** = quercetin 3-*O*-rutinoside (18.1 min), **3** = quercetin 3-*O*-glucoside (19.1 min), **4** = kaempferol 3-*O*-rutinoside (20.5 min), **5** = isorhamnetin 3-*O*-rutinoside (21.1 min).

Table 1. ^1H and ^{13}C NMR data of **1** from *P. sikki-mensis* var. *hopeana*

Position	Pyridine- d_5	
	δH	δC
<i>Quercetin</i>		
2		157.6
3		134.7
4		178.7
5		162.9
6	6.67 <i>d</i> , 1.8	99.6
7		165.5
8	6.60 <i>d</i> , 1.8	94.0
9		157.5
10		105.4
1'		123.1
2'	8.33 <i>d</i> , 2.0	117.8
3'		146.8
4'		150.4
5'	7.39 <i>d</i> , 8.3	116.3
6'	8.23 <i>dd</i> , 8.3, 2.1	122.8
<i>3-O-Glucose</i>		
1	6.42 <i>d</i> , 7.7	100.5
2	4.44 <i>dd</i> , 7.6, 8.8	82.4
3	4.34 <i>m</i>	78.6
4	4.34 <i>m</i>	72.5
5	4.03 <i>m</i>	77.3
6a	4.43 <i>dd</i> , 13.2, 2.9	68.0
6b	4.93 <i>m</i>	
<i>6"-O-Rhamnose</i>		
1	5.24 <i>brs</i>	102.3
2	4.34 <i>m</i>	72.1
3	3.95 <i>t</i> , 8.7	71.3
4	4.11 <i>m</i>	73.9
5	4.14 <i>m</i>	69.6
Me	1.47 <i>d</i> , 5.9	
<i>2"-O-Xylose</i>		
1	5.67 <i>d</i> , 6.2	105.7
2	4.24 <i>m</i>	74.9
3	4.24 <i>m</i>	77.1
4	4.21 <i>m</i>	71.0
5a	4.54 <i>dd</i> , 11.2, 4.1	66.6
5b	3.84 <i>dd</i> , 11.2, 8.5	

C-6 of glucose at δ_{C} 68.0 showed that rhamnose is attached to 6-position of glucose. Furthermore, β -xylopyranose was shown by the $^3J_{\text{H,C}}$ correlation of δ_{H} 4.54 (1H, *dd*, Xyl-5) with δ_{C} 105.7 (Xyl-1) and coupling constant ($J=6.2\text{ Hz}$) of anomeric proton signal. It was proved by HMBC (correlation of Xyl-1 proton and C-2 carbon signal of glucose at δ_{C} 82.4) that xylose is attached to 2-position of glucose. Significant HMBC correlations are shown in Fig. 2B. Thus, **1** was iden-

tified as quercetin 3-*O*- β -D-xylopyranosyl-(1 $\rightarrow$ 2)-*O*-[α -L-rhamnopyranosyl-(1 $\rightarrow$ 6)- β -D-glucopyranoside]. This flavonol glycoside has already been reported from the leaves of *Orixa japonica* Thunb. (Rutaceae) (Nishida *et al.*, 1990) and *Actinidia arguta* (Sieb. & Zucc.) var. *giraldii* (Diels) Vorosch. (Actinidiaceae) (Webby, 1991).

UV spectral properties showed that **2** and **3** are flavonols having free 5-, 7-, 3'- and 4'-hydroxyl and a substituted 3-hydroxyl groups. LC-MS of **2** showed that the molecular ion peaks, m/z 611 $[\text{M} + \text{H}]^+$ and 609 $[\text{M} - \text{H}]^-$, and fragment ion peaks, m/z 465 $[\text{M} - 146 + \text{H}]^+$ and 303 $[\text{M} - 146 - 162 + \text{H}]^+$, suggesting that **2** is quercetin which attached each 1 mol of hexose and rhamnose. LC-MS of **3** showed molecular ion peaks at m/z 465 $[\text{M} + \text{H}]^+$ and 463 $[\text{M} - \text{H}]^-$ and fragment ion peak at m/z 303 $[\text{M} - 162 + \text{H}]^+$, showing that the compound is quercetin which attached 1 mol hexose. Finally, they are identified as quercetin 3-*O*-rutinoside (**2**) and 3-*O*-glucoside (**3**) by direct HPLC and TLC comparisons with authentic samples.

Compound **4** was characterized as flavonol with free 5-, 7- and 4'-hydroxyl and a substituted 3-hydroxyl groups by UV spectral survey. LC-MS of **4** showed the molecular ion peaks at m/z 595 $[\text{M} + \text{H}]^+$ and 593 $[\text{M} - \text{H}]^-$, and fragment ion peaks at m/z 449 $[\text{M} - 162 + \text{H}]^+$ and m/z 287 $[\text{M} - 162 - 146 + \text{H}]^+$, suggesting that **4** is kaempferol which attached each 1 mol of hexose and rhamnose. From the results of HPLC and TLC comparison with authentic sample, **4** was identified as kaempferol 3-*O*-rutinoside.

Compound **5** was characterized as 5-, 7- and 4'-hydroxylated, and 3- and 3'-substituted flavonol by UV spectral survey. Since the molecular ion peak, m/z 623 $[\text{M} - \text{H}]^-$, and fragment ion peak, m/z 317 $[\text{M} - 162 - 146 + \text{H}]^+$ was shown by LC-MS, **5** is tetrahydroxy-monomethoxyflavone which attached each 1 mol of hexose and rhamnose. Finally, **5** was identified as isorhamnetin 3-*O*-rutinoside by HPLC and TLC comparison with authentic sample.

Some flavonols which were isolated from the

Primula species have been shown to be flavonols which were linearly attached sugars to 3-position, e.g. quercetin and isorhamnetin 3-*O*-glucosyl-(1→2)-gentiobiosides from the leaves of *Primula auricula* L. (Fico *et al.*, 2007.), quercetin 3-*O*-arabinosyl-(1→2)-arabinosyl-(1→4)-glucoside from the leaves of *Primula turkestanica* (Regel) E.A.White (Zakharov *et al.*, 1970), and quercetin and kaempferol 3-*O*-xylosyl-(1→2)-glucosyl-(1→6)-glucosides and kaempferol 3-*O*-glucosyl-(1→2)-glucosyl-(1→6)-glucoside from the leaves and flowers of *Primula sieboldii* (Hashimoto *et al.*, 2015). In this study, triglycosylated flavonol, quercetin 3-*O*-xylosyl-(1→2)-[rhamnosyl-(1→6)-glucoside] (**1**), was identified as branched flavonol glycoside. Although many *Primula* species are not surveyed for flavonoids, we presumed that **1** is the characteristic flavonol of *P. sikkimensis* var. *hopeana*. Flavonoid **1** from *Orixa japonica* was reported as oviposition deterrent of a Rutaceae-feeding swallowtail butterfly, *Papilio xuthus* (Nishida *et al.*, 1990). However, the function in *Primula* flowers is unknown.

Aglycone analyses of two *Primula* species by HPLC were shown in Fig. 8. These peaks were compared with authentic flavonol aglycones, i.e. quercetin, kaempferol and isorhamnetin. The proportion of each flavonol was followed, quercetin (Qu) 63.8%, kaempferol (Km) 17.7% and isorhamnetin (Is) 10.5% in *P. alpicola*, and Qu 82.8%, Km 4.0% and Is 9.2% in *P. sikkimensis* var. *hopeana*. Quercetin was highest amount than those of kaempferol and isorhamnetin in both *Primula* species. Thus, it was shown that major flavonoid in their species is quercetin (each 63.8% and 82.85%).

Yellow color due to 6- or 8-hydroxylated flavonols such as gossypetin and quercetagenin has been reported from *Centaurea* (Asteraceae) (Mishio *et al.*, 2006, 2015), *Rudbeckia* (Asteraceae) (Thompson *et al.*, 1972), *Eriophyllum* (Asteraceae) (Harborne and Smith, 1978), *Gossypium* (Malvaceae) (Harborne, 1966) and *Tagetes* flowers (Asteraceae) (Morita, 1957; Bhardwaj *et al.*, 1980). Harborne (1969) reported that




Fig. 8. Aglycone analysis of the flowers of *Primura alpicola* and *P. sikkimensis* var. *hopeana*. These aglycones were compared with authentic flavonols, quercetin (Qu, 11.7 min), kaempferol (Km, 13.2 min) and isorhamnetin (Is, 13.5 min) by HPLC.

yellowish *Primula* flowers in section Sikkimensis, are due to gossypetin. In this study, we isolated five flavonols, i.e. quercetin 3-*O*-xylosyl-(1→2)-[rhamnosyl-(1→6)-glucoside], 3-*O*-rutinoside and 3-*O*-glucoside, and kaempferol and isorhamnetin 3-*O*-rutinosides from two whitish flowers of *Primula* species.

Quercetin as yellow pigment has been reported from the yellow flowers of *Astrophytum* species (Cactaceae) (Iwashina *et al.*, 1988) and *Clematis* cultivars and species (Ranunculaceae) (Hashimoto *et al.*, 2008), *Chimonanthus praecox* (Calycanthaceae) (Iwashina *et al.*, 2001) and *Hibiscus rosa-sinensis* (Malvaceae) (Subramanian and Nair, 1972). In this study, the high quercetin concentration in the flowers of two *Primula* species was revealed by aglycone analysis. However, their *Primula* flowers are white to ivory white. As the results, we presumed that quercetin amount is quantitatively insufficient for expression of yellow.

Acknowledgments

The authors gratefully thank to Mrs. Aki Kiuchi, part-time assistant of National Museum of Nature and Science, Japan, for the technical assistance. This work was supported by Discretionary expenses of Director General of the National Museum of Nature and Science.

References

- Aitken, E., Grierson, A. J. C. and Ling, D. G. 1999. Primulaceae. In: Grierson, A. J. C. and Long, D. G. (Eds.), *Flora of Bhutan* Volume 2, part 2. pp. 515–568. Royal Botanic Garden Edinburgh, Edinburgh.
- Apel, L., Kammerer, D. R., Stintzing, F. C. and Spring, O. 2017. Comparative metabolite profiling of triterpenoid saponins and flavonoids in flower color mutations of *Primula veris* L. *International Journal of Molecular Sciences* 18: 153.
- Bhardwaj, D. K., Bisht, M. S., Uain, S. C., Mehta, C. K. and Sharma, G. C. 1980. Quercetagetin 5-methyl ether from the petals of *Tagetes patula*. *Phytochemistry* 19: 713–714.
- Fico, G., Rodondi, G., Flamini, G., Passarella, D. and Tomé, F. 2007. Comparative phytochemical and morphological analyses of three Italian *Primula* species. *Phytochemistry* 68: 1683–1691.
- Harborne, J. B. 1966. Comparative biochemistry of flavonoids—I. Distribution of chalcone and aurone pigments in plants. *Phytochemistry* 5: 111–115.
- Harborne, J. B. 1968. Comparative biochemistry of the flavonoids — VII, Correlations between flavonoid pigmentation and systematics in the family Primulaceae. *Phytochemistry* 7: 1215–1230.
- Harborne, J. B. 1969. Gossypetin and herbacetin as taxonomic markers in higher plants. *Phytochemistry* 8: 177–183.
- Harborne, J. B. and Smith, D. M. 1978. Anthochlors and other flavonoids as honey guides in the Compositae. *Biochemical Systematics and Ecology* 6: 287–291.
- Hashimoto, M., Iwashina, T. and Matsumoto, S. 2008. Flower color and flavonoid variation in *Clematis patens* (Ranunculaceae). *Bulletin of the National Museum of Nature and Science, Series B* 34: 43–51.
- Hashimoto, N., Ohsawa, R., Kitajima, J. and Iwashina, T. 2015. New flavonol glycosides from the leaves and flowers of *Primula sieboldii*. *Natural Product Communications* 10: 421–423.
- Iwashina, T. and Hatta, H. 1998. Flavonoids in the leaves of *Mastixia trichotoma*, *Camptotheca acuminata*, *Alangium platanifolium* var. *trilobum*, *A. premnifolium* and *Davidia involucrata*, and their implication with the genus *Cornus*. *Annals of the Tsukuba Botanical Garden* 17: 139–146.
- Iwashina, T., Konta, F. and Kitajima, J. 2001. Anthocyanins and flavonols of *Chimonanthus praecox* (Calycanthaceae) as flower pigments. *The Journal of Japanese Botany* 76: 166–172.
- Iwashina, T., López-Sáez, J. A. and Kitajima, J. 2008. Flavonoids from *Osyris alba*. *Biochemical Systematics and Ecology* 36: 146–147.
- Iwashina, T., Ootani, S. and Hayashi, K. 1988. On the pigmented spherical bodies and crystals in tepals of Cactaceous species in reference to the nature of betalains or flavonols. *The Botanical Magazine, Tokyo* 101: 175–184.
- Iwashina, T., Kitajima, J., Shiuchi, T. and Itou, Y. 2005. Chalcones and other flavonoids from *Asarum sensu lato* (Aristolochiaceae). *Biochemical Systematics and Ecology* 33: 571–584.
- Karl, C., Müller, G. and Pedersen, P. A. 1981. Flavonoids in the flowers of *Primula officinalis*. *Planta Medica* 41: 96–99.
- Mabry, T. J., Markham, K. R. and Thomas, M. B. 1970. *The Systematic Identification of Flavonoids*, Springer, New York.
- Markham, K. R. and Geiger, H. 1994. ^1H Nuclear magnetic resonance spectroscopy of flavonoids and their glycosides in hexadeuterodimethylsulfoxide. In: Harborne, J. B. (ed.), *The Flavonoids: Advances in Research since 1986*, pp. 441–498. Chapman & Hall, London.
- Mishio, T., Honma, T. and Iwashina, T. 2006. Yellow flavonoids in *Centaurea ruthenica* as flower pigments. *Biochemical Systematics and Ecology* 34: 180–184.
- Mishio, T., Takeda, K. and Iwashina, T. 2015. Anthocyanins and other flavonoids as flower pigments from eleven *Centaurea* species. *Natural Product Communications* 10: 447–450.

- Mizuno, T., Kamo, T., Sasaki, N., Yada, H., Murai, Y. and Iwashina, T. 2015. Novel C-xylosylflavones from the leaves and flowers of *Iris gracilipes*. *Natural Product Communications* 10: 441–444.
- Morita, N. 1957. Flavonoids of the flowers and the leaves of *Tagetes erecta*. *Yakugaku Zasshi* 77: 31–33.
- Nishida, R., Ohsugi, T., Fukami, H. and Nakajima, S. 1990. Oviposition deterrent of a Rutaceae-feeding swallowtail butterfly, *Papilio xuthus*, from a non-host Rutaceous plant, *Orixa japonica*. *Agricultural and Biological Chemistry* 54: 1265–1270.
- Richards, J. 1993. *Primula*. Timber Press, Portland.
- Saito, N., Yoda, K., Haruyama, H., Kuwano, H. and Honda, T. 1990. Structure determination of a violet-blue flower flavonoid quercetin 3-glucosyl (1→2) gentiobioside from *Primula polyantha*. *Heterocycles* 30: 759–764.
- Subramanian, S. S. and Nair, A. G. R. 1972. Flavonoids of four Malvaceous plants. *Phytochemistry* 11: 1518–1519.
- Tanaka, T., Nakashima, T., Ueda, T., Tomii, K. and Kouno, I. 2007. Facile discrimination of aldose enantiomers by reversed phase HPLC. *Chemical and Pharmaceutical Bulletin* 55: 899–901.
- Thompson, W. R., Meinwald, J., Aneshansley, D. and Eisner, T. 1972. Flavonols: pigments responsible for ultraviolet absorption in nectar guide of flower. *Science* 177: 528–530.
- Webby, R. F. 1991. A flavonol triglycoside from *Actinidia arguta* var. *giraldii*. *Phytochemistry* 30: 2443–2444.
- Zakharov, A. M., Glyzin, V. I. and Bankovskii, A. I. 1970. Primflaside: A new flavonol glycoside from *Primula turkestanica*. *Chemistry of Natural Compounds* 6: 486.