

Gaudryceras tombetsense Matsumoto, a Maastrichtian ammonoid from the Aridagawa area, Wakayama, southwestern Japan

YASUNARI SHIGETA¹, AKIHIRO MISAKI² and MASAOKI OHARA³

¹Department of Geology and Paleontology, National Museum of Nature and Science, 4-1-1 Amakubo, Tsukuba, Ibaraki 305-0005, Japan
(e-mail: shigeta@kahaku.go.jp)

²Kitakyushu Museum of Natural History & Human History, 2-4-1 Higashida, Yahatahigashi-ku, Kitakyushu, Fukuoka 805-0071, Japan

³Wakayama Prefectural Museum of Natural History, 370-1 Funao, Kainan, Wakayama 642-0001, Japan

Received December 8, 2011; Revised manuscript accepted March 14, 2012

Abstract. Discovery of *Gaudryceras tombetsense* Matsumoto, 1984 in the Futakawa Formation of the Sotoizumi Group in the Aridagawa area, Wakayama, southwestern Japan establishes that this formation includes sediments of early Late Maastrichtian age, which is the youngest record of the Sotoizumi Group distributed in Shikoku and the Kii Peninsula. The occurrence of the youngest fossils from the easternmost area of the Sotoizumi Group may support the suggestion that the depocenter of the Sotoizumi basin migrated eastward due to movement along the Kurosegawa Tectonic Zone (KTZ), which was still active during Campanian to Maastrichtian time.

Key words: ammonoid, Cretaceous, Futakawa Formation, *Gaudryceras tombetsense*, Sotoizumi Group, Wakayama

Introduction

Ammonoids attributable to *Gaudryceras* Grossouvre, 1894 are abundant in the Upper Cretaceous of the North Pacific realm and include strongly ornamented species that exhibit various ribbing styles, most of which have a restricted stratigraphic range (Matsumoto, 1995). Recent work involving the stratigraphy and fossil assemblages of the Upper Cretaceous of Sakhalin revealed that several species of *Gaudryceras* occur successively in Maastrichtian strata (Maeda *et al.*, 2005). Shigeta *et al.* (2010) demonstrated that *Gaudryceras izumiense* Matsumoto and Morozumi, 1980 is very useful for correlation of Lower Maastrichtian strata in Southwest Japan, Hokkaido and Alaska. Because of its widespread distribution in the circum-North Pacific realm, *Gaudryceras* is an ideal ammonoid for precise biostratigraphic correlation of Maastrichtian strata in this particular realm.

The Cretaceous Sotoizumi Group is widely distributed in the Aridagawa area, Wakayama, which is located in the Chichibu Belt in the Outer Zone of Southwest Japan. Various megafossils have been reported from the western

part and its biostratigraphy has been well documented (e.g. Matsumoto, 1947, 1954; Hirayama and Tanaka, 1956a, b; Obata and Ogawa, 1976; Tanaka, 1985; Ohara, 2005; Misaki *et al.*, 2008; Misaki and Maeda, 2009, 2010a, 2010b). In contrast, the combination of rare megafossils and a complicated geologic structure has precluded the establishment of a precise biostratigraphic correlation in the eastern part (Misaki and Ohara, 2012).

We have recently discovered several specimens referable to *Gaudryceras tombetsense* Matsumoto, 1984 in the Futakawa Formation of the eastern part of the Aridagawa area. These specimens are herein described and we discuss their biostratigraphic significance.

Geologic notes

Outcrops of the Futakawa Formation, which is the upper part of the Sotoizumi Group, are well exposed along a small tributary of the Aridagawa River in the western area of Kunohara (Figure 1). In the lower reaches of the tributary (Locs. 1–4), these outcrops include alternating beds of mudstone and sandstone, which dip 85–90°




Figure 1. Index maps (A and B) and route map (C) showing locality from which *Gaudryceras tombetsense* was collected. Modified after Taira (2001) and Yamakita and Otoh (2000a).

eastward and strike N45–60°E. The sandstones generally show turbiditic features such as graded bedding and current ripple lamination. Along the middle reaches (Locs. 5–12), exposures include dark gray bioturbated mudstones that dip 15–20° eastward and strike N15–20° W. The stratigraphic relationship between sediments in the lower and middle reaches is unclear because of poor exposures in this particular region. All specimens of *Gaudryceras tombetsense* were obtained from dark gray bioturbated mudstones at Loc. 6.

Paleontological description

Systematic descriptions basically follow the classification established by Klein *et al.* (2009). Morphological terms in the systematic description are those used in the Treatise on Invertebrate Paleontology (Moore, 1957). Quantifiers used to describe the shape of the ammonoid shell replicate those proposed by Matsumoto (1954, p. 246) and modified by Haggart (1989, table 8.1).

Institution abbreviations.—WMNH = Wakayama Prefectural Museum of Natural History, Kainan.




Figure 2. *Gaudryceras tombetsense* Matsumoto, 1984, from the Futakawa Formation at Loc. 6 in the Aridagawa area, Southwest Japan. **A**, WMNH-Ge-1140210134; **B**, WMNH-Ge-1140210132; **C**, WMNH-Ge-1140210136.

Superfamily Tetragonitoidea Hyatt, 1900
 Family Gaudryceratidae Spath, 1927
 Genus *Gaudryceras* Grossouvre, 1894

Type species.—*Ammonites mitis* Hauer, 1866.

Gaudryceras tombetsense Matsumoto, 1984

Figures 2–4

Gaudryceras sp. nov. Matsumoto, 1980, p. 289, pl. 46, figs. 3–5, pl. 47, figs. 3–6.

Gaudryceras tombetsense Matsumoto, 1984, p. 2, pl. 1, figs. 1–2, pl. 2, figs. 1–4; Shigeta *et al.*, 1999, pl. 6, fig. 1; Maeda *et al.*, 2005, p. 66, figs. 26.3, 27–31.

Gaudryceras hamanakense Matsumoto and Yoshida. Zonova, 1990, p. 33, pl. 1, figs. 1–3, pl. 2, figs. 1–5, pl. 3, figs. 1–6; Yazykova, 1992, p. 196, pl. 111, figs. 1–2; Yazykova, 1994, p. 292, pl. 3, figs. 2–3.

Gaudryceras denmanense Whiteaves. Yazykova, 1993, p. 153, pl. 88,

fig. 2; Yazykova, 1994, p. 291, pl. 6, fig. 3.

?*Gaudryceras* cf. *tombetsense* Matsumoto. Maeda *et al.*, 2005, p. 67, fig. 40.1–40.3; Tsujino *et al.*, 2010, p. 682, fig. 4.

Holotype.—GK.H5991, figured by Matsumoto (1984, p. 2, pl. 1, figs. 1–2), from the Maastrichtian Heitarozawa Formation of the Yezo Group exposed along the Kikusui-gawa Valley, a branch of the Tombetsu River, in the Nakatombetsu area, northern Hokkaido, Japan.

Material examined.—Eight specimens were collected from dark gray bioturbated mudstones at Loc. 6 along a small tributary of the Aridagawa River in the western area of Kunohara. Three specimens, WMNH-Ge-1140210132 (Figure 2B), 1140210134 (Figure 2A), 1140210136 (Figure 2C), are immature shells (60–70 mm in diameter), whereas the others are probably mature shells (over 150 mm in diameter). WMNH-Ge-1140210279 consists of a partly



Figure 3. *Gaudryceras tombetsense* Matsumoto, 1984, from the Futakawa Formation at Loc. 6 in the Aridagawa area, Southwest Japan. A, WMNH-Ge-1140210279; B, WMNH-Ge-1140210059-b. Black arrow indicates position of last septum.

preserved phragmocone and most of the body chamber (Figure 3A). WMNH-Ge-1140210135-b consists of the phragmocone and a partly preserved body chamber (Figure 4A). Specimens WMNH-Ge-1140210135-d (Figure 4B), WMNH-Ge-1140210059-b (Figure 3B) and WMNH-Ge-1140210088 consist only of body chamber fragments. All specimens are significantly deformed and crushed.

Description.—Early whorls (up to 50 mm in diameter, Figure 2): Very evolute shell with moderately inflated whorl section characterized by arched venter and convex flanks. Umbilicus fairly wide with shallow umbilicus and rounded shoulders. Ornamentation consists of very fine, dense lirae, which arise at umbilical seam and approach a sigmoidal pattern before passing over the venter in a broad convex arch. Intercalation of lirae occurs on umbilical shoulder and lower flank. Each whorl has vari-

able close or distant, rounded, collarlike ribs, running parallel to lirae.

Middle whorls (50–100 mm in diameter, Figure 3A): As size increases, whorl section tends to become slightly compressed. Lirae gradually develop into slightly more distant, narrowly raised ribs, which increase in strength and frequency as diameter increases. These ribs follow the same sigmoidal pattern as those in the early whorls. Collarlike ribs become more frequent.

Later whorls (over 100 mm in diameter, Figures 3, 4): As shell grows larger, whorl section becomes even more compressed. Ribs become much coarser and more distant and collarlike ribs become more frequent. These ribs tend to become more sinuous as diameter increases. Suture line finely and deeply incised, partly visible (Figure 4A).

Remarks.—These specimens exhibit remarkable onto-




Figure 4. *Gaudryceras tombetsense* Matsumoto, 1984, from the Futakawa Formation at Loc. 6 in the Aridagawa area, Southwest Japan. **A**, WMNH-Ge-1140210135-b; **B**, WMNH-Ge-1140210135-d. Black arrow indicates position of last septum.

genetic changes in ribbing ornamentation, which is characteristic of *Gaudryceras tombetsense* as described by Matsumoto (1984) and Maeda *et al.* (2005). Therefore, we assign them with confidence to *G. tombetsense*.

As Matsumoto (1984, p. 4) pointed out, specimens described as *Gaudryceras* sp. nov. from the Sukumo-Nakamura area in western Shikoku, Japan (Matsumoto 1980) are conspecific with *G. tombetsense*. Specimens assigned to *Gaudryceras hamanakense* Matsumoto and Yoshida (1979) by Zonova (1990), Yazykova (1992) and Yazykova (1994) are identical to juvenile shells of *G. tombetsense*. A large specimen described as *G. denmanense* Whiteaves (1901) by Yazykova (1993) and Yazykova (1994) closely matches the adult shell of *G. tombetsense*.

Maeda *et al.* (2005, p. 67) and Tsujino *et al.* (2010) described *Gaudryceras* cf. *tombetsense* from Sakhalin and eastern Shikoku based respectively on a single juvenile specimen. However, juvenile shells of *Gaudryceras*

resemble each other and a proper species assignment can sometimes be very difficult. Their attribution of this specimen to *G. tombetsense* may be correct, but additional specimens showing the different growth stages are necessary in order to make a species assignment with a higher degree of confidence.

Inouye (1933) reported *Gaudryceras crassicoatum* (Jimbo, 1894) from a locality in the Kunohara area, which may in fact be the same fossil locality as ours. Hirayama and Tanaka (1956b) also listed *Gaudryceras*(?) sp. from the Itao area, about 3 km east of Kunohara. However, since they provide neither an illustration nor a detailed description, it is very difficult to restudy them.

Discussion

Gaudryceras tombetsense has been reported from the Maastrichtian of various areas in the Northwest Pacific,

e.g. Shikoku, Hokkaido and Sakhalin. Details of their stratigraphic position and associated fauna are understood very well in the Makarov area of southern Sakhalin (Maeda *et al.*, 2005).

Gaudryceras tombetsense occurs abundantly in Unit K2 of the Krasnoyarka Formation in the Makarov area together with inoceramid *Sphenoceras hetonaianus* (Matsumoto, 1952), which is diagnostic of the upper Lower to lower Upper Maastrichtian (Toshimitsu *et al.*, 1995). Maeda *et al.* (2005, Fig. 14.1, 14.4) illustrated a specimen that they attributed to *Pachydiscus gracilis* Matsumoto, 1979 from the same horizon, but we herein refer it to a peculiar morphotype belonging to *P. flexuosus* Matsumoto, 1979 by having a much compressed whorl section. *P. flexuosus* is normally indicative of the Upper Maastrichtian (Toshimitsu *et al.*, 1995). *P. flexuosus* seems to appear in the Lower Maastrichtian, while the stratigraphic range obviously extends upward to the Upper Maastrichtian (Maeda *et al.*, 2005; Maeda and Shigeta, 2005). Such associated fossils suggest that the *G. tombetsense*-bearing bed should be correlated with the lower Upper Maastrichtian.

The occurrence of *Gaudryceras tombetsense* in the Futakawa Formation in the Kunohara area demonstrates that this particular horizon is of early Late Maastrichtian age, which is the youngest record of the Sotoizumi Group distributed in Shikoku and the Kii Peninsula. This evidence supports the suggestion by Tashiro (1985, 1993) that the age of the Sotoizumi Group becomes younger toward the eastern part of the basin because our fossil locality occurs in the easternmost area of the Sotoizumi basin.

According to Taira *et al.* (1981, 1983), sediments of the Cretaceous Period, including the Sotoizumi Group, were deposited in a strike-slip basin along the Chichibu Belt formed by movement along the Kurosegawa Tectonic Zone (KTZ). Tashiro (1985, 1994) and Yamakita and Otoh (2000a) thought that the movement occurred mainly during the Early Cretaceous, but Misaki and Maeda (2009) pointed out that the movement was still active during Campanian to Maastrichtian time. The discovery of early Late Maastrichtian fossils in the Futakawa Formation may support the suggestion of Misaki and Maeda (2009) that the depocenter of the Sotoizumi basin migrated eastward by movement along the KTZ, which was still active during Campanian to Maastrichtian time.

The Campanian-Maastrichtian Izumi Group is distributed along the Median Tectonic Line (MTL) of south-central Japan. This group is considered to have been deposited in a strike-slip basin formed by left-lateral strike-slip movement along the MTL (e.g. Taira *et al.*, 1981, 1983; Miyata, 1990; Yamakita and Ito, 1999; Yamakita and Otoh, 2000b). The results of Misaki and Maeda (2009)

as well as those from this study suggest that there were two parallel strike-slip basins along the eastern margin of Asia during latest Cretaceous time. This suggestion may provide an important key for reconstruction of the geotectonic history of southwestern Japan during the Cretaceous.

Acknowledgments

We thank Yasuyuki Tsujino (Tokushima Prefectural Museum, Tokushima), René Hoffmann (Ruhr University, Bochum) and Haruyoshi Maeda (Kyushu University, Fukuoka) for their valuable comments on the first draft. We are also grateful to Kuniharu Kamei (Hannan, Osaka) and Yukio Sako (Kushimoto, Wakayama) for donating specimens and providing information about their collection locality. Thanks are extended to Jim Jenks (West Jordan, Utah) for his helpful suggestions and improvement of the English text. This study was supported by a grant from the National Museum of Nature and Science in 2011 to Y. Shigeta (No. 20112001).

References

- Grossouvre, A. de., 1894: Recherches sur la Craie supérieure. Deuxième partie: Paléontologie. Les ammonites de la Craie supérieure. *Mémoires du Service de la Carte Géologique Détaillée de la France* (1893), p. 1–264.
- Haggart, J. W., 1989: New and revised ammonites from the Upper Cretaceous Nanaimo Group of British Columbia and Washington State. *Geological Survey of Canada Bulletin*, vol. 396, p. 181–221.
- Hauer, F. R. von., 1866: Neue Cephalopoden aus den Gosaugebilden der Alpen. *Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften in Wien*, vol. 53, p. 300–308.
- Hirayama, K. and Tanaka, K., 1956a: *Explanatory Text of the Geological Map of Japan "Kainan"*, Scale 1:50,000, 62 p. Geological Survey of Japan, Kawasaki. (in Japanese with English abstract)
- Hirayama, K. and Tanaka, K., 1956b: *Explanatory text of the Geological Map of Japan "Todorogi"*, Scale 1:50,000, 37 p. Geological Survey of Japan, Kawasaki. (in Japanese with English abstract)
- Hyatt, A., 1900: Cephalopoda. In: Zittel K. A. ed., *Textbook of Palaeontology, English ed., translated by C. R. Eastman*, p. 502–592. Macmillan, London and New York.
- Inouye, J., 1933: Geology of the Aridagawa valley, Wakayama Prefecture (2). *Chikyu*, vol. 19, p. 415–431. (in Japanese)
- Jimbo, K., 1894: Beiträge zur Kenntniss der Fauna der Kreideformation von Hokkaido. *Palaeontologische Abhandlungen, Neue Folge*, Bd. 2, p. 1–48.
- Klein, J., Hoffmann, R., Joly, B., Shigeta, Y. and Vašiček, Z., 2009: *Fossilium Catalogus I: Animalia Pars 146, Lower Cretaceous Ammonites IV, Boreophylloceratoidea, Phylloceratoidea, Lytoceratoidea, Tetragonitoidea, Haploceratoidea including the Upper Cretaceous Representatives*, 416 p. Buckhuys Publishers, Leiden.
- Maeda, H. and Shigeta, Y., 2005: Maastrichtian ammonoid fauna from the Pugachevo area, southern Sakhalin, Russian Far East. *National Science Museum Monographs*, vol. 31, p. 121–136.
- Maeda, H., Shigeta, Y., Fernando, A. G. S. and Okada, H., 2005: Stratigraphy and fossil assemblages of the Upper Cretaceous System in the Makarov area, southern Sakhalin, Russian Far East. *National*

- Science Museum Monographs*, vol. 31, p. 25–120.
- Matsumoto, T., 1947: The geologic research of the Aritagawa valley, Wakayama Prefecture, a contribution to the tectonic history of the Outer Zone of south-west Japan. *Science Reports of the Faculty of Science, Kyushu University, Geology*, vol. 2, p. 1–12. (in Japanese)
- Matsumoto, T., 1952: Appendix. Notes on a new species of *Inoceramus*. *Journal of Geography*, vol. 61, p. 67–72. (in Japanese with English abstract)
- Matsumoto, T., 1954: *The Cretaceous System in the Japanese Islands*, 324 p. Japan Society for the Promotion of Science, Tokyo.
- Matsumoto, T., 1979: Palaeontological descriptions Part 1. Some new species of *Pachydiscus* from Tombetsu and the Hobetsu Valleys. *Memoirs of the Faculty of Science, Kyushu University, Series D, Geology*, vol. 24, p. 47–73.
- Matsumoto, T., 1980: Cephalopods from the Shimanto Belt of Kochi Prefecture (Shikoku). In: Taira, A. and Tashiro, M. eds., *Geology and Paleontology of the Shimanto Belt – Selected Papers in Honor of Prof. Jiro Katto*, p. 283–298. Rinyakosaikai Press, Kochi. (in Japanese)
- Matsumoto, T., 1984: Some gaudryceratid ammonites from the Campanian and Maastrichtian of Hokkaido. Part 1. *Science Report of the Yokosuka City Museum*, no. 32, p. 1–10.
- Matsumoto, T., 1995: Notes on gaudryceratid ammonites from Hokkaido and Sakhalin. *Palaeontological Society of Japan, Special Paper*, no. 35, p. 1–152.
- Matsumoto, T. and Morozumi, Y., 1980: Late Cretaceous ammonites from the Izumi Mountains, Southwest Japan. *Bulletin of the Osaka Museum of Natural History*, no. 33, p. 1–31.
- Matsumoto, T. and Yoshida, S., 1979: A new gaudryceratid ammonite from eastern Hokkaido. *Transactions and Proceedings of the Palaeontological Society of Japan, New Series*, no. 114, p. 65–76.
- Misaki, A., Kadota, H. and Maeda, H., 2008: Discovery of mid-Cretaceous ammonoids from the Aridagawa area, Wakayama, southwest Japan. *Paleontological Research*, vol. 12, p. 19–26.
- Misaki, A. and Maeda, H., 2009: Lithostratigraphy and biostratigraphy of the Campanian-Maastrichtian Toyajo Formation in Wakayama, southwestern Japan. *Cretaceous Research*, vol. 30, p. 1398–1414.
- Misaki, A. and Maeda, H., 2010a: Two Campanian (Late Cretaceous) nosteroceratid ammonites from the Toyajo Formation in Wakayama, Southwest Japan. In: Tanabe, K., Shigeta, Y., Sasaki, T. and Hirano, H. eds., *Cephalopods – Present and Past*, p. 223–231. Tokai University Press, Tokyo.
- Misaki, A. and Maeda, H., 2010b: Stratigraphy of the mid- to upper Cretaceous System in the Aridagawa area, Wakayama, Southwest Japan. *Island Arc*, vol. 19, p. 517–529.
- Misaki, A. and Ohara, M., 2012: Discovery of *Ainoceras* (Ammonoidea) from the Upper Cretaceous Futakawa Formation in the Aridagawa area, Wakayama, southwestern Japan. *Paleontological Research*, vol. 15, p. 240–246.
- Miyata, T., 1990: Slump strain indicative of paleoslope in Cretaceous Izumi sedimentary basin along Median Tectonic Line, Southwest Japan. *Geology*, vol. 18, p. 392–394.
- Moore, R. C., 1957: *Treatise on Invertebrate Paleontology, Part L, Mollusca 4, Cephalopoda, Ammonoidea*, 490 p. Geological Society of America, New York, and University of Kansas Press, Lawrence.
- Obata, I. and Ogawa, Y., 1976: Ammonite biostratigraphy of the Cretaceous Arida Formation, Wakayama Prefecture. *Bulletin of the National Science Museum, Series C*, vol. 2, p. 93–110. (in Japanese with English summary)
- Ohara, M., 2005: Lower Cenomanian bivalve fossils from the Cretaceous of Kibi area, Wakayama Prefecture, Japan. *Annual Report of Wakayama Prefectural Museum of Natural History*, no. 23, p. 37–43. (in Japanese)
- Shigeta, Y., Maeda, H., Uemura, K. and Solov'yov, A. V., 1999: Stratigraphy of the Upper Cretaceous System in the Kril'on Peninsula, South Sakhalin, Russia. *Bulletin of the National Science Museum, Series C*, vol. 25, p. 1–27.
- Shigeta, Y., Tanabe, K. and Izukura, M., 2010: *Gaudryceras izumiense* Matsumoto and Morozumi, a Maastrichtian ammonoid from Hokkaido and Alaska and its biostratigraphic implications. *Paleontological Research*, vol. 14, p. 202–211.
- Spath, L. F., 1927: Revision of the Jurassic cephalopod fauna of Kachh (Cutch), part 1. *Memoirs of the Geological Survey of India, Palaeontologia Indica, New Series*, vol. 9, p. 1–71.
- Taira, A., 2001: Tectonic evolution of the Japanese island arc system. *Annual Review of Earth and Planetary Sciences*, vol. 29, p. 109–134.
- Taira, A., Saito, Y. and Hashimoto, M., 1981: The fundamental processes of geologic evolution of Japan. *Kagaku*, vol. 51, p. 508–515. (in Japanese)
- Taira, A., Saito, Y. and Hashimoto, M., 1983: The role of oblique subduction and strike-slip tectonics in the evolution of Japan. In: Hilde, T. W. C. and Uyeda, S. eds., *Geodynamics of the western Pacific-Indonesian Regions. Geodynamics Series Volume 11*, p. 303–316. American Geophysical Union, Washington, D.C.
- Tanaka, K., 1985: Japanese Cretaceous fossils of Geological Survey of Japan. *Chishitsu News*, no. 371, p. 56–63. (in Japanese)
- Tashiro, M., 1985: The Cretaceous System of the Chichibu Belt in Shikoku – on the Early Cretaceous lateral fault in the Chichibu Belt. *Fossils (The Palaeontological Society of Japan)*, no. 38, p. 23–35. (in Japanese with English abstract)
- Tashiro, M., 1993: Bivalve faunas from Cretaceous of Japan – Part 1: On the bivalve faunas from the Chichibu and “Ryouke” Belts in Southwest Japan. *Research Reports of the Kochi University, Natural Science*, vol. 42, p. 105–155. (in Japanese with English abstract)
- Tashiro, M., 1994: Cretaceous tectonic evolution of southwest Japan from the bivalve faunal view-points. *Research Reports of the Kochi University, Natural Science*, vol. 43, p. 43–54. (in Japanese with English abstract)
- Toshimitsu, S., Matsumoto, T., Noda, M., Nishida, T. and Maiya, S., 1995: Towards an integrated mega-, micro- and magneto-stratigraphy of the Upper Cretaceous in Japan. *Journal of the Geological Society of Japan*, vol. 101, p. 19–29. (in Japanese with English abstract)
- Tsujino, Y., Ishida, K., Wada, H. and Hirayama, M., 2010: The second discovery of a Late Cretaceous ammonite from the Shimanto Belt in Mugi, Tokushima Prefecture, eastern Shikoku, Japan. *Journal of the Geological Society of Japan*, vol. 116, p. 680–685. (in Japanese with English abstract)
- Whiteaves, J., 1901: Note on a supposed new species of *Lytoceras* from the Cretaceous rocks at Denman Island, in the strait of Georgia. *Ottawa Naturalist*, vol. 15, p. 31–32.
- Yamakita, S. and Ito, T., 1999: A numerical model of the subsidence process of the Izumi sedimentary basin. *Structural Geology*, vol. 43, p. 1–9. (in Japanese with English abstract)
- Yamakita, S. and Otoh, S., 2000a: Cretaceous rearrangement processes of pre-Cretaceous geologic units of the Japanese Island by MTL-Kurosegawa left-lateral strike-slip fault system. *Memoirs of the Geological Society of Japan*, vol. 56, p. 23–38. (in Japanese with English abstract)
- Yamakita, S. and Otoh, S., 2000b: Estimation of the amount of Late Cretaceous left-lateral strike-slip displacement along the Median Tectonic Line and its implications in the juxtaposition of some geologic belts of southwest Japan. In: Ishiga, H., Takasu, A. and Hayasaka, Y. eds., *Tectonic division and correlation of terranes*

- of Southwest Japan, Monograph 49*, p. 49–104. Association for the Geological Collaboration in Japan, Tokyo. (*in Japanese with English abstract*)
- Yazikova [Yazykova], E. A., 1994: Maastrichtian ammonites and biostratigraphy of the Sakhalin and the Shikotan Islands, Far East Russia. *Acta Geologica Polonica*, vol. 44, p. 277–303.
- Yazykova, E. A., 1992: Upper Cretaceous ammonites from East USSR. *Trudy Vsesoyunogo Nauchno-Issledovatel'skogo Geologicheskogo Instituta (VSEGEI), New Series*, vol. 350, p. 192–200. (*in Russian; original title translated*)
- Yazykova, E. A., 1993: Class Cephalopoda. In, Zonova, T. D., Kazintsova, L. I. and Yazykova, E. A. eds., *Atlas of Index Fossils in the Cretaceous Fauna of Sakhalin*, p. 144–174. Nedra, St. Petersburg. (*in Russian; original title translated*)
- Zonova, T. D., 1990: The *Gaudryceras hamanakense* Zone in Sakhalin. In., Ablaev, A. G. ed., *New data on stratigraphy of Far East and Pacific Ocean*, p. 31–36. Akademiya Nauk SSSR, Dal'nevostochnoe Otdelenie, Tikhookeanskii Okeanologicheskii Institut, Vladivostok. (*in Russian; original title translated*)