

A New Species of *Zingiber* (Zingiberaceae) from Mt. Victoria, Western Myanmar

Mu Mu Aung¹, Nobuyuki Tanaka^{2,*} and Nao Miyake³

¹Forest Research Institute, Forest Department, Ministry of Environmental Conservation and Forestry, Yezin, Nay Pyi Taw, Myanmar

²Department of Botany, National Museum of Nature and Science, Amakubo 4–1–1, Tsukuba, Ibaraki 305–0005, Japan

³Laboratory of Plant Ecology, Division of Studies in Applied Sciences, Graduate School of Integrated Arts and Sciences, Kochi University, Akebono-cho 2–5–1, Kochi 780–8520, Japan

*E-mail: nobuyuki_tanaka@kahaku.go.jp

(Received 16 February 2017; accepted 22 March 2017)

Abstract A new species, *Zingiber matupiense* M.M.Aung & Nob.Tanaka (Zingiberaceae) is described from the north of Mt. Victoria, western Myanmar. This species is assignable to *Zingiber* Mill. section *Cryptanthium* Horan., and is similar to the *Z. cernuum* Dalz., and *Z. squarrosum* Roxb. However, it is markedly distinguished from these species by its erect leafy stem, yellowish white, recurved bracts, longer floral tubes, and bright yellow lateral staminodes.

Key words : Myanmar, new species, sect. *Cryptanthium*, *Zingiber*, Zingiberaceae.

Introduction

The genus *Zingiber* Mill. is the second largest genus of Zingiberaceae with 100–150 species (Kishor and Leong-Škorničková, 2013) and occurs from China and Japan through south and southeast Asia to northern Australia, with its center of diversity in monsoonal continental Asia (Larsen, 2005). So far 20 taxa of *Zingiber* have been recorded from Myanmar (Kress *et al.*, 2003; Tanaka, 2012a, b; Aung *et al.*, 2015). This small number, however, reflects the lack of recent explorations in Myanmar and possibility of more taxa in this region.

In the course of taxonomic study to revise the genus *Zingiber*, and to clarify their species diversity in Myanmar, field explorations in various sites covering representative regions were carried out from the years 2013 through 2015. During this exploration in 2013 to Natma Taung (local name of Mt. Victoria) National Park located in

Chin State, western Myanmar, the authors discovered a hitherto-undescribed species of *Zingiber*. The plant was growing at the margin of evergreen forest at 1,660 m altitude of the foothill of Mt. Victoria.

The plant is assignable to *Zingiber* sect. *Cryptanthium* Horan. on the basis of inflorescence on procumbent peduncles arising from the rhizome. After examination of its morphology based on the herbarium and spirit specimens and the study of related literature in the neighbouring regions, such as India, China and monsoonal southeast Asia (Dalzell, 1852; Baker, 1892; Gagnepain, 1908; Win, 1975; Wu and Larsen, 2000; Theilade, 1999; Larsen and Larsen, 2006; Tripathi and Singh, 2006; Chaveerach *et al.*, 2007; Kishor and Leong-Škorničková, 2013; Thongam and Kongsam, 2014; Triboun *et al.*, 2014; Leong-Škorničková *et al.*, 2015), it has been found that the present collection is an undescribed species similar to, *Z. cernuum* Dalz. in south India and *Z.*

squarrosus Roxb. in India, Nepal and Myanmar. Hence it is described here as a new species with figures based on living and herbarium materials.

Zingiber matupiense M.M.Aung & Nob. Tanaka, *sp. nov.* (Figs. 1 & 2).

Diagnosis: This new species is most similar to *Z. cernuum* Dalz., but differs by its erect leafy

stem, yellowish white recurved bracts with translucent margin, longer floral tubes, lateral staminodes and bright yellow anther crest. On the other hand, it differs from *Z. squarrosus* by its lanceolate, involute bracts and larger yellow labellum.

Rhizomatous herb, up to 1.2 m high. Rhizomes thick, 5 cm in diameter, externally pale brown,




Fig. 1. *Zingiber matupiense* M.M.Aung & Nob.Tanaka, *sp. nov.* A. Flower with recurved bracts. B. Rhizome. C. Leaves and leafy stem.




Fig. 2. Floral parts of *Zingiber matupiense* M.M.Aung & Nob.Tanaka, sp. nov. A. Bract. B. Bracteole. C. Calyx. D. Labellum. E. Lateral corolla lobes. F. Dorsal corolla lobe. G. Style and stamen. H. Epigynous glands. (Scale bar = 1 cm)

internally creamy white, slightly aromatic, roots many, strong and fleshy. Leafy shoots erect, with up to 20 leaves, basal 1/3 of pseudostem leafless, fleshy, the lower part reddish green, upper part green; bladeless sheaths 3 or 4, dark red. Leaves 4–10 pairs, petiole almost sessile, pale green, pulvinate at base; ligules white, membranous,

sometimes reddish on the lower leaves, c. 1 cm long, glabrous, apex round, scarious in mature; lamina broadly ovate to oblong-lanceolate, 20–40 cm long, 10–15 cm wide, almost smooth, glabrous, mid-vein basally enlarged, base cordate to cuneate, apex acute to acuminate; adaxially dark green, abaxially pale green, margin more or

less undulate. Inflorescence radical; peduncle procumbent, very short, up to 8 cm long, covered by sheaths, lower ones smaller than upper, glabrous; spike obovate to sub-globose, 8 cm long, 5 cm wide, consisting of 20–25 bracts, coriaceous. Bracts ovate-lanceolate, 4.5–5 cm long, 2–2.8 cm wide, outer bracts greenish white, inner bracts yellowish white, loosely imbricate, recurved, apex acute or shortly acuminate, glabrous, margin hyaline, semi-translucent white, enclosing single flower. Bracteoles lanceolate, white, 3.5 cm long, 1 cm wide, glabrous, apex shortly trilobed. Flowers 7–8 cm long, yellowish white; calyx 1.5–2 cm long, apex shortly tridentate, tubular, membranous, unilaterally split, glabrous; floral tube slender, 4.5–5 cm long, 0.6 cm in diameter at mouth, pale yellow to white towards the base, glabrous, corolla lobes subequal; dorsal corolla lobe c. 3 cm long, 1.2 cm wide, translucent cream at base, yellowish white at apex, ovate-lanceolate, curved; lateral corolla lobes c. 2.7 cm long, 0.7 cm wide, colored as the dorsal lobe; labellum broadly ovate, maroon with creamy yellow blotches, glabrous, central lobe oblong, c. 2.2 cm long, 1.3 cm wide, margin slightly undulate, apex slightly notched or emarginate; lateral staminodes 1.4 cm long, 0.5 cm wide, bright yellow, connate to labellum by basal 1/2–1/3. Anther yellow, c. 2.7 cm long with a crest, anther crest c. 1.4 cm long, wrapped around stigma; connective appendage c. 1.3 cm, yellow; thecae 1.3 cm long, parallel, dehiscing along the entire length. Style white, filiform, glabrous, stigma white with a circular apical aperture surrounded by stiff hairs. Epigynous glands two, c. 6 mm long, linear, pale yellow. Ovary c. 4 mm long, 2 mm wide, white, trilocular with many ovules on axile placenta, densely pubescent. Mature fruits not seen.

Type: MYANMAR. Chin State: Along the roadside between Taingsi village and Hakha road, near the foothill of Mt. Victoria, Matupi Township, Natma Taung National Park, 1,660 m altitude, 21°50.349'N 93°28.343'E, 12 July 2013, *Mu Mu Aung et al.* 094216 (holotype-TNS, isotype-RAF).

Phenology: Flowering from June to August; fruiting period unknown. Flowers open in the morning.

Etymology: The specific epithet is derived from the name of type locality, Matupi.

Distribution: Known only from the type locality.

Habitat and ecology: Evergreen forest margins in shade, c. 1,660 m above sea level.

Note: Although its character has some similarities to *Z. cernuum* Dalz. and *Z. squarrosum* Roxb., the key diagnostic characters are given in Table 1. *Zingiber cernuum* known as the “curved-stem ginger” (Cooke, 1908) which was described from South India and considered there to be a common endemic (Kanivita *et al.*, 2007). *Zingiber squarrosum* was described by Roxburgh (1810), and he mentioned that the plant was a native of Pegu (Bago Region), lower part of Myanmar. Gross morphology of the new species distinguishes it from *Z. cernuum* by erect leafy shoots, larger and ovate ligules, prominently recurved bracts, bright yellow lateral staminodes, and slightly notched or emarginated apex of the labellum.

Acknowledgments

This study was partially supported by Ri-Yo-Kan research grant from Graduate School of Integrated Arts and Sciences, Kochi University, and JSPS KAKENHI Grant Numbers 23770101 and K07207 (to N. Tanaka). Thanks are due to John Mood for his useful comments. Authors are grateful to Forest Department, Ministry of Natural Resources and Environmental Conservation of Myanmar for permission to conduct this study in Natma Taung National Park, and for their support and collaboration.

References

- Aung, M. M., Tanaka, N. and Miyake, N. 2015. Two Gingers, *Zingiber orbiculatum* and *Z. flavomaculosum* (Zingiberaceae), Newly Recorded from Myanmar. Bulletin of the National Museum of Nature and Science, Series B, 41(3): 1–6.

- Baker, J. G. 1892. *Zingiber*. In: Hooker, J. D. (ed.), *Flora of British India* 6, pp. 243–249. L. Reeve and Co, London, U.K.
- Cooke, T. 1908. *Zingiber cernuum*. In: *The Flora of the Presidency of Bombay* 2: 734. Taylor and Francis, London, U.K.
- Chaveerach, A., Mokkamul, P., Sudmoon, R. and Tanee, T. 2007. A New Species of *Zingiber* (Zingiberaceae) from Northern Thailand. *Taiwania* 52(2): 159–163.
- Dalzell, N. A. 1852. Contributions to the Botany of Western India. *Hooker's Journal of Botany and Kew Garden Miscellany* 4: 341–347.
- Gagnepain, F. 1908. Zingibéracées. In: Lecomte, H. (ed.) *Flore générale de l'Indo-Chine* 6, pp 25–121. Masson & Co., Paris.
- Kavitha, P. G., Nair, P., Nair, R. A., Jayachandran, B. K., Sabu, M. and Thomas, G. 2007. AFLP polymorphism and Phytiun response in *Zingiber* species. In: Kes-havachandran, R., Nazeem, P., Girija, D., John, P. S. and Peter, K. V. (eds.), *Recent trends in horticultural biotechnology* 2: 497–503. ICAR national symposium on biotechnological interventions for improvement of horticultural crops: issues and strategies, Vellanikkara, Kerala, India, 10–12 January. New India Publishing, New Dehli, India.
- Kishor, R. and Leong-Škorničková, J. 2013. *Zingiber kangleipakense* (Zingiberaceae): a new species from Manipur, India. *Gardens' Bulletin Singapore* 65: 39–46.
- Kress, W. J., DeFilipps, R. A., Farr, E. and Kyi, D. Y. Y. 2003. A checklist of the trees, shrubs, herbs, and climbers of Myanmar. *Contributions from the U.S. National Herbarium* 45: 1–590.
- Larsen, K. and Larsen, S. S. 2006. *Gingers of Thailand*, 184 pp. Queen Sirikit Botanic Garden, Chiang Mai.
- Leong-Škorničková, J., Binh, N. Q., Dang, T. H., Sida, O., Rybkova, R. and Vuong, T. V. 2015. Nine new *Zingiber* species (Zingiberaceae) from Vietnam. *Phytotaxa* 219 (3): 201–220.
- Rahman, M. A. and Yusuf, M. 2013. *Zingiber salarkhanii* (Zingiberaceae), A New Species from Bangladesh. *Bangladesh Journal of Plant Taxonomy* 20(2): 239–242.
- Roxburgh, W. 1810. Description of Monandrous Plants. *Asiatic Research* 11: 348. Calcutta.
- Tanaka, N. 2012a. Studies of Zingiberaceae in Myanmar I: Newly Recorded Species of *Zingiber*, *Zingiber idae*. *Makinoa New Series* 10: 153–157.
- Tanaka, N. 2012b. Studies of Zingiberaceae in Myanmar II: A New Species, *Zingiber popaense* from Mt. Popa. *Makinoa New Series* 10: 159–163.
- Theilade, T. 1999. A synopsis in the genus *Zingiber* (Zingiberaceae) in Thailand. *Nordic Journal of Botany* 19: 389–410.
- Thongam, B and Kansom, B. 2014. A new species of *Zingiber* (Zingiberaceae) from Nagaland, India. *Phytotaxa* 178(3): 221–224.
- Triboun, P., Larsen, K., and Chantaranothai, P. 2014. A key to the genus *Zingiber* (Zingiberaceae) in Thailand with descriptions of 10 new taxa. *Thai Forest Bulletin, Botany* 6(1): 53–77.
- Tripathi, S. and Singh, K. K. 2006. Taxonomic Revision of the genus *Zingiber* Boehm. in North-East India. *Journal of Economic and Taxonomic Botany* 30: 520–530.
- Win, T. T. 1975. *Zingiberaceae of Burma in part*. Department of Botany, pp. 149. Arts and Science University, Rangoon.
- Wu, T. L. and Larsen, K. 2000. Zingiberaceae. In: Wu, Z.-Y. and Raven, P. H. (eds.), *Flora of China* vol. 24, pp. 333–346. Science Press, Beijing and Missouri Botanical Garden Press, St. Louis.