

Chrysomelid Beetles of Nepal Collected by the Himalaya Expedition 1979 of the National Science Museum, Tokyo (Part 2)¹⁾

Shinsaku Kimoto²⁾ and Haruo Takizawa³⁾

²⁾ 7–97, Mikunigaoka, Ogori City, Fukuoka, 838–0103 Japan

³⁾ Kami 2–7–16, Hasuda City, Saitama, 349–0122 Japan

Abstract This is the result of our study on the chrysomelid beetles of Nepal, which were collected by the National Science Museum Expedition to East Nepal in 1979. A small number of material collected by the 1981 survey is also dealt with. All the specimens examined including the holotypes to be designated are deposited in the collection of the National Science Museum, Tokyo, except for some duplicates retained at the authors' hands.

Key words: Chrysomelidae, Nepal, Taxonomy

In this paper, materials are classified into 45 species, one of which is described as new species and 6 species are recorded for the first time from Nepal.

Subfamily Chrysomelinae

Phaedon kimotoi Daccordi, 1977

Specimen examined: Tingsan La, 3,320 m, Sindhu Distr., 1 ex., 12–13. xi. 1979, M. Satô.

Distribution: Nepal.

Subfamily Galerucinae

Mandarella birmanica Bryant, 1954

Specimen examined: Godawari, 1,500 m, 1 ex., 28. ix. 1979, M. Satô.

Remarks: In the type specimen, the dorsal surface is entirely blackish and the legs are entirely brownish. Our specimen is slightly different from the type in having the dorsal surface entirely bluish.

Distribution: Nepal, Burma.

Monolepta signata (Olivier, 1808)

Specimens examined: Dobhan, 780 m, Terhathum Distr., 2 exs., 27. x. 1979, M. Satô.

Distribution: Nepal, India, Ceylon, Burma, Bhutan, Thailand, Cambodia, Vietnam, S. China, Hainan, Java.

Subfamily Alticinae

Amphimeloides sexmaculatus

Kimoto et Takizawa, sp. nov.

(Fig. 1)

Description: Ovate, convex; generally yellowish brown; pronotum with a pair of blackish transversely subtriangular markings basally; elytron with three large blackish markings, viz. basi-interior, lateral and subapical markings; head with a blackish marking on vertex; ventral surface with meso- and metathoraces and middle of first to third abdominal segments pitchy black; antenna pitchy black with four basal segments yellowish brown.

Head with vertex smooth and impunctate; frontal tubercle not distinctly raised. Antenna not close together, but inserted close to eye, and thickened towards apex; nearly a half as long as length of body; first segment elongate, club-

¹⁾ This study is supported by the Grants-in-aid for Overseas Scientific Survey Nos. 404101 and 57043055 from the Ministry of Education, Science and Culture, Japan. (Part 1): *Bull. Natn. Sci. Mus., Tokyo, ser. A*, 9(2): 83–96 (1983).




Fig. 1. a, *Amphimeloides sexmaculatus* Kimoto et Takizawa, sp. nov.; b, *A. bistrispunctatus* Chen.

shaped; second slightly shorter and thicker than third; fourth and fifth subequal in length; from sixth to eleventh gradually thickened; eleventh pointed at apex. Prothorax much broader than long; anterior margin emarginate, basal margin widely rounded, sides rounded; surface convex, smooth and distinctly impressed by fine punctures. Elytra broader than prothorax, surface smooth, shining, confusedly and strongly punctate.

Length 2.2–2.5 mm.

Holotype: Taplejung, 1,790 m, Taplejung Distr., Nepal, 27. x. 1979, M. Tomokuni. *Paratype*: 1 ex., same data as the holotype.

Remarks: This new species somewhat resembles *Amphimeloides bistrispunctatus* Chen, 1933, from Japan, in having the elytron with some distinct blackish markings but differs in having the antenna generally pitchy black with four basal segments yellowish brown.

***Amydus castaneus* Chen, 1935**

Specimens examined: Chitre, 2,500 m, Terhathum Distr., 5 exs., 1. xi. 1979, M. Tomokuni.

Distribution: Nepal.

***Amydus* sp. 1.**

Specimen examined: Minchhindhap Danda, 2,300–2,950 m, Terhathum Distr., 1 ex., 28. x. 1979, M. Tomokuni.

***Amydus* sp. 2.**

Specimen examined: Tinjura Danda, 2,600–3,000 m, Terhathum Distr., 1 ex., 31. x. 1979, M. Tomokuni.

***Aphthona gardneri* Bryant, 1941**

Specimens examined: Minchhindhap Danda, 2,300–2,950 m, Terhathum Distr., 4 exs., 28. x. 1979, M. Tomokuni. Khari La, 3,081 m—Kharikhora, 2,050 m, Solukhumbu Distr., 1 ex., 7. x. 1979, M. Tomokuni. Junbesi, 2,670 m—Lamjura La, 3,530 m, Solukhumbu Distr., 1 ex., 12. x. 1979, M. Tomokuni. Godawari, 1,500 m, 2 ex s., 28. ix. 1979, M. Satô. Surke, 2,290 m, Solukhumbu Distr., 2 exs., 5. x. 1979, M. Satô. Kemche Danda, 2,400–3,080 m, Solukhumbu Distr., 1 ex., 9. x. 1979, M. Satô. Kurdi, 3,000 m—Chumrung, 2,000 m, nr. Ghandrung, 2 exs., 21. x. 1981, M. Sakai. Landrung, 1,600 m—Damphus, 1,350 m, nr. Pokhara, 1 ex., 23. x. 1981, M. Sakai.

Distribution: Nepal, India.

Aphthona kanaraensis Jacoby, 1896

Specimens examined: Bhalko, 2,450 m, Dhankuta Distr., 1 ex., 1. xi. 1979, M. Tomokuni. Palati, 1,200 m, Sindhu Distr., 3 exs., 9. xi. 1979, M. Satô.

Distribution: Nepal.

Aphthona opaca Allard, 1889

Annls. Soc. ent. France, (6), 9: 305 (Cambodia).

Specimen examined: Khari La, 3,081 m — Kharikhola, 2,050 m, Solukhumbu Distr., 1 ex., 7. x. 1979, M. Tomokuni.

Distribution: Nepal, Thailand, Cambodia, Vietnam, SW China.

Remarks: This species is here recorded for the first time from Nepal.

Batophila castanea Scherer, 1989

Specimens examined: Gufa, 2,950–3,000 m, Terhathum Distr., 1 ex., 29. x. 1979, M. Tomokuni, 1 ex., 28–29. x. 1979, M. Satô. Tinjura Danda, 2,600–3,000 m, Terhathum Distr., 1 ex., 31. x. 1979, M. Tomokuni.

Distribution: Nepal, India.

Chaetocnema (Chaetocnema) nepalensis

Scherer, 1969

Specimens examined: Hile, 2,070 m, Dhankuta Distr., 5 exs., 2. xi. 1979, M. Satô. Palati, 1,200 m, Sindhu Distr., 1 ex., 9. xi. 1979, M. Satô. Dhankuta, 1,100 m, Dhankuta Distr., 1 ex., 2. xi. 1979, light trap, M. Tomokuni. Drumthali, 2,420 m, Sindhu Distr., 1 ex., 13. xi. 1979, light trap, M. Tomokuni. Kabre, 1,760 m, Dolokha, Janakpur, 1 ex., 17. x. 1979, S. Uéno.

Distribution: Nepal.

Chaetocnema (Chaetocnema) singala

Maulik, 1926

Specimens examined: Kenja, 1,720 m, Solukhumbu Distr., 1 ex., 13. x. 1979, M. Tomokuni. Drumthali, 2,420 m, Sindhu Distr., 1

ex., 13. xi. 1979, light trap, M. Tomokuni. Shera, 1,420 m, Dolakha Distr., 2 exs., 19. x. 1970, light trap, M. Tomokuni. Changma, 2,200 m, Ramechhap Distr., 1 ex., 14. x. 1979, M. Tomokuni. Dolangsa Khola, 2,340 m, Sindhu Distr., 1 ex., 13. xi. 1979, M. Satô. Hile, 2,070 m, Dhankuta Distr., 5 exs., 2. xi. 1979, M. Satô.

Distribution: Nepal, India, Ceylon, Thailand, Cambodia.

Chaetocnema (Tlanoma) sp.

(nr. *indica* Weise)

Specimen examined: Palati, 1,200 m, Sindhu Distr., 1 ex., 9. xi. 1979, M. Satô.

Hemipyxis fulvipennis (Illiger, 1807)

Specimen examined: Dharan Bazar, 350–370 m, Sunsari Distr., 1 ex., 6. xi. 1979, M. Tomokuni.

Distribution: Nepal, India, Burma, Vietnam, S. China.

Hemipyxis patkaia (Maulik, 1926)

Specimens examined: Dharan Bazar, 350–370 m, Sunsari Distr., 29 exs., 6. xi. 1979, M. Tomokuni. Sardhu Khola, 350 m, Sunsari Distr., 9 exs., 6. xi. 1979, M. Satô.

Distribution: Nepal, India.

Hespera cyanea Maulik, 1926

Specimens examined: Minchhindhap Danda, 2,300–2,950 m, Terhathum Distr., 2 exs., 28. x. 1979, M. Tomokuni. Kenja, 1,620 m, Solukhumbu Distr. — Changma, 2,200 m, Ramechhap Distr., 2 exs., 13. x. 1979, M. Tomokuni. Phakding, 2,580 m — Nangbug, 2,550 m, Solukhumbu Distr., 1 ex., 5. x. 1979, M. Tomokuni. Surke, 2,290 m, Solukhumbu Distr., 1 ex., 6. x. 1979, M. Satô.

Distribution: Nepal, India, Burma, Vietnam, W. China.

Hespera fulvimenbris Kung et Chen, 1954

Act. ent. sin., 4: 154 (N. Vietnam).

Specimens examined: Lukla, 2830 m—Phakding, 2,580 m, Solukhumbu Distr., 5 exs., 29. ix. 1979, M. Tomokuni. Phakding, 2,580 m—Namche Bazar, 3,440 m, Solukhumbu Distr., 1 ex., 30. ix. 1979, M. Tomokuni. Kathmandu, 1 ex., 15. x. 1979, M. Satô.

Distribution: Nepal, Vietnam.

Remarks: This species is here recorded for the first time from Nepal.

Hespera krishna Maulik, 1926

Specimens examined: Drumthali, 2,100–2,420 m, Sindhu Distr., 18 exs., 14. xi. 1979, M. Tomokuni. Phakding, 2,580 m—Namche Bazar, 3,440 m, Solukhumbu Distr., 2 exs., 30. ix. 1979, M. Tomokuni. Patashe Danda Gonda, 2,200–2,400 m, Dolakha, Tanakpur, 3 exs., 15. x. 1979, Y. Nishikawa. Godavari, 1,500 m, 1 ex., 21. ix. 1979, M. Satô.

Distribution: Nepal, India, Burma, Bhutan.

Hespera kumaonensis Scherer, 1969

Specimens examined: Silvalaya, 1,800 m, Ramechhap Distr., 14 exs., 15. x. 1979, M. Tomokuni. Phakding, 2,580 m, Solukhumbu Distr., 3 exs., 30. ix. 1979, M. Tomokuni.

Distribution: Nepal, India.

Hespera lomasa Maulik, 1926

Fn. Brit. India, Chrys. & Halt.: 142 (S. India, Ceylon).

Specimens examined: Dharapani, 1,000 m, Dhankuata Distr., 12 exs., 4. xi. 1979, M. Tomokuni.

Distribution: Nepal, India, Ceylon, Burma, Thailand, Laos, Vietnam, China, Taiwan, Ryukyu Is.

Remarks: This species is here recorded for the first time from Nepal.

Hespera naini Scherer, 1969

Pacif. Ins. Mon., 22: 45 (India).

Specimens examined: Phakding, 2,580 m—Nangbug, 2,550 m, Solukhumbu Distr., 2 exs., 5.

x. 1979, M. Tomokuni. Phakding, 2,580 m—Namche Bazar, 3,440 m, Solukhumbu Distr., 6 exs., 30. ix. 1979, M. Tomokuni. Namche Bazar, 3,440 m—Monjo, 2,880 m, Solukhumbu Distr., 1 ex., 4. x. 1979, M. Tomokuni. Jorsale, 2,800 m, Solukhumbu Distr., 3 exs., 4. x. 1979, M. Satô. Surke, 2,290 m, Solukhumbu Distr., 1 ex., 6. x. 1979, M. Satô. Chaunrikharka, 2,670 m, Solukhumbu Distr., 1 ex., 5. x. 1979, M. Satô. Monjo, 2,800 m, Solukhumbu Distr., 1 ex., 4. x. 1979, Y. Nishikawa.

Distribution: Nepal, India.

Remarks: This species is here recorded for the first time from Nepal.

Hespera sericea Weise, 1889

Specimens examined: Dharapani, 1,000 m, Dhankuta Distr., 7 exs., 4. xi. 1979, M. Tomokuni. Shivalaya, 1,800 m, Ramechhap Distr., 1 ex., 15. x. 1979, M. Tomokuni.

Distribution: Nepal, India, Thailand, Vietnam, S. China.

Hespera strigiceps Kimoto, 2001

Specimen examined: Mahidingma, 2,240 m, Solukhumbu Sagarmatha., 1 ex., 8. x. 1979, Y. Nishikawa.

Distribution: Nepal.

Hespera violaceipennis Scherer, 1969

Specimen examined: Pekarnasa, 1,480 m, Ramechhap, Janakpur, 1 ex., 18. x. 1979, Y. Nishikawa.

Distribution: Nepal, India.

Hyphasis indica (Baly, 1879)

Specimen examined: Birthani—Ulleri, 1 ex., 30. v. 1969, T. Miyashita. Ulleri, 1 ex., 30. v. 1969, T. Miyashita.

Distribution: Nepal, India.

Longitarsus birmanicus Jacoby, 1892

Specimens examined: Dhankuta, 1,100 m,

Dhankuta Distr., 5 exs., 2. xi. 1979, S. Uéno, 2 exs., 2. xi. 1979, light trap, M. Tomokuni. Dhankuta, 1,200 m, Kusi Zone, Dhankuta Distr., 1 ex., 2. xi. 1979, M. Satô. Hile, 2,070 m, Dhankuta Distr., 5 exs., 2. xi. 1979, M. Satô. Dharapani, 1,000 m, Dhankuta Distr., 5 exs., 4. xi. 1979, M. Tomokuni. Minchhindhap Danda, 2,300–2,950 m, Terhathum Distr., 3 exs., 28. x. 1979, M. Tomokuni. Palati, 1,200 m, Sindhu Distr., 1 ex., 9. xi. 1979, light trap, M. Tomokuni, 1 ex., 9. xi. 1979, M. Satô. Shivalaya, 1,800 m, Ramechhap Distr., 2 exs., 15. x. 1979, M. Tomokuni. Drumthali, 2,420 m, Sindhu Distr., 1 ex., 13. xi. 1979, M. Tomokuni. Drumthali, 2,100–2,420 m, Sindhu Distr., 1 ex., 14. xi. 1979, M. Tomokuni. Kharte Danda, 2,700–3,081 m, Solukhumbu Distr., 2 exs., 7. x. 1979, M. Satô. Poyan Khola, 2,730 m, Solukhumbu Distr., 1 ex., 7. x. 1979, M. Satô. Gonda, 2,200–2,300 m, Dolakha Distr., 1 ex., 15. x. 1979, M. Tomokuni. Bhalko, 2,450 m, Dhankuta Distr., 1 ex., 1. ex., 1. xi. 1979, M. Tomokuni. Sardhu Khola, 490–580 m, Dhankuta Distr., 3 exs., 5. xi. 1979, M. Tomokuni. Chauki, 2,750 m, Terhathum Distr., 1 ex., 30. x. 1979, Y. Nishikawa.

Distribution: Nepal, India, Ceylon, Burma, Thailand, Vietnam, Java, Bali.

***Longitarsus cheni* Scherer, 1969**

Pacif. Ins. Mon., 22: 62 (India).

Specimen examined: Kemche Danda, 2,400–3,080 m, Solukhumbu Distr., 1 ex., 9. x. 1979, M. Satô.

Distribution: Nepal, India.

Remarks: This species is here recorded for the first time from Nepal.

***Lomgotarsus championi* Maulik, 1926**

Specimen examined: Dolangsa, 2,050–2,600 m, Sindhu Distr., 1 ex., 11. xi. 1979, M. Tomokuni.

Distribution: Nepal, India.

***Longitarsus cyanipennis* Bryant, 1924**

Specimens examined: Namche Bazar, 3,440 m—Monjo, 2,800 m, Solukhumbu Distr., 1 ex., 4. x. 1979, M. Tomokuni. Godavari, 1,700 m, 1 ex., 16. x. 1979, M. Satô.

Distribution: Nepal, India, Thailand, Vietnam, China, Taiwan.

***Longitarsus gressitti* Scherer, 1969**

Specimens examined: Pakding, 2,580 m—Namche Bazar, 3,440 m, Solukhumbu Distr., 1 ex., 30. ix. 1979, M. Tomokuni. Khari La, 3,081 m—Kharikhola, 2,050 m, Solukhumbu Distr., 1 ex., 5. xi. 1979, M. Tomokuni. Gorjagong, 2,080 m, Terhathum Distr., 1 ex., 28. x. 1979, M. Satô. Kemche Danda, 2,400–3,080 m, Solukhumbu Distr., 1 ex., 9. x. 1979, M. Satô. Palati, 1,200 m, Sindhu Distr., 2 exs., 9. xi. 1979, M. Satô.

Distribution: Nepal.

***Longitarsus hsiensis* Chen, 1939**

Specimens examined: Phakding, 2,580 m—Namche Bazar, 3,440 m, Solukhumbu Distr., 2 exs., 30. ix. 1979, M. Tomokuni. Lukia, 2,830 m—Phakding, 2,580 m, Solukhumbu Distr., 1 ex., 29. ix. 1979, M. Tomokuni.

Distribution: Nepal, India, Vietnam, China.

***Manobia merkii* Medvedev, 1998**

Specimens examined: Dharapani, 1,000 m, Dhankuta Distr., 2 exs., 4. xi. 1979, M. Tomokuni. Jorpati, 2,200–2,300 m, Dhankuta Distr., 1 ex., 1. xi. 1979, M. Tomokuni.

Distribution: Nepal, India.

***Manobia violaceipennis* Medvedev, 1992**

Specimens examined: Bhalko, 2,450 m, Dhankuta Distr., 3 exs., 1. xi. 1979, M. Tomokuni, 1 ex., 1. xi. 1979, M. Satô. Tinjura Danda, 2,600–3,000 m, Terhathum Distr., 1 ex., 31. x. 1979, M. Tomokuni. Shivalaya, 1,800 m, Ramechhap Distr., 1 ex., 15. x. 1979, M. Tomokuni. Chitre, 2,500 m, Terhathum Distr., 2 exs., 1. xi.

1979, M. Tomokuni. Dharapani, 1,000 m, Dhankuta Distr., 1 ex., 4. xi. 1979, M. Tomokuni. Phakding, 2,580 m—Nangbug, 2,550 m, Solukhumbu Distr., 4 exs., 5. x. 1979, M. Tomokuni.

Distribution: Nepal.

***Manobia* sp. 1**

Specimen examined: Gufa, 2,950–3,000 m, Terhathum Distr., 1 ex., 29. x. 1979, M. Tomokuni.

***Manobia* sp. 2**

Specimen examined: Nangbung, 2,600 m, Solukhumbu Distr., 1 ex., 5. x. 1979, M. Satô.

***Maulika bengalensis* Lopatin, 1984**

Specimens examined: Lamjura Danda, E. side, 3,220 m to W. side, 3,300 m, Sagarmatha Distr., 1 ex., 12. x. 1979, M. Owada. Kemche Danda, Tragsindo La, 2,930–3,070 m, Solukhumbu Distr., 1 ex., 9. x. 1979, Y. Nishikawa.

Distribution: Nepal.

***Manobia* sp. (nr. *simplicithorax* Chen)**

Specimen examined: Hile, 2,070 m, Dhankuta Distr., 1 ex., 2. xi. 1979, M. Satô.

Remarks: This specimen closely resembles *M. simplicithorax* Chen, but differs in having the punctures of the elytron stronger.

***Nepalicropis dajeelingensis* Scherer, 1969**

Specimens examined: Chitre, 2,500 m, Terhathum Distr., 2 exs., 1. xi. 1979, M. Tomokuni.

Distribution: Nepal, India.

***Nonarthra variabilis* Baly, 1862**

Specimens examined: Lukia, 2,830 m—Phakding, 2,580 m, Solukhumbu Distr., 1 ex., 29. ix. 1979, M. Tomokuni. Phadking, 2,580 m—Namche Bazar, 3,440 m, Solukhumbu Distr., 1 ex., 30. ix. 1979, M. Tomokuni. Yarsa, 1,960 m,

Dolakha Distr., 2 exs., 17. x. 1979, M. Owada.

Distribution: Nepal, Afghanistan, India, Ceylon, Burma, Thailand, Laos, Vietnam, China, Hainan, Taiwan, Ryukyu Is., Malaya, Philippines.

***Phyllotreta chotanica* Duvivier, 1892**

Specimen examined: Shivalaya, 1,800 m, Ramechhap Distr., 1 ex., 15. x. 1979, M. Tomokuni.

Distribution: Nepal, India.

***Psylliodes viridana* Motschulsky, 1858**

Specimens examined: Gonda, 2,200 m–2,300 m, Dolakha Distr., 1 ex., 15. x. 1979, M. Tomokuni. Dolangsa, 2,050–2,600 m, Sindhu Distr., 1 ex., 11. xi. 1979, M. Tomokuni.

Distribution: Nepal, India, Ceylon.

***Sphaeroderma luteipenne* Weise, 1922**

Tijdschr. Ent., 65: 118 (Tonkin).

Specimens examined: Ringmo, 2,780 m—Khamje, 2,800 m, Solukhumbu Distr., 1 ex., 11. x. 1979, M. Tomokuni. Phakding, 2,580 m—Nangbug, 2,550 m, Solukhumbu Distr., 1 ex., 5. x. 1979, M. Tomokuni. Lukia, 2,830—Phakding, 2,580 m, Solukhumbu Distr., 1 ex., 29. ix. 1979, M. Tomokuni.

Distribution: Nepal, Thailand, Vietnam.

Remarks: This species is here recorded for the first time from Nepal.

***Taizonia andreevi* Gruev, 1985**

Specimen examined: Sangridara, 1,250 m, Dhankuta Distr., 1 ex., 4. xi. 1979, Y. Nishikawa.

Distribution: Nepal.

***Taizonia minima* (Scherer, 1969)**

Specimens examined: Dolangsa, 2,700 m, Sindhu Distr., Bagmati, 2 exs., 11. xi. 1979, Y. Nishikawa.

Distribution: Nepal, India.

References

- Allard, M., 1889. Contributions à la faune Indo-Chinoise, 5^e Memoire. Galerucides et Alticides. *Annls. Soc. ent. France*, **9**(6): 303–312.
- Kung, Y. T., & S. H. Chen, 1954. Notes on the Chinese and Tonkinese species of Halticinae genus *Hespera* Weise. *Acta ent. sin.*, **4**: 149–163.
- Maulik, S., 1926. Coleoptera, Chrysomelidae (Chrysomelinae and Halticinae). Fauna of British India, including Ceylon and Burma. Taylor and Francis, London. 442 pp.
- Medvedev, L. N. & Eva Sprecher-Uebersax, 1999. Katalog der Chrysomelidae von Nepal. *Ent. basil.*, **21**: 261–354.
- Scherer, G., 1969. Die Alticinae des indischen Subkontinentes (Coleoptera, Chrysomelidae, Alticinae). *Pacif. Ins. Mon.*, **22**: 1–251.
- Weise, J., 1922. Chrysomelidae der Indo-Malayischen Region. *Tijdschr. Ent.*, **65**: 39–130.

