

How striations of ophiuroid and asteroid trace fossils were produced—Observations of tube-feet movement in living ophiuroids and asteroids

YOSHIAKI ISHIDA¹, TOSHIHIKO FUJITA², MASATO KIYOMOTO³, MARTIN RÖPER⁴,
TOSHIFUMI KOMATSU⁵, KOKI KATO⁶, KOTARO KAMADA⁷, YASUNARI SHIGETA² AND TARO KUMAGAE⁸

¹2-20-13 Kamiogi, Suginami-ku, Tokyo 167-0043, Japan (e-mail: y-ishida@msi.biglobe.ne.jp)

²National Museum of Nature and Science, 4-1-1 Amakubo, Tsukuba, Ibaraki 305-0005, Japan

³Ochanomizu University, 11 Koyatsu, Tateyama, Chiba 294-0301, Japan

⁴Bürgermeister-Müller-Museum, 8 Bahnhofstrasse, Solnhofen 91807, Germany

⁵Kumamoto University, 2-39-1 Kurokami, Chuo-ku, Kumamoto 860-8555, Japan

⁶Graduate School of Science and Technology, Kumamoto University, 2-39-1 Kurokami, Chuo-ku, Kumamoto 860-8555, Japan

⁷Hirosaki University, 1 Bunkyo-cho, Hirosaki, Aomori 036-8560, Japan

⁸Japan Petroleum Exploration Corporation, 1-7-12 Marunouchi, Chiyoda-ku, Tokyo 100-0005, Japan

Received February 21, 2015; Revised manuscript accepted March 16, 2016

Abstract. To clarify the process of producing the striations in the trace fossil *Asteriacites*, we observed the behavior of living ophiuroids and asteroids in aquariums. When ophiuroids stopped crawling, they buried themselves shallowly in the substratum, removing substratum under their arms and discs by using their tube-feet. The basal-arm tube-feet produce well spaced, fine, parallel striations that are perpendicular to the arm axis. The oral tube-feet produce fine, radial striations in the central depression. When the ophiuroids resumed crawling, they raised their disc and four arms above the substratum and dragged one arm backward. The one backward arm erased the striations, and parallel fine striations remained in four arm depressions. Similarly, asteroids also produced wide and shallow striations perpendicular to the arm axis by tube-feet movement. When the asteroids started to move again, they bulldozed the substratum under the one preceding arm, where the striations were erased. Since the asteroids crawled raising the other four arms by their tube feet which produced rough and deep striations, the wide and shallow striations remained only in a half of each arm depression. The striations of *Asteriacites lumbricalis* and *A. quinquefolius* were similar in shape to the striations produced by movement of tube feet of living ophiuroids and asteroids, respectively.

Key words: *Asteriacites*, brittle star, sea star, striation, trace fossil, tube-foot

Introduction

Ichnospecies *Asteriacites lumbricalis* Schlotheim, 1820 and *A. quinquefolius* (Quenstedt, 1876) are well known star-shaped trace fossils. In previous studies, these two ichnospecies have been distinguished by the following features. *Asteriacites lumbricalis* has a distinct central subcircular area with narrow arms and faint striations in the arms (Seilacher, 1953; Hakes, 1976; Twitchett and Wignall, 1996; Ishida *et al.*, 2004), and *A. quinquefolius* has wide arms from an undifferentiated center and dense striations in the arms (Seilacher, 1953; Patel *et al.*, 2008; Ishida *et al.*, 2013).

The producers and their producing processes have been

studied based mainly on the shape of traces (e.g. Seilacher, 1953; Mángano *et al.*, 1999; Ishida *et al.*, 2004, 2013). However, the detailed morphology of the traces including striations (bioglyphs; Seilacher, 2007) and how the traces are produced have not been clarified well enough. For the paleontological implications of trace fossils, to specify the producing organisms and to clarify the producing processes are the most important, and observations of morphology and behavior of living organisms are effective methods to interpret trace fossils. *Asteriacites lumbricalis* was interpreted as a resting trace of ophiuroids (Seilacher, 1953; Chamberlain, 1971; Mikuláš, 1990; West and Ward, 1990; Mángano *et al.*, 1999; Wilson and Rigby, 2000; Ishida *et al.*, 2004, 2013), while *A. quinquefolius*




Figure 1. Sampling localities in Japan and Russia (A) and Germany (B). Star-shaped symbols indicate sampling localities, and letters near the symbols denote the site number.

was interpreted as a resting trace of asteroids (Seilacher, 1953; Patel *et al.*, 2008; Ishida *et al.*, 2013).

Both named fossils have characteristic striations on their arms. Seilacher (1953, 2007) observed the traces produced by living ophiuroids and asteroids in an aquarium and suggested the striations were marked by tube-feet, but morphological details of the striations and the behavior of the ophiuroids and asteroids producing them have never been studied. In the present study, we observed the tube-feet movement and the striations produced by living ophiuroids and asteroids in an aquarium as well as *in situ*. We compared their striations with those of *Asteriacites lumbricalis* and *A. quinquefolius* from four formations, and the producers of these trace fossils were precisely specified. In addition, the morphologies of striations of previously reported *Asteriacites* fossils were reconsidered based on our observations.

Material and methods

Two extant ophiuroid species *Ophiura kinbergi* (Ljungman, 1866) and *Stegophiura sterea* (H. L. Clark, 1908) and one extant asteroid species *Astropecten scoparius* Müller and Troschel, 1842 were used for the laboratory observations. For *O. kinbergi*, six specimens (7.9–8.9 mm in disc diameter) were recovered at a depth of 80 m in the Sea of Japan (Site A in Figure 1A) during the research cruise of T. S. *Toyoshio-maru* of Hiroshima University in July 2001 and four specimens (2.5–3.9 mm

in disc diameter) were recovered at a depth of 6–15 m in May 2012 in Tateyama Bay, Japan (Site D in Figure 1A), by the research boat *Sea-star* of Ochanomizu University. One specimen (6.6 mm in disc diameter) of *S. sterea* was recovered at a depth of 200 m in the Sea of Japan (Site B in Figure 1A) during the research cruise of R. V. *Mizuho-maru* of the Japan Sea National Fisheries Research Institute in May 2002. For *A. scoparius*, one specimen (72 mm in arm length) was recovered at a depth of 45–49 m in Manazuru Bay, Japan (Site C in Figure 1A) during the research cruise of R. V. *Ryokuyo-maru* of the Field Science Education and Research Center, Kyoto University in March 2012, and one specimen (115 mm in arm length) was recovered from a depth of 6–15 m in May 2012 in Tateyama Bay (Site D in Figure 1A), by the research boat *Sea-star* of Ochanomizu University. Additionally, one specimen (32 mm in arm length) was collected by hand from the intertidal zone in Aburatsubo Bay, Japan (Site E in Figure 1A) in August 2012.

The animals were kept alive in laboratories before observation. Behavior of one individual of *Stegophiura sterea*, eight individuals of *Ophiura kinbergi* and two individuals of *Astropecten scoparius* in a glass beaker (17 cm in diameter and 9.5 cm in height) or that of two individuals of *O. kinbergi* and one individual of *A. scoparius* in a plastic tray (30 × 22 × 5 cm in size) was observed at room temperature (12–25°C). Two different kinds of substratum were filled in to a thickness of 10 mm: beach sand (0.2–0.5 mm in grain size equivalent to fine to medium




Figure 2. Tube-feet movement of extant ophiuroids, *Ophiura kinbergi* (A, B) and *Stegophiura sterea* (C, D), when burrowing into beach sand observed in the glass beaker. Video images and sketches of them are shown in B and D. **A**, resting posture, upside view, specimen from Site A. **B**, successive underside images when burrowing (time interval 0.3 sec), specimen from Site D. A basal-arm tube-foot is extended and retracted. **C**, resting posture, upside view, specimen from Site B. **D**, underside image when burrowing, specimen from Site B. Most of basal-arm tube-feet and oral tube-feet are extended. Abbreviations: Ar, arm; Mo, mound of substratum; Ot1, first oral tube-foot; Ot2, second oral tube-foot; Tu, tube-foot. Scale bars equal 5 mm (A, C) and 1 mm (B, D).

sand) in a glass beaker and aluminiferous abrasive (#1500 in grain size equivalent to mud) in a plastic tray. Tube-feet movement of the ophiuroids and asteroid was recorded by a video camera from the under and lateral sides of the glass beaker. Their traces that remained on the surface of the substratum were photographed from above the water surface. Sizes of oral and basal-arm tube-feet of one individual of *O. kinbergi*, one individual of *S. sterea* and of tube-feet of one individual of *A. scoparius* were measured on the video by a caliper square.

Star-shaped traces on the sea bottom produced by the living ophiuroid and asteroid were reported on photographs of the deep-sea floor in Japan (Ishida *et al.*, 2004, 2013). The trace of the ophiuroid *Amphiophiura penichra* (H. L. Clark, 1911) was taken on the silt bottom at 730 m deep off northern Japan in July 1984 (Site F in Figure 1A) and that of an astropectinid asteroid on the silt bottom at 861–923 m deep off central Japan in May 1986 (Site G in Figure 1A), by a deep-sea camera during cruises of R. V. *Tansei-maru* of the Ocean Research Institute, University of Tokyo. Striated traces were observed in the star-shaped traces in the underwater photographs.

Three *Asteriacites* fossils already reported in previous

papers from two formations were reexamined, in particular, with respect to their striations: a concave epirelief of *Asteriacites lumbricalis* (diameter of central depression 13.2 mm, depth of central depression 0.5 mm, arm length 22.1 mm) in the mudstone of the Olenekian (Lower Triassic) Hiraiso Formation in Miyagi Prefecture, Japan (Site H in Figure 1A) (Kamada and Takizawa, 1992; Ishida *et al.*, 2004); a concave epirelief of *A. lumbricalis* (diameter of central depression 7.7 mm, depth of central depression 0.3 mm, arm length 23.4 mm) in the fine-grained limestone of the lower Tithonian (Upper Jurassic) Hienheim Formation at Ried, Germany (Site J in Figure 1B) (Röper and Rothgaenger, 1995, 1998; Ishida *et al.*, 2013); a concave epirelief of *A. quinquefolius* (depth of central depression 5 mm, arm length 53 mm) in the fine-grained limestone of the lower Tithonian (Upper Jurassic) Hienheim Formation at Ried, Germany (Site J in Figure 1B) (Röper and Rothgaenger, 1995, 1998; Ishida *et al.*, 2013). Additionally, two unidentified star-shaped trace fossils from two formations were also examined: a convex hyporelief (diameter of central ridge 7.1 mm, height of central ridge 0.5 mm, arm length 16.9 mm) in the medium sandstone of the Spathian (Lower Triassic) Lazurnaya Bay

Table 1. Measurements of tube-feet and space between them of *Ophiura kinbergi* (disc diameter 3.9 mm), *Stegophiura sterea* (disc diameter 6.6 mm) and *Astropecten scoparius* (arm length 115 mm). Mean and range are shown for basal-arm tube-feet. Length and width were measured when extended during burying into the substratum.

Species	First oral tube-feet ($n = 1$)			Second oral tube-feet ($n = 1$)			Basal-arm tube-feet ($n = 4$)			
	Length (mm)	Width (mm)	Width/Length	Length (mm)	Width (mm)	Width/Length	Length (mm)	Width (mm)	Width/Length	Space (mm)
<i>Ophiura kinbergi</i>	0.90	0.19	0.22	1.05	0.21	0.20	0.63 (0.59–0.65)	0.13 (0.16–0.21)	0.21 (0.10–0.12)	0.46 (0.46–0.46)
<i>Stegophiura sterea</i>	1.98	0.19	0.10	2.16	0.25	0.12	1.69 (1.61–1.80)	0.18 (0.16–0.21)	0.11 (0.11–0.12)	0.61 (0.61–0.61)
<i>Astropecten scoparius</i>	–	–	–	–	–	–	13.9 (13.3–15.0)	4.3 (4.0–4.9)	0.32 (0.30–0.33)	0 (0–0)

Formation on Russky Island, South Primorye, Russia (Site I in Figure 1A) (Shigeta *et al.*, 2009); a concave epirelif (diameter of central depression 34.4 mm, depth of central depression 4.0 mm, arm length 39.0 mm) in the fine sandstone of the Upper Cretaceous Himenoura Group on Makishima Island in Kumamoto Prefecture, Japan (Site K in Figure 1A) (Komatsu *et al.*, 2008; Kojo *et al.*, 2011). The striations of the ichnofossils were observed using a binocular microscope or by tracing them onto transparent paper placed upon the fossil-bearing sediment.

The trace fossil specimens from the Hienheim Formation are housed at the Bürgermeister-Müller-Museum, Solnhofen, Germany (BMMS). The specimens from the Hiraiso Formation and Lazurnaya Bay Formation and the synthetic resin cast from the Himenoura Group are housed at the National Museum of Nature and Science, Tsukuba, Japan (NMNS PA 18300, 18301, 18302, respectively).

Results

Aquarium observations of extant species

In the aquarium, the ophiuroids *Ophiura kinbergi* and *Stegophiura sterea* usually crawled on the substratum or sat on it with their body shallowly buried. When they sat, *Ophiura kinbergi* buried itself more shallowly than *S. sterea* (Figure 2A, C). Both ophiuroids sat by burying the disc and basal arms into the sediment and thereby small mounds of the sediment were formed around the discs and along the basal arms (Figure 2A, C). They dug down into the substratum by mainly using basal-arm tube-feet and oral tube-feet (Figure 2B, D). Basal-arm tube-feet were slender and widely apart (Table 1). The ophiuroids repeated the motion of extending basal-arm tube-feet outward and retracting them inward to sweep away the substratum under the arms (Figure 2B). One extension and retraction of the basal-arm tube-feet took about 0.5

seconds for *O. kinbergi* and 2.0 seconds for *S. sterea*. On the other hand, longer first and second oral tube-feet in discs (Table 1) moved cyclically in a radial direction to sweep away the substratum under the disc (Figure 2D). One cycle of this movement of the first and second oral tube-feet took about 0.9 and 1.3 seconds for *O. kinbergi* and 2.2 and 3.3 seconds for *S. sterea*, respectively. The substratum swept away by basal-arm tube-feet and oral tube-feet formed mounds at the side of the body (Figures 2A, C, 3A).

When the ophiuroids started to emerge again from the substratum, they raised rapidly their disc and the basal parts of four arms above the substratum, and moved pushing the substratum with the distal parts of the four arms, dragging the remaining arm backward. Consequently, a disc and four-arm trace was formed (see also Ishida *et al.*, 2004 for production of resting traces of ophiuroids). The striations remaining in the resting trace were fine, spaced corresponding to the space between neighboring basal arm tube-feet, parallel to each other and perpendicular to the arm axis in the basal “arms.” Fine and radial striations remained in the central depression (Figure 3).

The asteroid *Astropecten scoparius* usually moved on the substratum or sat with its body shallowly buried in it. When the asteroid sat, it buried its arms into the sediment and small mounds of the sediment were formed along the arms. It dug down into the substratum by mainly using thick basal-arm tube-feet (Table 1). The tube-feet showed a cyclic movement, extending outward and retracting inward, to sweep away the substratum under the arms (Figure 4A). The tube-feet were very close or in contact with each other (Figure 4A, Table 1). One cycle of this movement was about 5 seconds. The sediment swept away by tube-feet formed mounds (Figure 5A). The asteroid moved with one preceding arm and four arms extended in a lateral direction (Figure 5A). So, under the four lateral arms, tube-feet moved perpendicularly




Figure 3. Producing process of striations on aluminiferous abrasive by *Ophiura kinbergi* (specimen from Site D), observed in the plastic tray. **A**, successive images when starting to move again from its resting posture (time interval 30 sec). One conspicuous trace marked by an oval was produced after the ophiuroid moved. **B**, sketch of striations of the conspicuous trace. Arrows indicate the direction which the ophiuroid moved. Abbreviations: Mo, mound of substratum; Stb, striation by basal-arm tube-feet; Sto, striation by oral tube-feet. Scale bar equals 1 mm.




Figure 4. Tube-feet movement of *Astropecten scoparius* when burrowing and starting to move again observed on beach sand in the glass beaker. Video images and sketches of them are shown in **A** and **B**. **A**, Successive underside images when burrowing (time interval 1 sec), specimen from Site D. Tube-feet were extended and retracted. **B**, lateral image when starting to move again, specimen from Site E. Tube-feet were stuck into the sediment. Arrows show the direction in which the asteroid moved. Abbreviations: Ims, infero-marginal spine; Su, substratum; Tu1–3, tube-foot, identical tube-feet are shown by the same numbers. Scale bars equal 1 cm.

to the arm axis (Figure 4A). When the asteroid started to shift again, it bulldozed the substratum under the one preceding arm, and the depression of the arm was completely destroyed. However, the other four depressions remained (see also Ishida *et al.*, 2013). The tube-feet were stuck deeply into the sediment in the front half of four arm depressions (Figure 4B), and the deep and irregular striations were redrawn there (Figure 5B). On the other hand, in their back half, wide and parallel striations contacting with each other, originally formed when digging, remained (Figure 5B).

In the basal part of the arm depressions of traces, the striations of both ophiuroids and asteroids were parallel

to each other and perpendicular to the arm axis, but the striation width was different between them. Moreover, ophiuroid traces had radial striations in the center of the depression (Figures 3, 5).

In situ observations of extant species

Striations in the star-shaped traces made by the ophiuroid *Amphiophiura penichra* were clearly visible in the four arm depressions, and they were well spaced, fine, parallel to each other and perpendicular to the arm axis (Figure 6). Fine radial striations were also found in the center of the depression (Figure 6). These striations were very similar to those of *Ophiura kinbergi* in the aquarium




Figure 5. Producing process of striations on aluminiferous abrasive by *Astropecten scoparius* (specimen from Site C) observed in the plastic tray. **A**, successive images when starting to move again from its resting posture (time interval 50 sec). Three arm depressions marked by ovals were produced after the asteroid moved. **B**, enlarged view of the striations and sketch of a part of it. Arrows show the direction in which the asteroid moved. Abbreviations: Mo, mound of substratum; Std, deep striation; Stw, wide striation. Scale bars equal 1 cm.




Figure 6. Striations in a star-shaped trace produced by the ophiuroid *Amphiophiura penichra* on the deep-sea floor (Site F). A star-shaped trace on the left photograph is sketched at right. Arrows show the estimated direction in which the ophiuroid moved. Striations are observed in arm depressions and in a central depression (marked by an oval). The bottom sediment is silt. Abbreviations: Stb, striation by basal-arm tube-feet; Sto, striation by oral tube-feet. Scale bars equal 1 cm. Modified from Figure 5B in Ishida *et al.* (2004).



Figure 7. Striations in a star-shaped trace produced by an astropectinid asteroid on the deep-sea floor (Site G). A star-shaped trace on the left photograph is sketched at right. Arrows show the estimated direction in which the asteroid moved. The bottom sediment is silt. Abbreviations: Std, deep striation; Stw, wide striation. Scale bars equal 5 cm. Modified from Figure 5G in Ishida *et al.* (2013).

observation.

Striations of an astropectinid asteroid were found in the four arm depressions, and they were perpendicular to the arm axis (Figure 7). In each arm depression, in one

half the striations were wide, regular, and in contact with each other, while in the other half they were deep and irregular (Figure 7). The striations are similar to those of *Astropecten scoparius* in the aquarium observation.




Figure 8. Striations of trace fossils of *Asteriacites lumbricalis*. Traces on the left photographs are sketched at right. **A**, Triassic Hiraiso Formation, mudstone (NMNS PA 18300, modified from Ishida *et al.*, 2004, fig. 3); **B**, Jurassic Hienheim Formation, fine grained limestone (BMMS); **C**, Triassic Lazurnaya Bay Formation, medium sandstone (NMNS PA 18301); **D**, Cretaceous Himenoura Group, fine sandstone (NMNS PA 18302, synthetic resin cast of this material). Arrows show the direction of travel of the ophiuroids that produced these traces. Striations are marked by ovals. Abbreviations: Stb, striation by basal-arm tube-feet; Sto, striation by oral tube-feet. Scale bars equal 5 mm (A, B, C) and 1 cm (D).

Trace fossils

The trace fossils of *Asteriacites lumbricalis* from the Lower Triassic Hiraiso Formation (Figure 8A) and the lower Tithonian (Upper Jurassic) Hienheim Formation (Figure 8B) had fine, well spaced striations, parallel to each other, and perpendicular to the arm axis in four arm depressions, and faint, radial, fine striations in the center depression. Star-shaped trace fossils from the Triassic Lazurnaya Bay Formation (Figure 8C) and the Cretaceous Himenoura Group (Figure 8D) were identified also as *A. lumbricalis* by having a small form, a central disc and relatively narrow arms. They also had fine, well spaced striations, parallel to each other and perpendicular to the arm axis, and radially fine striations. The striations of these fossils were similar to those of the resting trace of living ophiuroids.

The trace fossils of *Asteriacites quinquefolius* from the Upper Jurassic Hienheim Formation (Figure 9) had both wide, regular and contact striations and deep and irregu-

lar striations in the arm depressions. The striations were similar to those of the resting trace of living asteroids in shape, direction and location.

Discussion

The detailed morphology of striations and their producing process were clarified in this study. Similar striations to those of this study were found also in the trace fossils reported in previous papers. Striations similar to those of resting traces of extant ophiuroids were observed in *Asteriacites lumbricalis*, from the Lower Devonian of the Union of South Africa (Fourie, 2009, figure on p. 33), from the upper Carboniferous of U. S. A. (Hakes, 1976, pl. 1, fig. 1c), from the Lower Triassic of U. S. A. (Wilson and Rigby, 2000, fig. 3) and from the Lower Jurassic of Germany (Seilacher, 1953, fig. 1). Moreover, *Asteriacites* sp. from the Carboniferous of U. S. A. (Hakes, 1976, *loc. cit.*) and an unidentified star-shaped trace fossil from the




Figure 9. Striations of trace fossils of *Asteriacites quinquefolius*, Jurassic Hienheim Formation, fine-grained limestone (BMMS). A trace on the left photograph is sketched at right. Arrows show the direction of travel of the asteroid that produced the resting trace. Striations are marked by ovals. Abbreviations: Std, deep striation; Stw, wide striation. Scale bars equal 1 cm. Modified from Ishida *et al.* (2013, fig. 3).

Upper Jurassic of Germany (Röper and Rothgaenger, 1998, fig. 135) were probably *A. lumbricalis* due to their small form with slender arm depressions, a central sub-circular depression and the strikingly similar striations. Striations similar to those of resting traces of extant asteroids were also found in *A. quinquefolius* from the Middle Jurassic of Switzerland (Seilacher, 1953, table 10, fig. 2). The star-shaped trace fossil from the Upper Jurassic of Germany (Röper and Rothgaenger, 1998, fig. 146) was probably *A. quinquefolius* due to its large form with the wide arm depressions and the typical striations.

Seilacher (1953) suggested that the striations of *Asteriacites lumbricalis* and *A. quinquefolius* were marked by tube-feet of ophiuroids and asteroids, respectively, by comparing the shape of striations remaining on the substrate in aquarium observations, but he did not observe the tube-feet movement. In his observations, striations made by ophiuroids were parallel to each other in the arm depressions and radial in the central depression, whereas striations made by asteroids were irregular in the arm depressions, and the morphological details of the striations were not made clear (Seilacher, 1953). In this study, the movement of tube-feet to produce the striations and the morphology of the striations produced by the movement were observed in detail using extant animals in an aquarium for the first time. Additionally, striations were observed on *in situ* photographs of star-shaped traces on the sea floor. Based on these observations, we showed that the morphology of striations was completely different between *A. lumbricalis* and *A. quinquefolius*, and that these different striations were produced by different producers, ophiuroids and asteroids, by the different morphology and behavior of the tube-feet.

When ophiuroids buried themselves, not only basal-arm tube-feet but also oral tube-feet were used for sweeping away the substratum under their body (Figure 10A, 1). This behavior of oral tube-feet digging in the substratum has never been reported. The oral tube-feet behavior when burrowing into the substratum was recently observed also in a possibly infaunal species of Ophiomycetidae (Masanori Okanishi *et al.*, unpublished data).

The striations in the resting traces of ophiuroids and asteroids found on the deep-sea floor agreed with the striations of those in aquariums morphologically (Figures 6, 7), but some differences were recognized. The striations made by ophiuroids on the silty deep-sea floor (Figure 6) were fainter than those in the aquarium (Figure 3) and the asteroids' deep irregular striations in the front half of arm depressions on the silty deep-sea floor (Figure 7) were deeper than those in the aquarium (Figure 5), probably because the substratum is finer on the deep-sea floor than in the aquarium. Similar striation morphology of ophiuroids and asteroids were observed both in the aquarium and on the sea floor, thus strengthening the identification of the producers of the trace fossil *Asteriacites*.

In previous studies, two ichnospecies of *Asteriacites* have been identified only by the shape of traces and the direction and density of the striations in the arm depressions (e.g. Seilacher, 1953; Hakes, 1976; Twitchett and Wignall, 1996; Wilson and Rigby, 2000; Fourie, 2009; Bernardi and Avanzini, 2011; Ishida *et al.*, 2004, 2013). Based on the detailed observations of morphology of the striations in this study, the two ichnospecies of *Asteriacites* were more clearly distinguished. The striations of *Asteriacites lumbricalis* (Figure 8) were fine, parallel to each other, perpendicular to the arm axis, well spaced




Figure 10. Horizontal and vertical schematic diagrams of the process of production of the striations of *Asteriacites lumbricalis* by ophiuroids in (A) and of *A. quinquefolius* by asteroids in (B). Thin and thick arrows show the movement of tube-feet and the direction of travel of the animals, respectively. Abbreviations: Ar, arm; Di, disc; Mo, mound of substratum; Ot1, first oral tube-foot; Ot2, second oral tube-foot; Stb, striation by basal-arm tube-feet; Std, deep striation; Sto, striation by oral tube-feet; Stw, wide striation; Su, substratum; Tu, tube-foot. Modified from Ishida *et al.* (2004, fig. 4) and Ishida *et al.* (2013, fig. 4).

in the four arm depressions, and fine, radial in the center depression, while those of *A. quinquefolius* (Figure 9) were deep and irregular in the front half of each arm depression, and wide, regular, and in contact in the other half. Furthermore, the analyses of the producing process of the striations by ophiuroids and asteroids in aquariums and *in situ* (Figure 10) clarified that the striations of *A. lumbricalis* and *A. quinquefolius* were produced by the movement of basal-arm tube-feet and oral tube-feet of fossil ophiuroids and of the tube-feet of fossil asteroids, respectively. Consequently, this study unequivocally corroborated that *A. lumbricalis* was, indeed, produced by ophiuroids, and *A. quinquefolius* by asteroids.

Acknowledgements

We would like to thank Hisanori Kohtsuka of the University of Tokyo, Yoichi Kogure of the Japan Sea National Fisheries Research Institute, Mamoru Yamaguchi of Ochanomizu University, Yoshiaki Kai of Kyoto University, Suguru Ohta emeritus professor of the University of Tokyo, and the staff and crew of the R. V. *Tansei-maru*, the T. S. *Toyoshio-maru*, the R. V. *Ryokuyo-maru*, and

the R. V. *Mizuho-maru* for their kind help in sampling. Many thanks go to Shiro Hasegawa, emeritus professor of Kumamoto University for his comments on the trace fossils. Thanks are due to Takumu Kato for his kind help in analysis of the movies. Thanks are extended to David L. Meyer of the University of Cincinnati, Frederick H. C. Hotchkiss of the Marine and Paleobiological Research Institute, Vineyard Haven, Massachusetts, U. S. A., Ben Thuy of the National Museum of Natural History, Luxembourg and an anonymous reviewer for greatly improving an earlier draft of this manuscript. A part of this work was financially supported by JSPS KAKENHI Grant Numbers 25540226 to Fujita and 16K05593 to Komatsu.

References

- Bernardi, M. and Avanzini, M., 2011: *Asteriacites lumbricalis* from the Anisian (Middle Triassic) of Vallarsa (Southern Trentino, NE Italy). *Studi Tridentino di Scienze Naturali*, vol. 88, p. 187–190.
- Chamberlain, C. K., 1971: Morphology and ethology of trace fossils from the Ouachita Mountains, southeast Oklahoma. *Journal of Paleontology*, vol. 45, p. 212–246.
- Clark, H. L., 1908: Some Japanese and East Indian echinoderms. *Bul-*

- letin of the Museum of Comparative Zoology*, vol. 51, p. 279–311.
- Clark, H. L., 1911: North Pacific ophiurans in the collection of the United States National Museum. *Smithsonian Institution United States National Museum Bulletin*, vol. 75, p. 1–302.
- Fourie, P. H., 2009: Unique asterozoan trace fossils of the Gydo Formation (Bokkeveld Group, Cape Supergroup) near Nieuwoudtville, Northern Cape Province. In, Beukes, N. and Rajesh, H. M. eds., *The Annual Report of the Paleoproterozoic Mineralization Research (PPM) Group*, p. 33. Department of Geology, University of Johannesburg, Johannesburg.
- Hakes, W. G., 1976: Trace fossils and depositional environment of four clastic units, Upper Pennsylvanian megacyclothems, north-east Kansas. *University of Kansas Paleontological Contributions*, article 63, p. 1–46.
- Ishida, Y., Fujita, T. and Kamada, K., 2004: Ophiuroid trace fossils in the Triassic of Japan compared to the resting behavior of extant brittle stars. In, Heinzeller, T. and Nebelsick, J. H. eds., *Echinoderms: München*, p. 433–438. Taylor and Francis Group, London.
- Ishida, Y., Röper, M. and Fujita, T., 2013: Comparison of asteroid and ophiuroid trace fossils in the Jurassic of Germany with resting behavior of extant asteroids and ophiuroids. In, Johnson, C. ed., *Echinoderms in a Changing World*, p. 87–94. Taylor and Francis Group, London.
- Kamada, K. and Takizawa, F., 1992: *Geology of the Osu District. With Geological Sheet Map at 1:50,000*, 69 p. Geological Survey of Japan, Tsukuba. (in Japanese with English abstract)
- Kojo, Y., Komatsu, T., Iwamoto, T., Takashima, R., Takahashi, O. and Nishi, H., 2011: Stratigraphy and detailed age of the Upper Cretaceous Himenoura Group in the eastern part of Amakusa-Kamishima Island, Kumamoto, Japan. *Journal of the Geological Society of Japan*, vol. 117, p. 398–416. (in Japanese with English abstract)
- Komatsu, T., Ono, M., Naruse, H. and Kumagae, T., 2008: Upper Cretaceous depositional environments and bivalve assemblages of Far-East Asia: the Himenoura Group, Kyushu, Japan. *Cretaceous Research*, vol. 29, p. 489–508.
- Ljungman, A. V., 1866: Om några nya arter af Ophiurider. *Öfversigt af Kongliga Vetenskaps-Akademiens Förhandlingar*, vol. 23, p. 163–166.
- Mángano, M. G., Buatois, L. A., West, R. R. and Maples, C. G., 1999: The origin and paleoecologic significance of the trace fossil *Asteriacites* in the Pennsylvanian of Kansas and Missouri. *Lethaia*, vol. 32, p. 17–30.
- Mikuláš, R., 1990: The ophiuroid *Taeniaster* as a tracemaker of *Asteriacites*, Ordovician of Czechoslovakia. *Ichnos*, vol. 1, p. 133–137.
- Müller, J. H. and Troschel, F. H., 1842: *System der Asteriden*, p. 1–134. Vieweg, Braunschweig.
- Patel, S. J., Bhatt, N. Y. and Desai, B. G., 2008: *Asteriacites quinquefolius*—asteroid trace maker from the Bhuj Formation (Lower Cretaceous) of the Mainland Kachchh, western India. *Journal of the Geological Society of India*, vol. 71, p. 129–132.
- Quenstedt, F. A., 1876: *Petrefaktenkunde Deutschlands. 4. Die Asteriden und Encriniden nebst Cysti- und Blastoideen*, p. 1–742. Fuess, Leipzig.
- Röper, M. and Rothgaenger, M., 1995: Eine neue Fossilagerstätte in den Ostbayerischen Oberjura-Plattenkalken bei Brunn/Oberpfalz. *Jahresbericht 1994 und Mitteilungen der Freunde der Bayerischen Staatssammlung für Paläontologie und Historische Geologie München e. V.*, 23, p. 32–46.
- Röper, M. and Rothgaenger, M., 1998: *Die Plattenkalke von Hienheim*, 110 p. Eichendorf Verlag, Eichendorf.
- Schlotheim, E. F. von, 1820: *Die Petrefaktenkunde auf ihrem jetzigen Standpunkte durch die Beschreibung seiner Sammlung Versteinerter und Fossiler Überreste des Thier- und Pflanzenreichs der Vorwelt Erläutert*, 437 p. Becher, Gotha.
- Seilacher, A., 1953: Studien zur Palichnologie II. Die fossilen Ruhespuren (Cubichnia). *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen*, Band 98, p. 87–124.
- Seilacher, A., 2007: *Trace Fossil Analysis*, 226 p. Springer, Berlin.
- Shigeta, Y., Zakharov, Y. D., Maeda, H. and Popov, A. M. eds., 2009: The Lower Triassic System in the Abrek Bay area, South Primorye, Russia. *National Museum of Nature and Science Monographs*, no. 38, p. 1–218.
- Twitchett, R. J. and Wignall, P. B., 1996: Trace fossils and the aftermath of the Permo-Triassic mass extinction: evidence from northern Italy. *Palaeogeography, Palaeoclimatology, Palaeoecology*, vol. 124, p. 137–151.
- West, R. R. and Ward, E. L., 1990: *Asteriacites lumbricalis* and a protasterid ophiuroid. In, Boucot, A. J. ed., *Evolutionary Paleobiology of Behavior and Coevolution*, p. 321–327. Elsevier, Amsterdam.
- Wilson, M. A. and Rigby, J., 2000: *Asteriacites lumbricalis* von Schlotheim 1820: ophiuroid trace fossils from the Lower Triassic Thaynes Formation, central Utah. *Ichnos*, vol. 7, p. 43–49.